

Lecture 5

September 6, 2018

Last class

- ▶ Recall the least squares estimator defined by $\hat{\beta} = (X^T X)^{-1} X^T Y$.

Last class

- ▶ Recall the least squares estimator defined by $\hat{\beta} = (X^T X)^{-1} X^T Y$.
- ▶ This estimates β in the linear model $Y = X\beta + e$ where $\mathbb{E}(e|X) = 0$.

Last class

- ▶ Recall the least squares estimator defined by $\hat{\beta} = (X^T X)^{-1} X^T Y$.
- ▶ This estimates β in the linear model $Y = X\beta + e$ where $\mathbb{E}(e|X) = 0$.
- ▶ We saw in the last class that $\hat{\beta}$ is linear in Y and conditionally unbiased i.e $\mathbb{E}(\hat{\beta}) = \beta$.

Last class

- ▶ Recall the least squares estimator defined by $\hat{\beta} = (X^T X)^{-1} X^T Y$.
- ▶ This estimates β in the linear model $Y = X\beta + e$ where $\mathbb{E}(e|X) = 0$.
- ▶ We saw in the last class that $\hat{\beta}$ is linear in Y and conditionally unbiased i.e $\mathbb{E}(\hat{\beta}) = \beta$.
- ▶ Under the additional assumption of homoskedasticity i.e., $\text{Cov}(e|X) = \sigma^2 I_n$, we also calculated the covariance matrix of $\hat{\beta}$ to be $\text{Cov}(\hat{\beta}) = \sigma^2 (X^T X)^{-1}$.

Gauss-Markov Theorem and Proof

- ▶ The Gauss-Markov Theorem states that $\hat{\beta}$ is BLUE (Best Linear Unbiased Estimator) under homoskedasticity.

Gauss-Markov Theorem and Proof

Four key words,
cannot delete
anyone

- ▶ The Gauss-Markov Theorem states that $\hat{\beta}$ is BLUE (Best Linear Unbiased Estimator) under homoskedasticity.
- ▶ This means that $\hat{\beta}$ is the “best” estimator among all linear and unbiased estimators of β .

Gauss-Markov Theorem and Proof

- ▶ The Gauss-Markov Theorem states that $\hat{\beta}$ is BLUE (Best Linear Unbiased Estimator) under homoskedasticity.
- ▶ This means that $\hat{\beta}$ is the “best” estimator among all linear and unbiased estimators of β .
- ▶ Here “best” is in terms of variance. This implies that $\hat{\beta}_i$ has the smallest variance among all linear and unbiased estimators of β_i for every i .

Gauss-Markov Theorem and Proof

- ▶ The Gauss-Markov Theorem states that $\hat{\beta}$ is BLUE (Best Linear Unbiased Estimator) under homoskedasticity.
- ▶ This means that $\hat{\beta}$ is the “best” estimator among all linear and unbiased estimators of β .
- ▶ Here “best” is in terms of variance. This implies that $\hat{\beta}_i$ has the smallest variance among all linear and unbiased estimators of β_i for every i .
- ▶ The Gauss-Markov theorem is actually quite simple to prove. Suppose $\tilde{\beta} = AY$ is any other linear unbiased estimator for β .

Gauss-Markov Theorem and Proof

- ▶ We get back the least squares estimator if $A = (X^T X)^{-1} X^T$ so let us assume that $B := A - (X^T X)^{-1} X^T \neq 0$.

Gauss-Markov Theorem and Proof

- ▶ We get back the least squares estimator if $A = (X^T X)^{-1} X^T$ so let us assume that $B := A - (X^T X)^{-1} X^T \neq 0$.
- ▶ Now the expectation of $\tilde{\beta}$ equals $\beta + BX\beta$.

Gauss-Markov Theorem and Proof

- ▶ We get back the least squares estimator if $A = (X^T X)^{-1} X^T$ so let us assume that $B := A - (X^T X)^{-1} X^T \neq 0$.
- ▶ Now the expectation of $\tilde{\beta}$ equals $\beta + BX\beta$.
- ▶ Because $\tilde{\beta}$ is unbiased, this must mean that $BX = 0$.

Gauss-Markov Theorem and Proof

- ▶ We get back the least squares estimator if $A = (X^T X)^{-1} X^T$ so let us assume that $B := A - (X^T X)^{-1} X^T \neq 0$.
- ▶ Now the expectation of $\tilde{\beta}$ equals $\beta + BX\beta$.
- ▶ Because $\tilde{\beta}$ is unbiased, this must mean that $BX = 0$.
- ▶ The covariance matrix of $\tilde{\beta}$ is

Gauss-Markov Theorem and Proof

- ▶ We get back the least squares estimator if $A = (X^T X)^{-1} X^T$ so let us assume that $B := A - (X^T X)^{-1} X^T \neq 0$.
- ▶ Now the expectation of $\tilde{\beta}$ equals $\beta + BX\beta$.
- ▶ Because $\tilde{\beta}$ is unbiased, this must mean that $BX = 0$.
- ▶ The covariance matrix of $\tilde{\beta}$ is

$$\text{Cov}(\tilde{\beta}) = \text{Cov}(AY)$$

Gauss-Markov Theorem and Proof

- ▶ We get back the least squares estimator if $A = (X^T X)^{-1} X^T$ so let us assume that $B := A - (X^T X)^{-1} X^T \neq 0$.
- ▶ Now the expectation of $\tilde{\beta}$ equals $\beta + BX\beta$.
- ▶ Because $\tilde{\beta}$ is unbiased, this must mean that $BX = 0$.
- ▶ The covariance matrix of $\tilde{\beta}$ is

$$\begin{aligned} \text{Cov}(\tilde{\beta}) &= \text{Cov}(AY) \\ &= ACov(Y)A^T \end{aligned}$$

Gauss-Markov Theorem and Proof

- ▶ We get back the least squares estimator if $A = (X^T X)^{-1} X^T$ so let us assume that $B := A - (X^T X)^{-1} X^T \neq 0$.
- ▶ Now the expectation of $\tilde{\beta}$ equals $\beta + BX\beta$.
- ▶ Because $\tilde{\beta}$ is unbiased, this must mean that $BX = 0$.
- ▶ The covariance matrix of $\tilde{\beta}$ is

$$\begin{aligned} \text{Cov}(\tilde{\beta}) &= \text{Cov}(AY) \\ &= A \text{Cov}(Y) A^T \\ &= [B + (X^T X)^{-1} X^T] [\sigma^2 I_n] [B + (X^T X)^{-1} X^T]^T \end{aligned}$$

Gauss-Markov Theorem and Proof

- ▶ We get back the least squares estimator if $A = (X^T X)^{-1} X^T$ so let us assume that $B := A - (X^T X)^{-1} X^T \neq 0$.
- ▶ Now the expectation of $\tilde{\beta}$ equals $\beta + BX\beta$.
- ▶ Because $\tilde{\beta}$ is unbiased, this must mean that $BX = 0$.
- ▶ The covariance matrix of $\tilde{\beta}$ is

$$\begin{aligned} \text{Cov}(\tilde{\beta}) &= \text{Cov}(AY) \\ &= A\text{Cov}(Y)A^T \\ &= [B + (X^T X)^{-1} X^T][\sigma^2 I_n][B + (X^T X)^{-1} X^T]^T \\ &= \sigma^2 [B + (X^T X)^{-1} X^T][B^T + X(X^T X)^{-1}] \end{aligned}$$

Gauss-Markov Theorem and Proof

- ▶ We get back the least squares estimator if $A = (X^T X)^{-1} X^T$ so let us assume that $B := A - (X^T X)^{-1} X^T \neq 0$.
- ▶ Now the expectation of $\tilde{\beta}$ equals $\beta + BX\beta$.
- ▶ Because $\tilde{\beta}$ is unbiased, this must mean that $BX = 0$.
- ▶ The covariance matrix of $\tilde{\beta}$ is

$$\begin{aligned} \text{Cov}(\tilde{\beta}) &= \text{Cov}(AY) \\ &= ACov(Y)A^T \\ &= [B + (X^T X)^{-1} X^T][\sigma^2 I_n][B + (X^T X)^{-1} X^T]^T \\ &= \sigma^2 [B + (X^T X)^{-1} X^T][B^T + X(X^T X)^{-1}] \\ &= \sigma^2 [(X^T X)^{-1} + BX(X^T X)^{-1} + (X^T X)^{-1} X^T B^T + BB^T] \end{aligned}$$

Gauss-Markov Theorem and Proof

- ▶ Because $BX = 0$, we deduce from the previous expression that

$$\text{Cov}(\tilde{\beta}) = \sigma^2(X^T X)^{-1} + \sigma^2 B B^T.$$

Gauss-Markov Theorem and Proof

- ▶ Because $BX = 0$, we deduce from the previous expression that

$$\text{Cov}(\tilde{\beta}) = \sigma^2(X^T X)^{-1} + \sigma^2 B B^T.$$

- ▶ Recall that $\sigma^2(X^T X)^{-1}$ is the covariance matrix of the least squares estimator $\hat{\beta}$.

Gauss-Markov Theorem and Proof

- ▶ Because $BX = 0$, we deduce from the previous expression that

$$\text{Cov}(\tilde{\beta}) = \sigma^2(X^T X)^{-1} + \sigma^2 B B^T.$$

- ▶ Recall that $\sigma^2(X^T X)^{-1}$ is the covariance matrix of the least squares estimator $\hat{\beta}$. We thus have

$$\text{Cov}(\tilde{\beta}) = \text{Cov}(\hat{\beta}) + \sigma^2 B B^T.$$

Gauss-Markov Theorem and Proof

- ▶ Because $BX = 0$, we deduce from the previous expression that

$$\text{Cov}(\tilde{\beta}) = \sigma^2(X^T X)^{-1} + \sigma^2 B B^T.$$

- ▶ Recall that $\sigma^2(X^T X)^{-1}$ is the covariance matrix of the least squares estimator $\hat{\beta}$. We thus have

$$\text{Cov}(\tilde{\beta}) = \text{Cov}(\hat{\beta}) + \sigma^2 B B^T.$$

- ▶ It is important to note that the Gauss-Markov theorem requires the assumption of homoskedasticity.

The case when $X^T X$ is not invertible

The case when $X^T X$ is not invertible

- Note first that $X^T X$ is not invertible if and only if there exists a column of X that can be written as the linear combination of all the other columns.

The case when $X^T X$ is not invertible

- ▶ Note first that $X^T X$ is not invertible if and only if there exists a column of X that can be written as the linear combination of all the other columns.
- ▶ To see this, just note that $X^T X$ is non-invertible if and only if there exists a non-zero vector w for which $X^T X w = 0$ which is equivalent to $X w = 0$.

The case when $X^T X$ is not invertible

- ▶ Note first that $X^T X$ is not invertible if and only if there exists a column of X that can be written as the linear combination of all the other columns.
- ▶ To see this, just note that $X^T X$ is non-invertible if and only if there exists a non-zero vector w for which $X^T X w = 0$ which is equivalent to $X w = 0$.
- ▶ This is because

$$X^T X w = 0 \Rightarrow w^T X^T X w = 0 \Rightarrow \|X w\|_2^2 = 0 \rightarrow X w = 0.$$

The case when $X^T X$ is not invertible

- ▶ Note first that $X^T X$ is not invertible if and only if there exists a column of X that can be written as the linear combination of all the other columns.
- ▶ To see this, just note that $X^T X$ is non-invertible if and only if there exists a non-zero vector w for which $X^T X w = 0$ which is equivalent to $X w = 0$.
- ▶ This is because

$$X^T X w = 0 \Rightarrow w^T X^T X w = 0 \Rightarrow \|X w\|_2^2 = 0 \rightarrow X w = 0.$$

- ▶ In the case when $X^T X$ is not invertible, we obviously cannot solve the normal equations $X^T X b = X^T Y$ as $b = (X^T X)^{-1} X^T Y$.

The case when $X^T X$ is not invertible

- ▶ Note first that $X^T X$ is not invertible if and only if there exists a column of X that can be written as the linear combination of all the other columns.
- ▶ To see this, just note that $X^T X$ is non-invertible if and only if there exists a non-zero vector w for which $X^T X w = 0$ which is equivalent to $X w = 0$.
- ▶ This is because

$$X^T X w = 0 \Rightarrow w^T X^T X w = 0 \Rightarrow \|X w\|_2^2 = 0 \rightarrow X w = 0.$$

- ▶ In the case when $X^T X$ is not invertible, we obviously cannot solve the normal equations $X^T X b = X^T Y$ as $b = (X^T X)^{-1} X^T Y$.
- ▶ So how do we estimate β here?

The case when $X^T X$ is not invertible

- ▶ Note first that $X^T X$ is not invertible if and only if there exists a column of X that can be written as the linear combination of all the other columns.
- ▶ To see this, just note that $X^T X$ is non-invertible if and only if there exists a non-zero vector w for which $X^T X w = 0$ which is equivalent to $X w = 0$.
- ▶ This is because

$$X^T X w = 0 \Rightarrow w^T X^T X w = 0 \Rightarrow \|X w\|_2^2 = 0 \rightarrow X w = 0.$$

- ▶ In the case when $X^T X$ is not invertible, we obviously cannot solve the normal equations $X^T X b = X^T Y$ as $b = (X^T X)^{-1} X^T Y$.
- ▶ So how do we estimate β here?
- ▶ It turns out that even when $X^T X$ is non-invertible, the normal equations have a solution.

The case when $X^T X$ is not invertible

- ▶ This is because the column space of $X^T X$ (denoted by $\mathcal{C}(X^T X)$) always equals the column space of X^T (denoted by $\mathcal{C}(X^T)$).

The case when $X^T X$ is not invertible

- ▶ This is because the column space of $X^T X$ (denoted by $\mathcal{C}(X^T X)$) always equals the column space of X^T (denoted by $\mathcal{C}(X^T)$).
- ▶ To see why $\mathcal{C}(X^T X) = \mathcal{C}(X^T)$, note that it is enough to prove that $\mathcal{C}(X^T X)^\perp = \mathcal{C}(X^T)^\perp$ where $\mathcal{C}(X^T X)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T X)$ (similarly $\mathcal{C}(X^T)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T)$).

The case when $X^T X$ is not invertible

- ▶ This is because the column space of $X^T X$ (denoted by $\mathcal{C}(X^T X)$) always equals the column space of X^T (denoted by $\mathcal{C}(X^T)$).
- ▶ To see why $\mathcal{C}(X^T X) = \mathcal{C}(X^T)$, note that it is enough to prove that $\mathcal{C}(X^T X)^\perp = \mathcal{C}(X^T)^\perp$ where $\mathcal{C}(X^T X)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T X)$ (similarly $\mathcal{C}(X^T)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T)$).
- ▶ Now a vector $u \in \mathcal{C}(X^T X)^\perp$ if and only if $u^T X^T X = 0$

The case when $X^T X$ is not invertible

- ▶ This is because the column space of $X^T X$ (denoted by $\mathcal{C}(X^T X)$) always equals the column space of X^T (denoted by $\mathcal{C}(X^T)$).
- ▶ To see why $\mathcal{C}(X^T X) = \mathcal{C}(X^T)$, note that it is enough to prove that $\mathcal{C}(X^T X)^\perp = \mathcal{C}(X^T)^\perp$ where $\mathcal{C}(X^T X)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T X)$ (similarly $\mathcal{C}(X^T)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T)$).
- ▶ Now a vector $u \in \mathcal{C}(X^T X)^\perp$ if and only if $u^T X^T X = 0$

The case when $X^T X$ is not invertible

- ▶ This is because the column space of $X^T X$ (denoted by $\mathcal{C}(X^T X)$) always equals the column space of X^T (denoted by $\mathcal{C}(X^T)$).
- ▶ To see why $\mathcal{C}(X^T X) = \mathcal{C}(X^T)$, note that it is enough to prove that $\mathcal{C}(X^T X)^\perp = \mathcal{C}(X^T)^\perp$ where $\mathcal{C}(X^T X)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T X)$ (similarly $\mathcal{C}(X^T)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T)$).
- ▶ Now a vector $u \in \mathcal{C}(X^T X)^\perp$ if and only if $u^T X^T X = 0$ which is the same as $u^T X^T = 0$ i.e., $u \in \mathcal{C}(X^T)^\perp$.

The case when $X^T X$ is not invertible

- ▶ This is because the column space of $X^T X$ (denoted by $\mathcal{C}(X^T X)$) always equals the column space of X^T (denoted by $\mathcal{C}(X^T)$).
- ▶ To see why $\mathcal{C}(X^T X) = \mathcal{C}(X^T)$, note that it is enough to prove that $\mathcal{C}(X^T X)^\perp = \mathcal{C}(X^T)^\perp$ where $\mathcal{C}(X^T X)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T X)$ (similarly $\mathcal{C}(X^T)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T)$).
- ▶ Now a vector $u \in \mathcal{C}(X^T X)^\perp$ if and only if $u^T X^T X = 0$ which is the same as $u^T X^T = 0$ i.e., $u \in \mathcal{C}(X^T)^\perp$.
- ▶ Because $\mathcal{C}(X^T X) = \mathcal{C}(X^T)$, the normal equations $X^T X b = X^T Y$ always have a solution

The case when $X^T X$ is not invertible

- ▶ This is because the column space of $X^T X$ (denoted by $\mathcal{C}(X^T X)$) always equals the column space of X^T (denoted by $\mathcal{C}(X^T)$).
- ▶ To see why $\mathcal{C}(X^T X) = \mathcal{C}(X^T)$, note that it is enough to prove that $\mathcal{C}(X^T X)^\perp = \mathcal{C}(X^T)^\perp$ where $\mathcal{C}(X^T X)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T X)$ (similarly $\mathcal{C}(X^T)^\perp$ denotes the orthogonal complement of $\mathcal{C}(X^T)$).
- ▶ Now a vector $u \in \mathcal{C}(X^T X)^\perp$ if and only if $u^T X^T X = 0$ which is the same as $u^T X^T = 0$ i.e., $u \in \mathcal{C}(X^T)^\perp$.
- ▶ Because $\mathcal{C}(X^T X) = \mathcal{C}(X^T)$, the normal equations $X^T X b = X^T Y$ always have a solution (because $X^T Y \in \mathcal{C}(X^T) = \mathcal{C}(X^T X)$, we can always write $X^T Y$ as $X^T X b$ for some b).

The case when $X^T X$ is not invertible

- ▶ However, when $X^T X$ is non-invertible, the normal equations will have infinitely many solutions.

The case when $X^T X$ is not invertible

- ▶ However, when $X^T X$ is non-invertible, the normal equations will have infinitely many solutions. In this case, we shall call any solution of $X^T X b = X^T Y$ a least squares estimator of β .

The case when $X^T X$ is not invertible

- ▶ However, when $X^T X$ is non-invertible, the normal equations will have infinitely many solutions. In this case, we shall call any solution of $X^T X b = X^T Y$ a least squares estimator of β .
- ▶ When $X^T X$ is non-invertible, there will be infinitely many least squares estimators for β .

The case when $X^T X$ is not invertible

- ▶ However, when $X^T X$ is non-invertible, the normal equations will have infinitely many solutions. In this case, we shall call any solution of $X^T X b = X^T Y$ a least squares estimator of β .
- ▶ When $X^T X$ is non-invertible, there will be infinitely many least squares estimators for β .
- ▶ It might seem strange to have infinitely many least squares estimators. It turns out however that when $X^T X$ is non-invertible, estimation of β is not a well-defined problem.

The case when $X^T X$ is not invertible

- ▶ However, when $X^T X$ is non-invertible, the normal equations will have infinitely many solutions. In this case, we shall call any solution of $X^T X b = X^T Y$ a least squares estimator of β .
- ▶ When $X^T X$ is non-invertible, there will be infinitely many least squares estimators for β .
- ▶ It might seem strange to have infinitely many least squares estimators. It turns out however that when $X^T X$ is non-invertible, estimation of β is not a well-defined problem.
- ▶ To see this, note first that if X_{i*} denotes the i th row of X , then the i th observation Y_i can be written as $Y_i = X_{i*}\beta + e_i$. We therefore only have data on the linear combinations $X_{1*}\beta, \dots, X_{n*}\beta$.

The case when $X^T X$ is not invertible

- ▶ However, when $X^T X$ is non-invertible, the normal equations will have infinitely many solutions. In this case, we shall call any solution of $X^T X b = X^T Y$ a least squares estimator of β .
- ▶ When $X^T X$ is non-invertible, there will be infinitely many least squares estimators for β .
- ▶ It might seem strange to have infinitely many least squares estimators. It turns out however that when $X^T X$ is non-invertible, estimation of β is not a well-defined problem.
- ▶ To see this, note first that if X_{i*} denotes the i th row of X , then the i th observation Y_i can be written as $Y_i = X_{i*} \beta + e_i$. We therefore only have data on the linear combinations $X_{1*} \beta, \dots, X_{n*} \beta$.
- ▶ Therefore, we can only estimate a linear function $\lambda^T \beta$ if the vector λ can be written as a linear combination of the rows of X .

Estimability

- ▶ The previous discussion motivates the following definition of estimability:

Estimability

- ▶ The previous discussion motivates the following definition of estimability:

Definition

A linear function $\lambda^T \beta$ of β is said to be estimable if the vector λ is a linear combination of the rows of X .

Estimability

- ▶ The previous discussion motivates the following definition of estimability:

Definition

A linear function $\lambda^T \beta$ of β is said to be estimable if the vector λ is a linear combination of the rows of X .

- ▶ Equivalently, we say that $\lambda^T \beta$ is estimable if and only if $\lambda \in \mathcal{C}(X^T)$.

Estimability

- ▶ The previous discussion motivates the following definition of estimability:

Definition

A linear function $\lambda^T \beta$ of β is said to be estimable if the vector λ is a linear combination of the rows of X .

- ▶ Equivalently, we say that $\lambda^T \beta$ is estimable if and only if $\lambda \in \mathcal{C}(X^T)$.
- ▶ Because $\mathcal{C}(X^T) = \mathcal{C}(X^T X)$, estimability of $\lambda^T \beta$ is further equivalent to $\lambda \in \mathcal{C}(X^T X)$.

Estimability

- ▶ The previous discussion motivates the following definition of estimability:

Definition

A linear function $\lambda^T \beta$ of β is said to be estimable if the vector λ is a linear combination of the rows of X .

- ▶ Equivalently, we say that $\lambda^T \beta$ is estimable if and only if $\lambda \in \mathcal{C}(X^T)$.
- ▶ Because $\mathcal{C}(X^T) = \mathcal{C}(X^T X)$, estimability of $\lambda^T \beta$ is further equivalent to $\lambda \in \mathcal{C}(X^T X)$.
- ▶ It therefore only makes sense to estimate $\lambda^T \beta$ if $\lambda \in \mathcal{C}(X^T X)$.

Estimability

- ▶ The previous discussion motivates the following definition of estimability:

Definition

A linear function $\lambda^T \beta$ of β is said to be estimable if the vector λ is a linear combination of the rows of X .

- ▶ Equivalently, we say that $\lambda^T \beta$ is estimable if and only if $\lambda \in \mathcal{C}(X^T)$.
- ▶ Because $\mathcal{C}(X^T) = \mathcal{C}(X^T X)$, estimability of $\lambda^T \beta$ is further equivalent to $\lambda \in \mathcal{C}(X^T X)$.
- ▶ It therefore only makes sense to estimate $\lambda \beta$ if $\lambda \in \mathcal{C}(X^T X)$.
- ▶ If $X^T X$ is invertible, then every $\lambda^T \beta$ is estimable.

Estimability

- ▶ The previous discussion motivates the following definition of estimability:

Definition

A linear function $\lambda^T \beta$ of β is said to be estimable if the vector λ is a linear combination of the rows of X .

- ▶ Equivalently, we say that $\lambda^T \beta$ is estimable if and only if $\lambda \in \mathcal{C}(X^T)$.
- ▶ Because $\mathcal{C}(X^T) = \mathcal{C}(X^T X)$, estimability of $\lambda^T \beta$ is further equivalent to $\lambda \in \mathcal{C}(X^T X)$.
- ▶ It therefore only makes sense to estimate $\lambda \beta$ if $\lambda \in \mathcal{C}(X^T X)$.
- ▶ If $X^T X$ is invertible, then every $\lambda^T \beta$ is estimable.
- ▶ If $X^T X$ is non-invertible, then not every $\lambda^T \beta$ is estimable.

Estimability

- **Result:** If $\lambda^T \beta$ is estimable, then $\lambda^T \hat{\beta}_{ls}$ is the same for every solution $\hat{\beta}_{ls}$ of the normal equations. In other words, the least squares estimate of $\lambda^T \beta$ is unique.

Estimability

- ▶ **Result:** If $\lambda^T \beta$ is estimable, then $\lambda^T \hat{\beta}_{ls}$ is the same for every solution $\hat{\beta}_{ls}$ of the normal equations. In other words, the least squares estimate of $\lambda^T \beta$ is unique.
- ▶ **Proof.** Since $\lambda^T \beta$ is estimable, the vector λ lies in the column space of $X^T X$ and hence $\lambda = X^T X u$ for some vector u .

Estimability

- ▶ **Result:** If $\lambda^T \beta$ is estimable, then $\lambda^T \hat{\beta}_{ls}$ is the same for every solution $\hat{\beta}_{ls}$ of the normal equations. In other words, the least squares estimate of $\lambda^T \beta$ is unique.
- ▶ **Proof.** Since $\lambda^T \beta$ is estimable, the vector λ lies in the column space of $X^T X$ and hence $\lambda = X^T X u$ for some vector u . Therefore,

$$\lambda^T \hat{\beta}_{ls} = u^T X^T X \hat{\beta}_{ls}$$

Estimability

- ▶ **Result:** If $\lambda^T \beta$ is estimable, then $\lambda^T \hat{\beta}_{ls}$ is the same for every solution $\hat{\beta}_{ls}$ of the normal equations. In other words, the least squares estimate of $\lambda^T \beta$ is unique.
- ▶ **Proof.** Since $\lambda^T \beta$ is estimable, the vector λ lies in the column space of $X^T X$ and hence $\lambda = X^T X u$ for some vector u . Therefore,

$$\lambda^T \hat{\beta}_{ls} = u^T X^T X \hat{\beta}_{ls} = u^T X^T Y,$$

Estimability

- ▶ **Result:** If $\lambda^T \beta$ is estimable, then $\lambda^T \hat{\beta}_{ls}$ is the same for every solution $\hat{\beta}_{ls}$ of the normal equations. In other words, the least squares estimate of $\lambda^T \beta$ is unique.
- ▶ **Proof.** Since $\lambda^T \beta$ is estimable, the vector λ lies in the column space of $X^T X$ and hence $\lambda = X^T X u$ for some vector u . Therefore,

$$\lambda^T \hat{\beta}_{ls} = u^T X^T X \hat{\beta}_{ls} = u^T X^T Y,$$

where the last equality follows from the fact that $\hat{\beta}_{ls}$ satisfies the normal equations.

Estimability

- ▶ **Result:** If $\lambda^T \beta$ is estimable, then $\lambda^T \hat{\beta}_{ls}$ is the same for every solution $\hat{\beta}_{ls}$ of the normal equations. In other words, the least squares estimate of $\lambda^T \beta$ is unique.
- ▶ **Proof.** Since $\lambda^T \beta$ is estimable, the vector λ lies in the column space of $X^T X$ and hence $\lambda = X^T X u$ for some vector u . Therefore,

$$\lambda^T \hat{\beta}_{ls} = u^T X^T X \hat{\beta}_{ls} = u^T X^T Y,$$

where the last equality follows from the fact that $\hat{\beta}_{ls}$ satisfies the normal equations. Since u only depends on λ , this proves that $\lambda^T \hat{\beta}_{ls}$ does not depend on the particular choice of the solution $\hat{\beta}_{ls}$ of the normal equations.

Summary

The situation when $X^T X$ is non-invertible can be summarized as follows:

Summary

The situation when $X^T X$ is non-invertible can be summarized as follows:

- ▶ There are infinitely many solutions of the normal equations $X^T X b = X^T Y$.

Summary

The situation when $X^T X$ is non-invertible can be summarized as follows:

- ▶ There are infinitely many solutions of the normal equations $X^T X b = X^T Y$.
- ▶ Not every linear combination $\lambda^T \beta$ can be estimated from regression data. We say that $\lambda^T \beta$ is estimable if and only if $\lambda \in \mathcal{C}(X^T) = \mathcal{C}(X^T X)$.

Summary

The situation when $X^T X$ is non-invertible can be summarized as follows:

- ▶ There are infinitely many solutions of the normal equations $X^T X b = X^T Y$.
- ▶ Not every linear combination $\lambda^T \beta$ can be estimated from regression data. We say that $\lambda^T \beta$ is estimable if and only if $\lambda \in \mathcal{C}(X^T) = \mathcal{C}(X^T X)$.
- ▶ When $\lambda^T \beta$ is estimable, the value $\lambda^T \hat{\beta}_{ls}$ is the same for every solution $\hat{\beta}_{ls}$ of the normal equations $X^T X b = X^T Y$.