

Lecture 6

September 10, 2018

Regression Terminology

- ▶ In this lecture and the next, we shall go over some common regression technology:

Regression Terminology

- ▶ In this lecture and the next, we shall go over some common regression technology:
 1. Regression plane

Regression Terminology

- ▶ In this lecture and the next, we shall go over some common regression technology:
 1. Regression plane
 2. Fitted values and the hat matrix

Regression Terminology

- ▶ In this lecture and the next, we shall go over some common regression technology:
 1. Regression plane
 2. Fitted values and the hat matrix
 3. Residuals

Regression Terminology

- ▶ In this lecture and the next, we shall go over some common regression technology:
 1. Regression plane
 2. Fitted values and the hat matrix
 3. Residuals
 4. Leverages

Regression Terminology

- ▶ In this lecture and the next, we shall go over some common regression technology:
 1. Regression plane
 2. Fitted values and the hat matrix
 3. Residuals
 4. Leverages
 5. Residual Sum of Squares (RSS) and R^2

Regression Terminology

- ▶ In this lecture and the next, we shall go over some common regression technology:
 1. Regression plane
 2. Fitted values and the hat matrix
 3. Residuals
 4. Leverages
 5. Residual Sum of Squares (RSS) and R^2
 6. Residual Standard Error

Regression Terminology

- ▶ In this lecture and the next, we shall go over some common regression technology:
 1. Regression plane
 2. Fitted values and the hat matrix
 3. Residuals
 4. Leverages
 5. Residual Sum of Squares (RSS) and R^2
 6. Residual Standard Error
 7. Standardized or Studentized Residuals

The Regression Plane

- ▶ If we get a new subject whose explanatory variable values are x_1, \dots, x_p , then our prediction for its response variable value is

$$y = \hat{\beta}_0 + \hat{\beta}_1 x_{i1} + \dots + \hat{\beta}_p x_{ip}.$$

The Regression Plane

- ▶ If we get a new subject whose explanatory variable values are x_1, \dots, x_p , then our prediction for its response variable value is

$$y = \hat{\beta}_0 + \hat{\beta}_1 x_{i1} + \dots + \hat{\beta}_p x_{ip}.$$

- ▶ This equation represents a plane which we call the regression plane.

Fitted Values and the Hat Matrix

- ▶ These are the values predicted by the linear model for the n subjects.

Fitted Values and the Hat Matrix

- ▶ These are the values predicted by the linear model for the n subjects.
- ▶ The values of the explanatory variables are x_{i1}, \dots, x_{ip} for the i th subject.

Fitted Values and the Hat Matrix

- ▶ These are the values predicted by the linear model for the n subjects.
- ▶ The values of the explanatory variables are x_{i1}, \dots, x_{ip} for the i th subject.
- ▶ Thus the linear model prediction for the i th subject is

$$\hat{\beta}_0 + \hat{\beta}_1 x_{i1} + \dots + \hat{\beta}_p x_{ip}.$$

Fitted Values and the Hat Matrix

- ▶ These are the values predicted by the linear model for the n subjects.
- ▶ The values of the explanatory variables are x_{i1}, \dots, x_{ip} for the i th subject.
- ▶ Thus the linear model prediction for the i th subject is

$$\hat{\beta}_0 + \hat{\beta}_1 x_{i1} + \dots + \hat{\beta}_p x_{ip}.$$

- ▶ Because the value of the response variable for the i th subject is y_i , it makes sense to call the above prediction \hat{y}_i .
Thus

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_{i1} + \dots + \hat{\beta}_p x_{ip}, \quad \text{for } i = 1, \dots, p.$$

Fitted Values and the Hat Matrix

- ▶ These values $\hat{y}_1, \dots, \hat{y}_n$ are called fitted values and the vector $\hat{Y} = (\hat{y}_1, \dots, \hat{y}_n)^T$ is called the vector of fitted values.

Fitted Values and the Hat Matrix

- ▶ These values $\hat{y}_1, \dots, \hat{y}_n$ are called fitted values and the vector $\hat{Y} = (\hat{y}_1, \dots, \hat{y}_n)^T$ is called the vector of fitted values.
- ▶ This vector can be written succinctly as $\hat{Y} = X\hat{\beta}$.

Fitted Values and the Hat Matrix

- ▶ These values $\hat{y}_1, \dots, \hat{y}_n$ are called fitted values and the vector $\hat{Y} = (\hat{y}_1, \dots, \hat{y}_n)^T$ is called the vector of fitted values.
- ▶ This vector can be written succinctly as $\hat{Y} = X\hat{\beta}$.
- ▶ Because $\hat{Y} = X(X^T X)^{-1} X^T Y$, we can write

$$\hat{Y} = X(X^T X)^{-1} X^T Y$$

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.
 2. It is idempotent i.e., $H^2 = H$.

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.
 2. It is idempotent i.e., $H^2 = H$.
 3. $HX = X$.

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.
 2. It is idempotent i.e., $H^2 = H$.
 3. $HX = X$.
- ▶ These can be easily derived from the definition $H = X(X^T X)^{-1} X^T$.

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.
 2. It is idempotent i.e., $H^2 = H$.
 3. $HX = X$.
- ▶ These can be easily derived from the definition $H = X(X^T X)^{-1} X^T$.
- ▶ Because of these, we get

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.
 2. It is idempotent i.e., $H^2 = H$.
 3. $HX = X$.
- ▶ These can be easily derived from the definition $H = X(X^T X)^{-1} X^T$.
- ▶ Because of these, we get

$$\mathbb{E}(\hat{Y}) =$$

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.
 2. It is idempotent i.e., $H^2 = H$.
 3. $HX = X$.
- ▶ These can be easily derived from the definition $H = X(X^T X)^{-1} X^T$.
- ▶ Because of these, we get

$$\mathbb{E}(\hat{Y}) = \mathbb{E}(HY) =$$

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.
 2. It is idempotent i.e., $H^2 = H$.
 3. $HX = X$.
- ▶ These can be easily derived from the definition $H = X(X^T X)^{-1} X^T$.
- ▶ Because of these, we get

$$\mathbb{E}(\hat{Y}) = \mathbb{E}(HY) = H\mathbb{E}(Y) = HX\beta =$$

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.
 2. It is idempotent i.e., $H^2 = H$.
 3. $HX = X$.
- ▶ These can be easily derived from the definition $H = X(X^T X)^{-1} X^T$.
- ▶ Because of these, we get

$$\mathbb{E}(\hat{Y}) = \mathbb{E}(HY) = H\mathbb{E}(Y) = HX\beta =$$

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.
 2. It is idempotent i.e., $H^2 = H$.
 3. $HX = X$.
- ▶ These can be easily derived from the definition $H = X(X^T X)^{-1} X^T$.
- ▶ Because of these, we get

$$\mathbb{E}(\hat{Y}) = \mathbb{E}(HY) = H\mathbb{E}(Y) = HX\beta = X\beta =$$

Fitted Values and the Hat Matrix

- ▶ The vector of fitted values \hat{Y} is the (orthogonal) projection of Y onto the column space of X .
- ▶ Let $H = X(X^T X)^{-1} X^T$ so that $\hat{Y} = HY$.
- ▶ Because multiplication by H changes Y into \hat{Y} , the matrix H is called the Hat Matrix. It is very important in linear regression. It has the following three easily verifiable properties:
 1. It is a symmetric $n \times n$ matrix.
 2. It is idempotent i.e., $H^2 = H$.
 3. $HX = X$.
- ▶ These can be easily derived from the definition $H = X(X^T X)^{-1} X^T$.
- ▶ Because of these, we get

$$\mathbb{E}(\hat{Y}) = \mathbb{E}(HY) = H\mathbb{E}(Y) = HX\beta = X\beta = \mathbb{E}(Y).$$

Fitted Values and the Hat Matrix

- ▶ Thus \hat{Y} and Y have the same expectation.

Fitted Values and the Hat Matrix

- ▶ Thus \hat{Y} and Y have the same expectation.
- ▶ Also

Fitted Values and the Hat Matrix

- ▶ Thus \hat{Y} and Y have the same expectation.
- ▶ Also

$$\text{Cov}(\hat{Y}) = H\text{Cov}(Y)H^T =$$

Fitted Values and the Hat Matrix

- ▶ Thus \hat{Y} and Y have the same expectation.
- ▶ Also

$$\text{Cov}(\hat{Y}) = H\text{Cov}(Y)H^T = H(\sigma^2 I_n)H =$$

Fitted Values and the Hat Matrix

- ▶ Thus \hat{Y} and Y have the same expectation.
- ▶ Also

$$\text{Cov}(\hat{Y}) = H\text{Cov}(Y)H^T = H(\sigma^2 I_n)H = \sigma^2 H.$$

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

$$\hat{e} = \hat{Y} - Y$$

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

$$\hat{e} = Y - \hat{Y} = (I - H)Y.$$

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

$$\hat{e} = \hat{Y} - Y = (I - H)Y.$$

- ▶ The vector of residuals \hat{e} acts as a proxy for the unobserved error vector e .

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

$$\hat{e} = \hat{Y} - Y = (I - H)Y.$$

- ▶ The vector of residuals \hat{e} acts as a proxy for the unobserved error vector e .
- ▶ The most important fact about the residuals in the linear model is that they are orthogonal to the column space of X .

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

$$\hat{e} = \hat{Y} - Y = (I - H)Y.$$

- ▶ The vector of residuals \hat{e} acts as a proxy for the unobserved error vector e .
- ▶ The most important fact about the residuals in the linear model is that they are orthogonal to the column space of X .
- ▶ This happens because $HX = X$ so that

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

$$\hat{e} = \hat{Y} - Y = (I - H)Y.$$

- ▶ The vector of residuals \hat{e} acts as a proxy for the unobserved error vector e .
- ▶ The most important fact about the residuals in the linear model is that they are orthogonal to the column space of X .
- ▶ This happens because $HX = X$ so that
$$\hat{e}^T X =$$

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

$$\hat{e} = \hat{Y} - Y = (I - H)Y.$$

- ▶ The vector of residuals \hat{e} acts as a proxy for the unobserved error vector e .
- ▶ The most important fact about the residuals in the linear model is that they are orthogonal to the column space of X .
- ▶ This happens because $HX = X$ so that

$$\hat{e}^T X = ((I - H)Y)^T X =$$

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

$$\hat{e} = \hat{Y} - Y = (I - H)Y.$$

- ▶ The vector of residuals \hat{e} acts as a proxy for the unobserved error vector e .
- ▶ The most important fact about the residuals in the linear model is that they are orthogonal to the column space of X .
- ▶ This happens because $HX = X$ so that

$$\hat{e}^T X = ((I - H)Y)^T X = Y^T (I - H)X =$$

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

$$\hat{e} = \hat{Y} - Y = (I - H)Y.$$

- ▶ The vector of residuals \hat{e} acts as a proxy for the unobserved error vector e .
- ▶ The most important fact about the residuals in the linear model is that they are orthogonal to the column space of X .
- ▶ This happens because $HX = X$ so that

$$\hat{e}^T X = ((I - H)Y)^T X = Y^T (I - H)X = Y^T (X - HX) = 0.$$

Residuals

- ▶ The difference between y_i and \hat{y}_i is called the residual for the i th subject: $\hat{e}_i := y_i - \hat{y}_i$.
- ▶ The vector $\hat{e} = (\hat{e}_1, \dots, \hat{e}_n)^T$ is called the vector of residuals

$$\hat{e} = \hat{Y} - Y = (I - H)Y.$$

- ▶ The vector of residuals \hat{e} acts as a proxy for the unobserved error vector e .
- ▶ The most important fact about the residuals in the linear model is that they are orthogonal to the column space of X .
- ▶ This happens because $HX = X$ so that
$$\hat{e}^T X = ((I - H)Y)^T X = Y^T (I - H)X = Y^T (X - HX) = 0.$$
- ▶ As a result $\hat{e}^T Xu = 0$ for every vector u which means that \hat{e} is orthogonal to the column space of X .

Residuals

- ▶ The first column of X consists of ones.


Residuals

- ▶ The first column of X consists of ones.
- ▶ Because \hat{e} is orthogonal to everything in the column space of X , it must therefore be orthogonal to the vector of ones which means that

$$\sum_{i=1}^n \hat{e}_i = 0.$$

Residuals

- ▶ The first column of X consists of ones.
- ▶ Because \hat{e} is orthogonal to everything in the column space of X , it must therefore be orthogonal to the vector of ones which means that

$$\sum_{i=1}^n \hat{e}_i = 0.$$


- ▶ \hat{e} is also orthogonal to every column of X :

$$\sum_{i=1}^n \hat{e}_i x_{ij} = 0 \quad \text{for every } j.$$

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.
- ▶ Hence, although there are n residuals, there are effectively $n - p - 1$ of them. The number $n - p - 1$ is therefore referred to as the degrees of freedom of the residuals $\hat{e}_1, \dots, \hat{e}_n$.

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.
- ▶ Hence, although there are n residuals, there are effectively $n - p - 1$ of them. The number $n - p - 1$ is therefore referred to as the degrees of freedom of the residuals $\hat{e}_1, \dots, \hat{e}_n$.
- ▶ The expectation of \hat{e} is

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.
- ▶ Hence, although there are n residuals, there are effectively $n - p - 1$ of them. The number $n - p - 1$ is therefore referred to as the degrees of freedom of the residuals $\hat{e}_1, \dots, \hat{e}_n$.
- ▶ The expectation of \hat{e} is

$$\mathbb{E}(\hat{e}) = \mathbb{E}((I - H)Y) =$$

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.
- ▶ Hence, although there are n residuals, there are effectively $n - p - 1$ of them. The number $n - p - 1$ is therefore referred to as the degrees of freedom of the residuals $\hat{e}_1, \dots, \hat{e}_n$.
- ▶ The expectation of \hat{e} is

$$\mathbb{E}(\hat{e}) = \mathbb{E}((I - H)Y) = (I - H)\mathbb{E}(Y) =$$

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.
- ▶ Hence, although there are n residuals, there are effectively $n - p - 1$ of them. The number $n - p - 1$ is therefore referred to as the degrees of freedom of the residuals $\hat{e}_1, \dots, \hat{e}_n$.
- ▶ The expectation of \hat{e} is

$$\mathbb{E}(\hat{e}) = \mathbb{E}((I - H)Y) = (I - H)\mathbb{E}(Y) = (I - H)X\beta = 0.$$

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.
- ▶ Hence, although there are n residuals, there are effectively $n - p - 1$ of them. The number $n - p - 1$ is therefore referred to as the degrees of freedom of the residuals $\hat{e}_1, \dots, \hat{e}_n$.
- ▶ The expectation of \hat{e} is

$$\mathbb{E}(\hat{e}) = \mathbb{E}((I - H)Y) = (I - H)\mathbb{E}(Y) = (I - H)X\beta = 0.$$

- ▶ The Covariance matrix of \hat{e} is

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.
- ▶ Hence, although there are n residuals, there are effectively $n - p - 1$ of them. The number $n - p - 1$ is therefore referred to as the degrees of freedom of the residuals $\hat{e}_1, \dots, \hat{e}_n$.
- ▶ The expectation of \hat{e} is

$$\mathbb{E}(\hat{e}) = \mathbb{E}((I - H)Y) = (I - H)\mathbb{E}(Y) = (I - H)X\beta = 0.$$

- ▶ The Covariance matrix of \hat{e} is

$$\text{Cov}(\hat{e}) = \text{Cov}((I - H)Y) =$$

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.
- ▶ Hence, although there are n residuals, there are effectively $n - p - 1$ of them. The number $n - p - 1$ is therefore referred to as the degrees of freedom of the residuals $\hat{e}_1, \dots, \hat{e}_n$.
- ▶ The expectation of \hat{e} is

$$\mathbb{E}(\hat{e}) = \mathbb{E}((I - H)Y) = (I - H)\mathbb{E}(Y) = (I - H)X\beta = 0.$$

- ▶ The Covariance matrix of \hat{e} is

$$\text{Cov}(\hat{e}) = \text{Cov}((I - H)Y) = (I - H)\text{Cov}(Y)(I - H)^T =$$

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.
- ▶ Hence, although there are n residuals, there are effectively $n - p - 1$ of them. The number $n - p - 1$ is therefore referred to as the degrees of freedom of the residuals $\hat{e}_1, \dots, \hat{e}_n$.
- ▶ The expectation of \hat{e} is

$$\mathbb{E}(\hat{e}) = \mathbb{E}((I - H)Y) = (I - H)\mathbb{E}(Y) = (I - H)X\beta = 0.$$

- ▶ The Covariance matrix of \hat{e} is

$$\text{Cov}(\hat{e}) = \text{Cov}((I - H)Y) = (I - H)\text{Cov}(Y)(I - H)^T = \sigma^2(I - H).$$

Residuals

- ▶ The vector of fitted values belongs to the column space of X because $\hat{Y} = X\hat{\beta}$.
- ▶ Therefore \hat{e} is also orthogonal to \hat{Y} .
- ▶ Because $X^T \hat{e} = 0$, the residuals satisfy $p + 1$ linear equalities.
- ▶ Hence, although there are n residuals, there are effectively $n - p - 1$ of them. The number $n - p - 1$ is therefore referred to as the degrees of freedom of the residuals $\hat{e}_1, \dots, \hat{e}_n$.
- ▶ The expectation of \hat{e} is

$$\mathbb{E}(\hat{e}) = \mathbb{E}((I - H)Y) = (I - H)\mathbb{E}(Y) = (I - H)X\beta = 0.$$

- ▶ The Covariance matrix of \hat{e} is

$$\text{Cov}(\hat{e}) = \text{Cov}((I - H)Y) = (I - H)\text{Cov}(Y)(I - H)^T = \sigma^2(I - H).$$

- ▶ Note that the residuals have different variances.

Residuals

- Note that the residuals have different variances. (even though e_1, \dots, e_n are assumed to have the same variance σ^2).

Levarages

- ▶ The diagonal entries of the hat matrix are called leverages and these are denoted by $h_{ii}, i = 1, \dots, n$.

Leverages

- ▶ The diagonal entries of the hat matrix are called leverages and these are denoted by $h_{ii}, i = 1, \dots, n$.
- ▶ We will study them carefully later on under the topic of regression diagnostics. For now, let us just remark that the leverages are always between 0 and 1.

Levarages

- ▶ The diagonal entries of the hat matrix are called leverages and these are denoted by $h_{ii}, i = 1, \dots, n$.
- ▶ We will study them carefully later on under the topic of regression diagnostics. For now, let us just remark that the leverages are always between 0 and 1.
- ▶ This can be proved in many ways but a simple way is to observe that $h_{ii} = \text{var}(y_i) \geq 0, 1 - h_{ii} = \text{var}(\hat{e}_i) \geq 0$.