

# Partial regression plot

October 29, 2018

## Theorem

*Let  $\hat{\beta}$  be the vector of coefficients from the regression of  $Y$  on  $X$ , with residuals  $\hat{e}$ , and  $X^{(j)}$  the  $j$ -th column of  $X$ . Let  $b^{(j)}$  be the slope from the simple linear regression of  $Y^{(j)} := \text{Res}(Y, X^{(j)})$  on  $X^{(j)} := \text{Res}(X^{(j)}, X^{-j})$ , with residuals  $e^{(j)}$ . Then  $b^{(j)} = \hat{\beta}_j$ .*

## Theorem

*Let  $\hat{\beta}$  be the vector of coefficients from the regression of  $Y$  on  $X$ , with residuals  $\hat{e}$ , and  $X^{(j)}$  the  $j$ -th column of  $X$ . Let  $b^{(j)}$  be the slope from the simple linear regression of  $Y^{(j)} := \text{Res}(Y, X^{(j)})$  on  $X^{(j)} := \text{Res}(X^{(j)}, X^{-j})$ , with residuals  $e^{(j)}$ . Then  $b^{(j)} = \hat{\beta}_j$ .*



- ▶ We prove the case  $j = p$ ; the generalization is obvious.

## Theorem

Let  $\hat{\beta}$  be the vector of coefficients from the regression of  $Y$  on  $X$ , with residuals  $\hat{e}$ , and  $X^{(j)}$  the  $j$ -th column of  $X$ . Let  $b^{(j)}$  be the slope from the simple linear regression of  $Y^{(j)} := \text{Res}(Y, X^{(j)})$  on  $X^{(j)} := \text{Res}(X^{(j)}, X^{-j})$ , with residuals  $e^{(j)}$ . Then  $b^{(j)} = \hat{\beta}_j$ .



- ▶ We prove the case  $j = p$ ; the generalization is obvious.
- ▶ First, let's consider the simple linear regression of  $Y^{(p)}$  on  $X^{(p)}$ .

## Theorem

Let  $\hat{\beta}$  be the vector of coefficients from the regression of  $Y$  on  $X$ , with residuals  $\hat{e}$ , and  $X^{(j)}$  the  $j$ -th column of  $X$ . Let  $b^{(j)}$  be the slope from the simple linear regression of  $Y^{(j)} := \text{Res}(Y, X^{(j)})$  on  $X^{(j)} := \text{Res}(X, X^{-j})$ , with residuals  $e^{(j)}$ . Then  $b^{(j)} = \hat{\beta}_j$ .



- ▶ We prove the case  $j = p$ ; the generalization is obvious.
- ▶ First, let's consider the simple linear regression of  $Y^{(p)}$  on  $X^{(p)}$ .
- ▶ Let  $H(-p)$  be the projection matrix on all columns in  $X$  except  $X^{(p)}$ , the  $p$ th column.

## Theorem

Let  $\hat{\beta}$  be the vector of coefficients from the regression of  $Y$  on  $X$ , with residuals  $\hat{e}$ , and  $X^{(j)}$  the  $j$ -th column of  $X$ . Let  $b^{(j)}$  be the slope from the simple linear regression of  $Y^{(j)} := \text{Res}(Y, X^{(j)})$  on  $X^{(j)} := \text{Res}(X, X^{-j})$ , with residuals  $e^{(j)}$ . Then  $b^{(j)} = \hat{\beta}_j$ .



- ▶ We prove the case  $j = p$ ; the generalization is obvious.
- ▶ First, let's consider the simple linear regression of  $Y^{(p)}$  on  $X^{(p)}$ .
- ▶ Let  $H(-p)$  be the projection matrix on all columns in  $X$  except  $X^{(p)}$ , the  $p$ th column.
- ▶ Then  $Y^{(p)} = (I - H(-p))Y$  and  $X^{(p)} = (I - H(-p))X^{(p)}$ .

## Theorem

Let  $\hat{\beta}$  be the vector of coefficients from the regression of  $Y$  on  $X$ , with residuals  $\hat{e}$ , and  $X^{(j)}$  the  $j$ -th column of  $X$ . Let  $b^{(j)}$  be the slope from the simple linear regression of  $Y^{(j)} := \text{Res}(Y, X^{(j)})$  on  $X^{(j)} := \text{Res}(X^{(j)}, X^{-j})$ , with residuals  $e^{(j)}$ . Then  $b^{(j)} = \hat{\beta}_j$ .



- ▶ We prove the case  $j = p$ ; the generalization is obvious.
- ▶ First, let's consider the simple linear regression of  $Y^{(p)}$  on  $X^{(p)}$ .
- ▶ Let  $H(-p)$  be the projection matrix on all columns in  $X$  except  $X^{(p)}$ , the  $p$ th column.
- ▶ Then  $Y^{(p)} = (I - H(-p))Y$  and  $X^{(p)} = (I - H(-p))X^{(p)}$ .
- ▶ Since both  $Y^{(p)}$  and  $X^{(p)}$  are residuals from a fitted linear model with an intercept, their entries have sample mean 0; therefore this is simple linear regression through the origin.

► Hence:

$$b^{(p)} = \frac{(X^{(p)})^T Y^{(p)}}{\|X^{(p)}\|_2^2}.$$



► Hence:

$$b^{(p)} = \frac{(X^{(p)})^T Y^{(p)}}{\|X^{(p)}\|_2^2}.$$

► And so

$$b^{(p)} = \frac{Y^T (I - H(-p)) X(p)}{X(p)^T (I - H(-p)) X(p)}.$$

- ▶ Hence:

$$b^{(p)} = \frac{(X^{(p)})^T Y^{(p)}}{\|X^{(p)}\|_2^2}.$$

- ▶ And so

$$b^{(p)} = \frac{Y^T (I - H(-p)) X(p)}{X(p)^T (I - H(-p)) X(p)}.$$

- ▶ We need to prove that  $\hat{\beta}_p$  is also equal to this quantity.

- ▶ Hence:


$$b^{(p)} = \frac{(X^{(p)})^T Y^{(p)}}{\|X^{(p)}\|_2^2}.$$

- ▶ And so

$$b^{(p)} = \frac{Y^T (I - H(-p)) X(p)}{X(p)^T (I - H(-p)) X(p)}.$$

- ▶ We need to prove that  $\hat{\beta}_p$  is also equal to this quantity.
- ▶ To do so, consider fitting a linear regression under the constraint  $\beta_p = b$  for some fixed value  $b$ ; the corresponding minimization problem would be the following:

- ▶ Hence:


$$b^{(p)} = \frac{(X^{(p)})^T Y^{(p)}}{\|X^{(p)}\|_2^2}.$$

- ▶ And so

$$b^{(p)} = \frac{Y^T (I - H(-p)) X^{(p)}}{X^{(p)T} (I - H(-p)) X^{(p)}}.$$

- ▶ We need to prove that  $\hat{\beta}_p$  is also equal to this quantity.
- ▶ To do so, consider fitting a linear regression under the constraint  $\beta_p = b$  for some fixed value  $b$ ; the corresponding minimization problem would be the following:



$$\min_{\alpha \in \mathbb{R}^p} \|Y - V\alpha - X^{(p)}b\|_2^2$$

where  $V$  contains the first  $p$  columns of  $X$ . Let  $\alpha(b)$  denote the minimizer of the above problem.

- Note that  $\alpha(0)$  is the least squares regression of  $Y$  on  $V$ , while  $\alpha(\hat{\beta}_p)$  represents the first  $p$  coefficients of the full linear regression of  $Y$  on  $X$ .

- ▶ Note that  $\alpha(0)$  is the least squares regression of  $Y$  on  $V$ , while  $\alpha(\hat{\beta}_p)$  represents the first  $p$  coefficients of the full linear regression of  $Y$  on  $X$ .
- ▶ We want a formula for  $\alpha(b)$  for all  $b$ ; therefore we take the gradient in to obtain the stationary condition.

- ▶ Note that  $\alpha(0)$  is the least squares regression of  $Y$  on  $V$ , while  $\alpha(\hat{\beta}_p)$  represents the first  $p$  coefficients of the full linear regression of  $Y$  on  $X$ .
- ▶ We want a formula for  $\alpha(b)$  for all  $b$ ; therefore we take the gradient in to obtain the stationary condition. Which, upon solving for, yields:

$$\alpha(b) = (V^T V)^{-1} V^T (Y - X(p)b).$$

- ▶ Note that  $\alpha(0)$  is the least squares regression of  $Y$  on  $V$ , while  $\alpha(\hat{\beta}_p)$  represents the first  $p$  coefficients of the full linear regression of  $Y$  on  $X$ .
- ▶ We want a formula for  $\alpha(b)$  for all  $b$ ; therefore we take the gradient in to obtain the stationary condition. Which, upon solving for, yields:

$$\alpha(b) = (V^T V)^{-1} V^T (Y - X(p)b).$$

- ▶ Note that  $X\hat{\beta} = V\alpha(\hat{\beta}_p) + X(p)\hat{\beta}_p$ ; combined with the formula for  $\alpha(b)$  this yields:

$$\begin{aligned} X\hat{\beta} &= V(V^T V)^{-1} V^T (Y - X(p)\hat{\beta}_p) + X(p)\hat{\beta}_p \\ &= H(-p)Y + (I - H(-p))X(p)\hat{\beta}_p \\ &= HY, \end{aligned}$$

where the last equality is the fundamental equality of least squares (here  $H$  is the projection matrix on all columns of  $X$ ).



- Multiplying the above equalities through by  $X(p)^T$  yields:

$$\begin{aligned} X(p)^T Y &= X(p)^T H Y \\ &= X(p)^T H(-p) Y + X(p)^T (I - H(-p)) X(p) \hat{\beta}_p, \end{aligned}$$

solving for  $\hat{\beta}_p$  leads to the desired result.