

Lecture 23

November 7, 2018

Maximum Likelihood in Logistic Regression

- ▶ We have binary responses y_1, \dots, y_n and data on p explanatory variables x_{ij} , $i = 1, \dots, n$ and $j = 1, \dots, p$.

Maximum Likelihood in Logistic Regression

- ▶ We have binary responses y_1, \dots, y_n and data on p explanatory variables $x_{ij}, i = 1, \dots, n$ and $j = 1, \dots, p$.
- ▶ We assume that y_1, \dots, y_n are independent Bernoulli random variables with parameters p_1, \dots, p_n .

Maximum Likelihood in Logistic Regression

- ▶ We have binary responses y_1, \dots, y_n and data on p explanatory variables x_{ij} , $i = 1, \dots, n$ and $j = 1, \dots, p$.
- ▶ We assume that y_1, \dots, y_n are independent Bernoulli random variables with parameters p_1, \dots, p_n .
- ▶ We model the relationship between the response and explanatory variables by the formula

$$\log \frac{p_i}{1 - p_i} = \beta_0 + \beta_1 x_{i1} + \dots + \beta_p x_{ip}. \quad (1)$$

Maximum Likelihood in Logistic Regression

- ▶ We have binary responses y_1, \dots, y_n and data on p explanatory variables x_{ij} , $i = 1, \dots, n$ and $j = 1, \dots, p$.
- ▶ We assume that y_1, \dots, y_n are independent Bernoulli random variables with parameters p_1, \dots, p_n .
- ▶ We model the relationship between the response and explanatory variables by the formula

$$\log \frac{p_i}{1 - p_i} = \beta_0 + \beta_1 x_{i1} + \dots + \beta_p x_{ip}. \quad (1)$$

- ▶ Given data y_1, \dots, y_n and x_{ij} for $i = 1, \dots, n$ and $j = 1, \dots, p$, how can we estimate the parameters β_0, \dots, β_p .

- Note that the model can alternatively be written as

$$y_i \sim \text{Ber} \left(\frac{\exp(\beta_0 + \sum_{j=1}^p \beta_j x_{ij})}{1 + \exp(\beta_0 + \sum_{j=1}^p \beta_j x_{ij})} \right)$$

with y_1, \dots, y_n being independent.

- Note that the model can alternatively be written as

$$y_i \sim \text{Ber} \left(\frac{\exp(\beta_0 + \sum_{j=1}^p \beta_j x_{ij})}{1 + \exp(\beta_0 + \sum_{j=1}^p \beta_j x_{ij})} \right)$$

with y_1, \dots, y_n being independent.

- We use maximum likelihood to estimate β_0, \dots, β_p . The log-likelihood of the data y_1, \dots, y_n (we take X to be deterministic) is

$$\begin{aligned} \ell(\beta) &= \sum_{i=1}^n (y_i \log p_i + (1 - y_i) \log(1 - p_i)) \\ &= \sum_{i=1}^n \left[y_i (\beta_0 + \beta_1 x_{i1} \cdots + \beta_p x_{ip}) - \right. \\ &\quad \left. \log(1 + \exp(\beta_0 + \beta_1 x_{i1} \cdots + \beta_p x_{ip})) \right]. \end{aligned}$$

- ▶ The MLE of β is the maximizer of $\ell(\beta)$.

- ▶ The MLE of β is the maximizer of $\ell(\beta)$.
- ▶ The maximizer of $\ell(\beta)$ cannot be computed in closed form. We use Newton's method for maximizing $\ell(\beta)$.

- ▶ The MLE of β is the maximizer of $\ell(\beta)$.
- ▶ The maximizer of $\ell(\beta)$ cannot be computed in closed form. We use Newton's method for maximizing $\ell(\beta)$.
- ▶ Newton's method uses the iterative scheme

$$\beta^{(m+1)} = \beta^{(m)} - \left(H\ell(\beta^{(m)}) \right)^{-1} \nabla \ell(\beta^{(m)}) \quad (2)$$

where $\nabla \ell(\beta)$ and $H\ell(\beta)$ denote the gradient and Hessian of the function $\ell(\beta)$ respectively:

$\nabla \ell(\beta) := (\partial \ell(\beta) / \partial \beta_0, \dots, \partial \ell(\beta) / \partial \beta_p)^T$ and $H\ell(\beta)$ is the $(p+1) \times (p+1)$ matrix whose entries are second order derivatives of $\ell(\beta)$.

- For example, the $(1, 1)$ th entry of $H\ell(\beta)$ is $\partial^2 \ell(\beta) / \partial \beta_0^2$, the $(1, 2)$ th entry is $\partial^2 \ell(\beta) / \partial \beta_0 \partial \beta_1$ and so on.

- ▶ For example, the $(1, 1)$ th entry of $H\ell(\beta)$ is $\partial^2\ell(\beta)/\partial\beta_0^2$, the $(1, 2)$ th entry is $\partial^2\ell(\beta)/\partial\beta_0\partial\beta_1$ and so on.
- ▶ We saw in the last class that

$$\nabla\ell(\beta) = X^T(Y - p) \quad \text{and} \quad H\ell(\beta) = -X^T W X$$

where W is the $n \times n$ diagonal matrix whose i^{th} diagonal entry is $p_i(1 - p_i)$.

- ▶ For example, the $(1, 1)$ th entry of $H\ell(\beta)$ is $\partial^2\ell(\beta)/\partial\beta_0^2$, the $(1, 2)$ th entry is $\partial^2\ell(\beta)/\partial\beta_0\partial\beta_1$ and so on.
- ▶ We saw in the last class that

$$\nabla\ell(\beta) = X^T(Y - p) \quad \text{and} \quad H\ell(\beta) = -X^T W X$$

where W is the $n \times n$ diagonal matrix whose i^{th} diagonal entry is $p_i(1 - p_i)$.

- ▶ Newton's iterative scheme (2) therefore becomes

$$\beta^{(m+1)} = \beta^{(m)} + (X^T W X)^{-1} X^T (Y - p).$$

- ▶ For example, the $(1, 1)$ th entry of $H\ell(\beta)$ is $\partial^2 \ell(\beta) / \partial \beta_0^2$, the $(1, 2)$ th entry is $\partial^2 \ell(\beta) / \partial \beta_0 \partial \beta_1$ and so on.
- ▶ We saw in the last class that

$$\nabla \ell(\beta) = X^T(Y - p) \quad \text{and} \quad H\ell(\beta) = -X^T W X$$

where W is the $n \times n$ diagonal matrix whose i^{th} diagonal entry is $p_i(1 - p_i)$.

- ▶ Newton's iterative scheme (2) therefore becomes

$$\beta^{(m+1)} = \beta^{(m)} + (X^T W X)^{-1} X^T (Y - p).$$

- ▶ This can be rewritten as

$$\beta^{(m+1)} = (X^T W X)^{-1} X^T W Z \tag{3}$$

► where

$$Z = X\beta^{(m)} + W^{-1}(Y - p). \quad (4)$$

- ▶ where

$$Z = X\beta^{(m)} + W^{-1}(Y - p). \quad (4)$$

- ▶ The method of estimating β therefore proceeds iteratively as follows.

- ▶ where

$$Z = X\beta^{(m)} + W^{-1}(Y - p). \quad (4)$$

- ▶ The method of estimating β therefore proceeds iteratively as follows.
- ▶ First have an initial estimate of β_0, \dots, β_p .

- ▶ where

$$Z = X\beta^{(m)} + W^{-1}(Y - p). \quad (4)$$

- ▶ The method of estimating β therefore proceeds iteratively as follows.
- ▶ First have an initial estimate of β_0, \dots, β_p .
- ▶ Call this initial estimator $\hat{\beta}^{(0)}$. Use this estimator to calculate p_i via

$$p_i = \frac{\exp(\hat{\beta}_0^{(0)} + \hat{\beta}_1^{(0)}x_{i1} + \dots + \hat{\beta}_p^{(0)}x_{ip})}{1 + \exp(\hat{\beta}_0^{(0)} + \hat{\beta}_1^{(0)}x_{i1} + \dots + \hat{\beta}_p^{(0)}x_{ip})}.$$

- ▶ Use these values of p_i to create the response variable values Z_i via (6) and also use values of p_i to construct the matrix W . With Z and W , we can estimate β via

$$\hat{\beta}^{(1)} = (X^T W X)^{-1} X^T W Z.$$

- ▶ Use these values of p_i to create the response variable values Z_i via (6) and also use values of p_i to construct the matrix W . With Z and W , we can estimate β via

$$\hat{\beta}^{(1)} = (X^T W X)^{-1} X^T W Z.$$

- ▶ Now replace the initial estimator $\hat{\beta}^{(0)}$ by $\hat{\beta}^{(1)}$ and repeat this process. Keep repeating this until two successive estimates $\hat{\beta}^{(m)}$ and $\hat{\beta}^{(m+1)}$ do not change much. At that point, stop and report the estimate of β in the logistic regression model as $\hat{\beta}^{(m)}$.

- ▶ The expression $(X^T W X)^{-1} X^T W Z$ is reminiscent of the usual $(X^T X)^{-1} X^T Y$ which is the usual estimate of β in the linear model. In fact, this is the least squares estimate in a weighted least squares model as we shall describe next.

Weighted Least Squares

- ▶ Consider regression data in the usual set-up. Suppose we think that the right model is:

$$Y = X\beta + e \quad \text{where, } \mathbb{E}(e) = 0, \quad \text{and, } \text{Cov}(e) = \sigma^2 V$$

for some known (positive definite) matrix V .

Weighted Least Squares

- ▶ Consider regression data in the usual set-up. Suppose we think that the right model is:

$$Y = X\beta + e \quad \text{where, } \mathbb{E}(e) = 0, \quad \text{and, } \text{Cov}(e) = \sigma^2 V$$

for some known (positive definite) matrix V .

- ▶ What then is a good estimator of β ?

Weighted Least Squares

- ▶ Consider regression data in the usual set-up. Suppose we think that the right model is:

$$Y = X\beta + e \quad \text{where, } \mathbb{E}(e) = 0, \quad \text{and, } \text{Cov}(e) = \sigma^2 V$$

for some known (positive definite) matrix V .

- ▶ What then is a good estimator of β ?
- ▶ The difference from the usual situation is the presence of this matrix V .

Weighted Least Squares

- ▶ Consider regression data in the usual set-up. Suppose we think that the right model is:

$$Y = X\beta + e \quad \text{where, } \mathbb{E}(e) = 0, \quad \text{and, } \text{Cov}(e) = \sigma^2 V$$

for some known (positive definite) matrix V .

- ▶ What then is a good estimator of β ?
- ▶ The difference from the usual situation is the presence of this matrix V .
- ▶ It turns out the usual least squares estimator is not a good choice here for estimating β . It is better to use the weighted least squares estimator:

$$\hat{\beta}_{wls} := (X^T V^{-1} X)^{-1} X^T V^{-1} Y. \quad (5)$$

- ▶ It is not too hard to see that this estimator minimizes the weighted sum of squares

$$(Y - X\beta)^T V^{-1} (Y - X\beta)$$

over all β .

- ▶ It is not too hard to see that this estimator minimizes the weighted sum of squares

$$(Y - X\beta)^T V^{-1} (Y - X\beta)$$

over all β .

- ▶ Why is it sensible to use (5) for estimating β in this case?

- ▶ It is not too hard to see that this estimator minimizes the weighted sum of squares

$$(Y - X\beta)^T V^{-1} (Y - X\beta)$$

over all β .

- ▶ Why is it sensible to use (5) for estimating β in this case?
- ▶ The follows reasons motivate this choice:

- ▶ It is not too hard to see that this estimator minimizes the weighted sum of squares

$$(Y - X\beta)^T V^{-1} (Y - X\beta)$$

over all β .

- ▶ Why is it sensible to use (5) for estimating β in this case?
- ▶ The follows reasons motivate this choice:
- ▶ If e is multivariate normal, then (5) is the mle for β .

- Suppose V is diagonal. Then it is obvious that

$$(Y - X\beta)^T V^{-1} (Y - X\beta) = \sum_{i=1}^n \frac{(y_i - \beta_0 - \beta_1 x_{i1} - \cdots - \beta_p x_{ip})^2}{v_{ii}}$$

where v_{ii} denotes the i th diagonal entry of V . It is intuitively clear that minimizing this weighted sum of squares as opposed to the unweighted sum of squares is the right thing to do here.

- Suppose V is diagonal. Then it is obvious that

$$(Y - X\beta)^T V^{-1} (Y - X\beta) = \sum_{i=1}^n \frac{(y_i - \beta_0 - \beta_1 x_{i1} - \cdots - \beta_p x_{ip})^2}{v_{ii}}$$

where v_{ii} denotes the i th diagonal entry of V . It is intuitively clear that minimizing this weighted sum of squares as opposed to the unweighted sum of squares is the right thing to do here.

- For example, if v_{ii} is very high, it means that the i th observation is not very trustworthy and it therefore makes sense to give it low weight.

- ▶ Suppose V is diagonal. Then it is obvious that

$$(Y - X\beta)^T V^{-1} (Y - X\beta) = \sum_{i=1}^n \frac{(y_i - \beta_0 - \beta_1 x_{i1} - \dots - \beta_p x_{ip})^2}{v_{ii}}$$

where v_{ii} denotes the i th diagonal entry of V . It is intuitively clear that minimizing this weighted sum of squares as opposed to the unweighted sum of squares is the right thing to do here.

- ▶ For example, if v_{ii} is very high, it means that the i th observation is not very trustworthy and it therefore makes sense to give it low weight.
- ▶ In the same way, it makes sense to give large weight to the i th observation if v_{ii} is low.

- ▶ One can show that $\hat{\beta}_{wls}$ is the BLUE for β .

- ▶ One can show that $\hat{\beta}_{wls}$ is the BLUE for β .
- ▶ The expectation and the covariance matrix of $\hat{\beta}_{wls}$ can be easily calculated via:

- ▶ One can show that $\hat{\beta}_{wls}$ is the BLUE for β .
- ▶ The expectation and the covariance matrix of $\hat{\beta}_{wls}$ can be easily calculated via:



$$\mathbb{E}\hat{\beta}_{wls} = \beta \quad \text{and} \quad \text{Cov}(\hat{\beta}_{wls}) = \sigma^2(X^T V^{-1} X)^{-1}.$$

Iteratively Reweighed Least Squares for Logistic Regression Fitting

- ▶ Because of the similarity between (3) and (5), Newton's method for computing the maximum likelihood estimator in logistic regression can be seen as a sequence of weighted least squares estimators.

Iteratively Reweighed Least Squares for Logistic Regression Fitting

- ▶ Because of the similarity between (3) and (5), Newton's method for computing the maximum likelihood estimator in logistic regression can be seen as a sequence of weighted least squares estimators.
- ▶ That is why the iterative method is also called IRLS (Iteratively Reweighted Least Squares) or IWLS (Iteratively Weighted Least Squares).

Iteratively Reweighed Least Squares for Logistic Regression Fitting

- ▶ Because of the similarity between (3) and (5), Newton's method for computing the maximum likelihood estimator in logistic regression can be seen as a sequence of weighted least squares estimators.
- ▶ That is why the iterative method is also called IRLS (Iteratively Reweighted Least Squares) or IWLS (Iteratively Weighted Least Squares).
- ▶ Here is a more intuitive approach to understand IRLS. The goal is to fit the model (1) to the data.

Iteratively Reweighed Least Squares for Logistic Regression Fitting

- ▶ Because of the similarity between (3) and (5), Newton's method for computing the maximum likelihood estimator in logistic regression can be seen as a sequence of weighted least squares estimators.
- ▶ That is why the iterative method is also called IRLS (Iteratively Reweighted Least Squares) or IWLS (Iteratively Weighted Least Squares).
- ▶ Here is a more intuitive approach to understand IRLS. The goal is to fit the model (1) to the data.
- ▶ Because $p_i = \mathbb{E}y_i$, the equation (1) can be rewritten as

$$\log \frac{\mathbb{E}(y_i)}{1 - \mathbb{E}(y_i)} = \beta_0 + \beta_1 x_{i1} + \cdots + \beta_p x_{ip}.$$

- ▶ Because of the form above, a first idea to fit this model to data might be to try to fit a linear model to the response variable $\log(y_i/(1 - y_i))$ on the explanatory variables and then to estimate β_0, \dots, β_p by the estimated coefficients of that linear model.

- ▶ Because of the form above, a first idea to fit this model to data might be to try to fit a linear model to the response variable $\log(y_i/(1 - y_i))$ on the explanatory variables and then to estimate β_0, \dots, β_p by the estimated coefficients of that linear model.
- ▶ But because y_i is 0 or 1, the quantity $\log(y_i/(1 - y_i))$ is either $-\infty$ or ∞ and so this response variable would make no sense.

- ▶ Because of the form above, a first idea to fit this model to data might be to try to fit a linear model to the response variable $\log(y_i/(1 - y_i))$ on the explanatory variables and then to estimate β_0, \dots, β_p by the estimated coefficients of that linear model.
- ▶ But because y_i is 0 or 1, the quantity $\log(y_i/(1 - y_i))$ is either $-\infty$ or ∞ and so this response variable would make no sense.
- ▶ A way to fix this is to work with a response variable that is similar in spirit to $\log(y_i/(1 - y_i))$ but which actually makes sense.

- ▶ Because of the form above, a first idea to fit this model to data might be to try to fit a linear model to the response variable $\log(y_i/(1 - y_i))$ on the explanatory variables and then to estimate β_0, \dots, β_p by the estimated coefficients of that linear model.
- ▶ But because y_i is 0 or 1, the quantity $\log(y_i/(1 - y_i))$ is either $-\infty$ or ∞ and so this response variable would make no sense.
- ▶ A way to fix this is to work with a response variable that is similar in spirit to $\log(y_i/(1 - y_i))$ but which actually makes sense.
- ▶ Let $g(x) = \log(x/(1 - x))$. By a first order Taylor expansion to g around p_i , we can write

$$g(y_i) \approx g(p_i) + g'(p_i)(y_i - p_i) = \log \frac{p_i}{1 - p_i} + \frac{y_i - p_i}{p_i(1 - p_i)}$$

- The right hand side above makes sense as opposed to $g(y_i)$. So we let

$$Z_i = \log \frac{p_i}{1 - p_i} + \frac{y_i - p_i}{p_i(1 - p_i)} \quad (6)$$

and we can fit a linear model to Z_i based on the explanatory variables and estimate β by the estimated coefficients in that linear model.

- ▶ The right hand side above makes sense as opposed to $g(y_i)$. So we let

$$Z_i = \log \frac{p_i}{1 - p_i} + \frac{y_i - p_i}{p_i(1 - p_i)} \quad (6)$$

and we can fit a linear model to Z_i based on the explanatory variables and estimate β by the estimated coefficients in that linear model.

- ▶ Should we estimate the coefficients of that linear model by ordinary least squares or should we use weighted least squares?

- ▶ The right hand side above makes sense as opposed to $g(y_i)$. So we let

$$Z_i = \log \frac{p_i}{1 - p_i} + \frac{y_i - p_i}{p_i(1 - p_i)} \quad (6)$$

and we can fit a linear model to Z_i based on the explanatory variables and estimate β by the estimated coefficients in that linear model.

- ▶ Should we estimate the coefficients of that linear model by ordinary least squares or should we use weighted least squares? The variance of Z_i is:

$$\text{var}(Z_i) = \text{var} \left(\frac{y_i - p_i}{p_i(1 - p_i)} \right) = \frac{1}{p_i(1 - p_i)}.$$

- ▶ The right hand side above makes sense as opposed to $g(y_i)$. So we let

$$Z_i = \log \frac{p_i}{1 - p_i} + \frac{y_i - p_i}{p_i(1 - p_i)} \quad (6)$$

and we can fit a linear model to Z_i based on the explanatory variables and estimate β by the estimated coefficients in that linear model.

- ▶ Should we estimate the coefficients of that linear model by ordinary least squares or should we use weighted least squares? The variance of Z_i is:

$$\text{var}(Z_i) = \text{var} \left(\frac{y_i - p_i}{p_i(1 - p_i)} \right) = \frac{1}{p_i(1 - p_i)}.$$

- ▶ Therefore if W is a diagonal matrix whose i th diagonal entry is $p_i(1 - p_i)$, then

$$\text{Cov}(Z) = W^{-1}.$$

- ▶ Thus, while fitting a linear model to Z_i based on the explanatory variables, it is sensible to estimate the coefficients of the linear model by

$$(X^T W X)^{-1} X^T W Z.$$

- ▶ Thus, while fitting a linear model to Z_i based on the explanatory variables, it is sensible to estimate the coefficients of the linear model by

$$(X^T W X)^{-1} X^T W Z.$$

- ▶ This gives us the estimate of β in the logistic regression model:

$$\hat{\beta} := (X^T W X)^{-1} X^T W Z. \quad (7)$$

- ▶ Thus, while fitting a linear model to Z_i based on the explanatory variables, it is sensible to estimate the coefficients of the linear model by

$$(X^T W X)^{-1} X^T W Z.$$

- ▶ This gives us the estimate of β in the logistic regression model:

$$\hat{\beta} := (X^T W X)^{-1} X^T W Z. \quad (7)$$

- ▶ The obvious problem with the above approach is that we do not know p_i (p_i depends on the parameters β_0, \dots, β_p that we are trying to estimate) and so we cannot really compute the response variable Z_i or the matrix W .

- ▶ The natural solution to this is to use an iterative method. First have an initial estimate of β_0, \dots, β_p . Call this initial estimator $\hat{\beta}^{(0)}$. Use this estimator to calculate p_i via

$$p_i = \frac{\exp(\hat{\beta}_0^{(0)} + \hat{\beta}_1^{(0)} x_{i1} + \dots + \hat{\beta}_p^{(0)} x_{ip})}{1 + \exp(\hat{\beta}_0^{(0)} + \hat{\beta}_1^{(0)} x_{i1} + \dots + \hat{\beta}_p^{(0)} x_{ip})}.$$

- ▶ The natural solution to this is to use an iterative method. First have an initial estimate of β_0, \dots, β_p . Call this initial estimator $\hat{\beta}^{(0)}$. Use this estimator to calculate p_i via

$$p_i = \frac{\exp(\hat{\beta}_0^{(0)} + \hat{\beta}_1^{(0)} x_{i1} + \dots + \hat{\beta}_p^{(0)} x_{ip})}{1 + \exp(\hat{\beta}_0^{(0)} + \hat{\beta}_1^{(0)} x_{i1} + \dots + \hat{\beta}_p^{(0)} x_{ip})}.$$

- ▶ Use these values of p_i to create the response variable values Z_i via (6) and also use values of p_i to construct the matrix W . With Z and W , we can estimate β as in (7). Call this $\hat{\beta}^{(1)}$:

$$\hat{\beta}^{(1)} = (X^T W X)^{-1} X^T W Z.$$

- ▶ The natural solution to this is to use an iterative method. First have an initial estimate of β_0, \dots, β_p . Call this initial estimator $\hat{\beta}^{(0)}$. Use this estimator to calculate p_i via

$$p_i = \frac{\exp(\hat{\beta}_0^{(0)} + \hat{\beta}_1^{(0)} x_{i1} + \dots + \hat{\beta}_p^{(0)} x_{ip})}{1 + \exp(\hat{\beta}_0^{(0)} + \hat{\beta}_1^{(0)} x_{i1} + \dots + \hat{\beta}_p^{(0)} x_{ip})}.$$

- ▶ Use these values of p_i to create the response variable values Z_i via (6) and also use values of p_i to construct the matrix W . With Z and W , we can estimate β as in (7). Call this $\hat{\beta}^{(1)}$:

$$\hat{\beta}^{(1)} = (X^T W X)^{-1} X^T W Z.$$

- ▶ Now replace the initial estimator $\hat{\beta}^{(0)}$ by $\hat{\beta}^{(1)}$ and repeat this process. Keep repeating this until two successive estimates $\hat{\beta}^{(m)}$ and $\hat{\beta}^{(m+1)}$ do not change much. At that point, stop and report the estimate of β in the logistic regression model by $\hat{\beta}^{(m)}$.

- ▶ By what we have seen that this method is equivalent to computing the MLE by Newton's method.

Standard Errors for the MLE

- ▶ Is the MLE $\hat{\beta}$ unbiased (or at least approximately unbiased)?

Standard Errors for the MLE

- ▶ Is the MLE $\hat{\beta}$ unbiased (or at least approximately unbiased)? How do we compute its standard errors?

Standard Errors for the MLE

- ▶ Is the MLE $\hat{\beta}$ unbiased (or at least approximately unbiased)? How do we compute its standard errors? To answer these questions, consider the following simple heuristic argument.

Standard Errors for the MLE

- ▶ Is the MLE $\hat{\beta}$ unbiased (or at least approximately unbiased)? How do we compute its standard errors? To answer these questions, consider the following simple heuristic argument.
- ▶ Because $\hat{\beta}$ maximizes the loglikelihood, we have $\nabla \ell(\hat{\beta}) = 0$.

Standard Errors for the MLE

- ▶ Is the MLE $\hat{\beta}$ unbiased (or at least approximately unbiased)? How do we compute its standard errors? To answer these questions, consider the following simple heuristic argument.
- ▶ Because $\hat{\beta}$ maximizes the loglikelihood, we have $\nabla \ell(\hat{\beta}) = 0$. Let us now obtain a Taylor expansion of $\nabla \ell(\hat{\beta})$ around the true β :

$$0 = \nabla \ell(\hat{\beta}) \approx \nabla \ell(\beta) + H\ell(\beta) (\hat{\beta} - \beta).$$

Standard Errors for the MLE

- ▶ Is the MLE $\hat{\beta}$ unbiased (or at least approximately unbiased)? How do we compute its standard errors? To answer these questions, consider the following simple heuristic argument.
- ▶ Because $\hat{\beta}$ maximizes the loglikelihood, we have $\nabla \ell(\hat{\beta}) = 0$. Let us now obtain a Taylor expansion of $\nabla \ell(\hat{\beta})$ around the true β :

$$0 = \nabla \ell(\hat{\beta}) \approx \nabla \ell(\beta) + H\ell(\beta) (\hat{\beta} - \beta).$$

- ▶ Using the expressions $\nabla \ell(\beta) = X^T(Y - p)$ and $H\ell(\beta) = -X^T W X$, we obtain

$$0 \approx X^T(Y - p) - X^T W X(\hat{\beta} - \beta).$$

Standard Errors for the MLE

- ▶ Is the MLE $\hat{\beta}$ unbiased (or at least approximately unbiased)? How do we compute its standard errors? To answer these questions, consider the following simple heuristic argument.
- ▶ Because $\hat{\beta}$ maximizes the loglikelihood, we have $\nabla \ell(\hat{\beta}) = 0$. Let us now obtain a Taylor expansion of $\nabla \ell(\hat{\beta})$ around the true β :

$$0 = \nabla \ell(\hat{\beta}) \approx \nabla \ell(\beta) + H\ell(\beta) (\hat{\beta} - \beta).$$

- ▶ Using the expressions $\nabla \ell(\beta) = X^T(Y - p)$ and $H\ell(\beta) = -X^T W X$, we obtain

$$0 \approx X^T(Y - p) - X^T W X(\hat{\beta} - \beta).$$

- ▶ This gives

$$\hat{\beta} - \beta \approx (X^T W X)^{-1} X^T(Y - p).$$

- ▶ Because $\mathbb{E} Y = p$, this means that $\hat{\beta}$ is approximately unbiased for β .

- ▶ Because $\mathbb{E} Y = p$, this means that $\hat{\beta}$ is approximately unbiased for β . Also because $\text{Cov}(Y) = W$, we have

$$\text{Cov}(\hat{\beta}) \approx (X^T W X)^{-1}.$$

- ▶ Because $\mathbb{E} Y = p$, this means that $\hat{\beta}$ is approximately unbiased for β . Also because $\text{Cov}(Y) = W$, we have

$$\text{Cov}(\hat{\beta}) \approx (X^T W X)^{-1}. \quad \text{💬}$$

- ▶ Therefore the approximate standard error of $\hat{\beta}_j$ is obtained by the square root of the corresponding diagonal entry of $(X^T W X)^{-1}$.