

1

```
shoeSize <- c(6.5, 9.0, 8.5, 8.5, 10.5, 7.0, 9.5, 9.0, 13.0, 7.5, 10.5, 8.5, 12.0, 10.5, 13.0, 11.5,
8.5,5.0,10.0,6.5,7.5,8.5,10.5,8.5,10.5,11.0,9.0,13.0)
height <- c(66.0, 68.0, 64.5, 65.0, 70.0,64.0,70.0,71.0,72.0,64.0,74.5,67.0,71.0,71.0,77.0,72.0,59.0,62.0,72.0,66.0,64.0,67.0,73.0,69.0,
gender <- c("F", "F", "F", "F", "M", "F", "F", "F", "M", "F", "M", "F", "M", "M", "M", "M", "F", "F",
"M", "F", "F", "M", "M", "F", "M", "M", "M", "M")
householdData <- data.frame( ShoeSize = shoeSize, Height = height, Gender = gender )
householdData
```

1.a

In the data, there are three variables which are the shoe size, height, and gender. There are 28 observations

1.b

```
males <- householdData[householdData$Gender == "M",] males
females <- householdData[householdData$Gender == "F",] females
```

1.c

```
meanOfShoeSize <- mean(householdData$ShoeSize) meanOfShoeSize
meanOfHeight <- mean(householdData$Height) meanOfHeight
```

1.d

The relationship of the two is that the shoe size is directly proportional to the height. If the height is small, the shoe size is also small.

#—————

2

```
months_vector <- c("March", "April", "January", "November", "January", "September", "October", "September", "November", "Au
months_vector
factor_months_vector <- factor(months_vector)
factor_months_vector
```

3

```
summary(months_vector) summary(factor_months_vector)
```

In the summary of months_vector, it shows the number of observations, class, and mode of the vector.

In the summary of factor_months_vector, it shows the frequency of each months.

Both are useful in different cases where the no. of observations, class, mode, or the frequency is needed.

4

```
factor_data <- c("East", "West", "North") factor_frequency <- c(1,4,3)
new_order_data <- factor(factor_data,levels = c("East","West","North"))
print(new_order_data)
```

5

```
imported_table <- read.table(file = "/cloud/project/import_march.csv", header = TRUE, sep = ",")
imported_table
```

6

```
randomNum <- readline(prompt = "Enter number from 1 to 50:")
#cant knit if there is as.numeric #randomNum <- as.numeric(randomNum)
paste("The number you have chosen is", randomNum)
if (randomNum > 50) { paste("The number selected is beyond the range of 1 to 50") } else if (randomNum
== 20) { paste("TRUE") } else { paste(randomNum) }
```

7

```
minimumBills <- function(price) {
```

```
minBills <- price %/% 50 paste("The minimum no. of bills:", minBills) }
minimumBills(90)
```

8.a

```
names <- c("Annie", "Thea", "Steve", "Hanna") grade1 <- c(85,65,75,95) grade2 <- c(65,75,55,75) grade3 <-
c(85,90,80,100) grade4 <- c(100,90,85,90)

mathScore <- data.frame( Name = names, Grade1 = grade1, Grade2 = grade2, Grade3 = grade3, Grade4 =
grade4 )
```

8.b

```
mathScoreAverage <- -(mathScoreGrade1 + mathScoreGrade2 + mathScoreGrade3 + mathScore$Grade4)
/ 4

highscorers <- mathScore[mathScore$Average > 90,] highscorers

if (nrow(highscorers) > 0) { paste(highscorers$Name, "'saveragegradethissemesteris", highscorers$Average)
} else { paste("No students have an average math score over 90.") }
```

8.c

```
firstTest <- sum(mathScore$Grade1) / nrow(mathScore) firstTest
secondTest <- sum(mathScore$Grade2) / nrow(mathScore) secondTest
thirdTest <- sum(mathScore$Grade3) / nrow(mathScore) thirdTest
fourthTest <- sum(mathScore$Grade4) / nrow(mathScore) fourthTest

if (firstTest < 80) { paste("The 1st test was difficult.") } else if(secondTest < 80) { paste("The 2nd test
was difficult.") } else if(thirdTest < 80) { paste("The 3rd test was difficult.") } else if(fourthTest < 80) {
paste("The 4th test was difficult.") } else { paste("No test had an average score less than 80.") }
```

8.d

annie scores

```
if (mathScore[1,2] > mathScore[1,3] && mathScore[1,2] > mathScore[1,4] && mathScore[1,2] > math-
Score[1,5]) { annieHighest <- mathScore[1,2] } else if (mathScore[1,3] > mathScore[1,4] && mathScore[1,3]
> mathScore[1,5]) { annieHighest <- mathScore[1,3] } else if (mathScore[1,4] > mathScore[1,5] && math-
Score[1,2] > mathScore[1,5]) { annieHighest <- mathScore[1,4] } else { annieHighest <- mathScore[1,5]
}
```

thea scores

```
if (mathScore[2,2] > mathScore[2,3] && mathScore[2,2] > mathScore[2,4] && mathScore[2,2] > math-
Score[2,5]) { theaHighest <- mathScore[2,2] } else if (mathScore[2,3] > mathScore[2,4] && mathScore[2,3] >
mathScore[2,5]) { theaHighest <- mathScore[2,3] } else if (mathScore[2,4] > mathScore[2,5] && mathScore[2,2]
> mathScore[2,5]) { theaHighest <- mathScore[2,4] } else { theaHighest <- mathScore[2,5] }
```

steve scores

```
if (mathScore[3,2] > mathScore[3,3] && mathScore[3,2] > mathScore[3,4] && mathScore[3,2] > mathScore[3,5]) { steveHighest <- mathScore[3,2] } else if (mathScore[3,3] > mathScore[3,4] && mathScore[3,3] > mathScore[3,5]) { steveHighest <- mathScore[3,3] } else if (mathScore[3,4] > mathScore[3,5] && mathScore[3,4] > mathScore[3,2]) { steveHighest <- mathScore[3,4] } else { steveHighest <- mathScore[3,5] }
```

hanna scores

```
if (mathScore[4,2] > mathScore[4,3] && mathScore[4,2] > mathScore[4,4] && mathScore[4,2] > mathScore[4,5]) { hannaHighest <- mathScore[4,2] } else if (mathScore[4,3] > mathScore[4,4] && mathScore[4,3] > mathScore[4,5]) { hannaHighest <- mathScore[4,3] } else if (mathScore[4,4] > mathScore[4,5] && mathScore[4,4] > mathScore[4,2]) { hannaHighest <- mathScore[4,4] } else { hannaHighest <- mathScore[4,5] }
```

```
mathScore$HighestGrades <- c(annieHighest, theaHighest, steveHighest, hannaHighest)
```

```
above90 <- mathScore[mathScore$HighestGrades > 90,] above90
```

```
if (nrow(above90) > 0) { paste(above90Name, "shighestgradethissemesteris", above90HighestGrade) } else { paste("No students have an average math score over 90.") }
```