Supplementary Table S1. Univariable and Multivariable Cox Regression Analyses of the Risk Group Defined by the Proposed Model

Variable	Univariable		Multivariable	
	HR (95% CI)	*P*	HR (95% CI)	*P*
A. Overall survival (n = 238, 199 deaths)				
High-risk group (vs. low risk)	2.32 (1.74–3.10)	< 0.001	1.73 (1.27–2.36)	< 0.001
Extensive stage (vs. limited)	2.39 (1.71–3.32)	< 0.001	1.77 (1.23–2.54)	0.002
ECOG performance status ≥ 2 (vs. 0–1)	1.42 (0.94–2.15)	0.095	1.02 (0.66–1.58)	0.921
Age, per 10 years	1.59 (1.31–1.92)	< 0.001	1.67 (1.37–2.05)	< 0.001
Female sex (vs. male)	1.08 (0.73–1.60)	0.713	0.97 (0.65–1.46)	0.896
Liver metastasis	2.70 (1.94–3.74)	< 0.001	2.00 (1.41–2.84)	< 0.001
Brain metastasis at diagnosis	1.81 (1.30–2.54)	< 0.001	1.59 (1.11–2.26)	0.011
B. Progression-free survival (n = 238, 212 events)				
High-risk group (vs. low risk)	2.03 (1.54–2.67)	< 0.001	1.73 (1.29–2.33)	< 0.001
Extensive stage (vs. limited)	2.09 (1.54–2.85)	< 0.001	1.73 (1.24–2.43)	0.001
ECOG performance status ≥ 2 (vs. 0–1)	1.20 (0.79–1.81)	0.390	1.08 (0.70–1.66)	0.722
Age, per 10 years	1.23 (1.03–1.46)	0.022	1.17 (0.98–1.40)	0.078
Female sex (vs. male)	1.08 (0.73–1.59)	0.702	0.82 (0.54–1.24)	0.348
Liver metastasis	2.27 (1.65–3.13)	< 0.001	1.65 (1.17–2.34)	0.005
Brain metastasis at diagnosis	1.85 (1.33–2.58)	< 0.001	1.56 (1.10–2.20)	0.012

Risk group as in Table 5. Because the model already uses these clinical variables, the multivariable HR shows added prognostic information, not independence from them. CI = confidence interval, ECOG = Eastern Cooperative Oncology Group, HR = hazard ratio