

Supplementary Table 15. Blood Glucose During ArgTT Compared to Fasted					
Time Post-Arginine (min)	Difference in Blood Glucose (mM)	95% Credibility Interval (mM)		Strength of Evidence	
		Lower Bound	Upper Bound		
Female - FP					
7	1.1	0.3	1.9	Very strong	
15	-1.6	-2.6	-0.6	Decisive	
30	-3.0	-4.0	-2.0	Decisive	
60	-3.1	-4.2	-2.1	Decisive	
90	-1.6	-2.6	-0.6	Decisive	
Female - KC					
7	1.3	0.5	2.1	Decisive	
30	-1.6	-2.6	-0.6	Decisive	
60	-1.4	-2.5	-0.3	Very strong	
Female - TC					
30	-2.4	-3.4	-1.3	Decisive	
60	-3.1	-4.3	-2.0	Decisive	
90	-1.9	-3.0	-0.8	Decisive	
120	-1.1	-2.0	-0.2	Very strong	
Male - FP					
7	1.8	1.0	2.6	Decisive	
15	-1.3	-2.2	-0.3	Very strong	
30	-2.5	-3.5	-1.5	Decisive	
60	-2.7	-3.8	-1.5	Decisive	
90	-1.3	-2.3	-0.2	Very strong	
Male - KC					
7	1.6	0.7	2.4	Decisive	
15	-1.9	-2.9	-0.8	Decisive	
30	-3.7	-4.8	-2.6	Decisive	
60	-3.9	-5.1	-2.7	Decisive	
90	-2.7	-3.8	-1.6	Decisive	
120	-2.0	-2.9	-1.0	Decisive	
Male - TC					
15	-1.4	-2.6	-0.3	Very strong	
30	-2.9	-4.1	-1.8	Decisive	
60	-3.6	-4.9	-2.3	Decisive	
90	-1.9	-3.1	-0.7	Decisive	
120	-1.4	-2.4	-0.4	Decisive	
ArgTT=Arginine Tolerance Test. If no result is reported, the posterior probability did not exceed 95%.					