Ouppien	. 95% Credibility Interval (nM)				area to rastea
Weeks Post-Implant	Time Post-Glucose (min)	Difference in C-peptide (nM)		· · ·	Strength of Evidence
Female - FP		,			
Week 16	60	0.2	0.0	0.4	Strong
Week 16	90	0.2	0.0	0.4	Strong
Week 16	120	0.3	0.1		Very strong
Week 21	30	0.2	0.0	0.4	Strong
Week 21	60	0.4	0.2		Decisive
Week 21	90	0.4	0.3	0.6	Decisive
Week 21	120	0.5	0.3		Decisive
Female - KC					
Week 12	60	0.2	0.0	0.3	Strong
Week 16	30	0.3	0.1		Decisive
Week 16	60	0.4	0.3		Decisive
Week 16	90	0.4	0.3		Decisive
Week 16	120	0.4	0.3		Decisive
Week 10	30	0.7	0.6		Decisive
Week 21	60	0.6	0.5		Decisive
Week 21	90	0.3	0.3		Decisive
Week 21	120	0.3	0.2		Decisive
	120	0.3	0.2	0.4	Decisive
Female - TC		0.4	0.0	0.7	Otrono
Week 8	60	0.4	0.0		Strong
Week 10	60	0.7	0.3		Decisive
Week 12	30	0.5	0.1		Very strong
Week 12	60	0.8	0.5		Decisive
Week 16	30	0.8	0.4		Decisive
Week 16	60	0.6	0.3		Decisive
Week 16	90	0.4	0.1		Very strong
Week 21	30	1.8	1.5		Decisive
Week 21	60	1.0	0.8		Decisive
Week 21	90	0.5	0.2		Decisive
Week 21	120	0.6	0.3	0.9	Decisive
Male - FP					
Week 16	90	0.3	0.0		Strong
Week 21	60	0.4	0.2	0.6	Decisive
Week 21	90	0.4	0.2	0.7	Decisive
Week 21	120	0.4	0.2	0.7	Decisive
Male - KC					
Week 16	60	0.3	0.0	0.5	Strong
Week 16	90	0.3	0.1	0.6	Very strong
Week 16	120	0.4	0.1	0.6	Very strong
Week 21	30	0.3	0.1	0.5	Decisive
Week 21	60	0.5	0.3	0.7	Decisive
Week 21	90	0.5	0.3	0.8	Decisive
Week 21	120	0.4	0.2	0.7	Decisive
Male - TC					
Week 10	60	0.7	0.3	1.2	Decisive
Week 12	60	1.1	0.7	1.5	Decisive
Week 16	30	0.4	0.0	0.9	Strong
Week 16	60	0.6	0.2	1.1	Decisive
Week 16	90	0.5	0.1	1.0	Very strong
Week 21	30	1.2	0.8	1.6	Decisive
Week 21	60	0.7	0.3	1.2	Decisive
Week 21	120	0.5	0.1	1.0	Strong
If no result is rep	ported, the posterior prob	pability did not exce	ed 95%.		

Supplementary Table 7. Glucose-Stimulated C-peptide Compared to Fasted