



# Calculation of predictions for non-identical particle correlations in AA collisions at LHC energies from hydrodynamics-inspired models

MASTER OF SCIENCE THESIS

Author:

**Mateusz Wojciech Gałążyn**

Supervisor:

**Prof. Adam Kisiel**

Warsaw, 25th July 2014



# Obliczenia teoretycznych przewidywań korelacji cząstek nieidentycznych w zderzeniach AA przy energiach LHC pochodzących z modeli hydrodynamicznych

PRACA MAGISTERSKA

Autor:

**Mateusz Wojciech Gałążyn**

Promotor:

**dr hab. inż. Adam Kisiel, prof. PW**

Warszawa, 25 lipca 2014

## **Abstract**

## **Streszczenie**

# Contents

4	<b>1 Theory of heavy ion collisions</b>	<b>2</b>
5	1.1 The Standard Model . . . . .	2
6	1.2 Quantum Chromodynamics . . . . .	3
7	1.3 Relativistic heavy ion collisions . . . . .	3
8	<b>2 Therminator model</b>	<b>4</b>
9	2.1 (3+1)-dimensional viscous hydrodynamics . . . . .	4
10	2.2 Statistical hadronization . . . . .	5
11	2.2.1 Cooper-Frye formalism . . . . .	6
12	<b>3 Particle interferometry</b>	<b>8</b>
13	3.1 HBT interferometry . . . . .	8
14	3.2 Intensity interferometry in heavy ion collisions . . . . .	8
15	3.2.1 Theoretical approach . . . . .	8
16	3.2.2 Experimental approach . . . . .	8
17	3.3 Scaling of femtoscopic radii . . . . .	8
18	<b>4 Results</b>	<b>9</b>
19	4.1 Identical particles correlations . . . . .	9
20	4.2 Results of the fit . . . . .	9
21	4.3 Discussion of results . . . . .	9
22	<b>5 Summary</b>	<b>10</b>

# <sup>23</sup> Introduction

# Chapter 1

## Theory of heavy ion collisions

### 1.1 The Standard Model

In the 1970s, a new theory of fundamental particles and their interaction emerged. A new concept, which concerns the electromagnetic, weak and strong nuclear interactions between know particles. This theory is called *The Standard Model*. There are seventeen named particles in the standard model, organized into the chart shown below (Fig. 1.1). Fundamental particles are divided into two families: *fermions* and *bosons*.




Figure 1.1: The Standard Model of elementary particles [1].

32 Fermions are the building blocks of matter. They are divided into two groups.  
 33 Six of them, which must bind together are called *quarks*. Quarks are known to  
 34 bind into doublets (*mesons*), triplets (*baryons*) and recently confirmed four-quark  
 35 states.<sup>1</sup> Two of baryons, with the longest lifetimes, are forming a nucleus: a pro-  
 36 ton and a neutron. A proton is build from two up quarks and one down, and  
 37 neutron consists of two down quarks and one up. A proton is found to be a stable  
 38 particle (at least it has a lifetime larger than  $10^{35}$  years) and a free neutron has a  
 39 mean lifetime about  $8.8 \times 10^2$  s. Fermions, that can exist independently are called  
 40 *leptons*. Neutrinos are a subgroup of leptons, which are only influenced by weak  
 41 interaction. Fermions can be divided into three generations (three columns in  
 42 the Figure 1.1). Generation I particles can combine into hadrons with the longest  
 43 life spans. Generation II and III consists of unstable particles which form also  
 44 unstable hadrons.

45 Bosons are force carriers. There are four fundamental forces: weak - respons-  
 46 ible for radioactive decay, strong - coupling quarks into hadrons, electromagnetic  
 47 - between charged particles and gravity - the weakest, which causes the attraction  
 48 between particles with a mass. The Standard Model describes the first three. The  
 49 weak force is mediated by  $W^\pm$  and  $Z^0$  bosons, electromagnetic force is carried by  
 50 photons  $\gamma$  and the carriers of a strong interaction are gluons  $g$ . The fifth boson is  
 51 a Higgs boson which is responsible for giving other particles mass.

## 52 1.2 Quantum Chromodynamics

## 53 1.3 Relativistic heavy ion collisions

---

<sup>1</sup>The LHCb experiment at CERN in Geneva confirmed recently existence of  $Z(4430)$  - a particle consisting of four quarks [2].



## 54 Chapter 2

# 55 Terminator model

56 THERMINATOR [3] is a Monte Carlo event generator designed to investigate  
57 the particle production in the relativistic heavy ion collisions. The functionality  
58 of the code includes a generation of the stable particles and unstable resonances  
59 at the chosen hypersurface model. It performs the statistical hadronization which  
60 is followed by space-time evolution of particles and the decay of resonances. The  
61 key element of this method is an inclusion of a complete list of hadronic reson-  
62 ances, which contribute very significantly to the observables. The second version  
63 of THERMINATOR [4] comes with a possibility to incorporate any shape of freeze-  
64 out hypersurface and the expansion velocity field, especially those generated ex-  
65 ternally with various hydrodynamic codes.

## 66 2.1 (3+1)-dimensional viscous hydrodynamics

67 Most of the relativistic viscous hydrodynamic calculations are done in  
68 (2+1)-dimensions. Such simplification assumes boost-invariance of a matter  
69 created in a collision. Experimental data reveals that no boost-invariant region is  
70 formed in the collisions [5]. Hence, for the better description of created system a  
71 (3+1)-dimensional model is required.

72 In the four dimensional relativistic dynamics one can describe a system  
73 using a space-time four-vector  $x^\nu = (ct, x, y, z)$ , a velocity four-vector  
74  $u^\nu = \gamma(c, v_x, v_y, v_z)$  and a energy-momentum tensor  $T^{\mu\nu}$ . The particular  
75 components of  $T^{\mu\nu}$  have a following meaning:

- 76 •  $T^{00}$  - an energy density,
- 77 •  $cT^{0\alpha}$  - an energy flux across a surface  $x^\alpha$ ,
- 78 •  $T^{\alpha 0}$  - an  $\alpha$ -momentum flux across a surface  $x^\alpha$  multiplied by  $c$ ,
- 79 •  $T^{\alpha\beta}$  - components of momentum flux density tensor,

where  $\gamma = (1 - v^2/c^2)^{-1/2}$  is Lorentz factor and  $\alpha, \beta \in \{1, 2, 3\}$ . Using  $u^\nu$  one can express  $T^{\mu\nu}$  as follows [6]:

$$T_0^{\mu\nu} = (e + p)u^\mu u^\nu - pg^{\mu\nu} \quad (2.1)$$

where  $e$  is an energy density,  $p$  is a pressure and  $g^{\mu\nu}$  is an inverse metric tensor:

$$g^{\mu\nu} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}. \quad (2.2)$$

The presented version of energy-momentum tensor (2.1) can be used to describe dynamics of a perfect fluid. To take into account influence of viscosity, one has to apply the following corrections coming from shear  $\pi^{\mu\nu}$  and bulk  $\Pi$  viscosities [7]:

$$T^{\mu\nu} = T_0^{\mu\nu} + \pi^{\mu\nu} + \Pi(g^{\mu\nu} - u^\mu u^\nu). \quad (2.3)$$

The stress tensor  $\pi^{\mu\nu}$  and the bulk viscosity  $\Pi$  are solutions of dynamical equations in the second order viscous hydrodynamic framework [6]. The comparison of hydrodynamics calculations with the experimental results reveal, that the shear viscosity divided by entropy  $\eta/s$  has to be small and close to the AdS/CFT estimate  $\eta/s = 0.08$  [7, 8].

When using  $T^{\mu\nu}$  to describe system evolving close to local thermodynamic equilibrium, relativistic hydrodynamic equations in a form of:

$$\partial_\mu T^{\mu\nu} = 0 \quad (2.4)$$

can be used to describe the dynamics of the local energy density, pressure and flow velocity.

Hydrodynamic calculations are starting from the Glauber<sup>1</sup> model initial conditions. The collective expansion of a fluid ends at the freeze-out hypersurface. That surface is usually defined as a constant temperature surface, or equivalently as a cut-off in local energy density. The freeze-out is assumed to occur at the temperature  $T = 140$  MeV.

## 2.2 Statistical hadronization

Statistical description of heavy ion collision has been successfully used to describe quantitatively *soft* physics, i.e. the regime with the transverse momentum not exceeding 2 GeV. The basic assumption of the statistical approach of evolution of the quark-gluon plasma is that at some point of the space-time evolution of the fireball, the thermal equilibrium is reached. When

<sup>1</sup>The Glauber Model is used to calculate “geometrical” parameters of a collision like an impact parameter, number of participating nucleons or number of binary collisions.

the system is in the thermal equilibrium the local phase-space densities of particles follow the Fermi-Dirac or Bose-Einstein statistical distributions. At the end of the plasma expansion, the freeze-out occurs. The freeze-out model incorporated in the THERMINATOR model assumes, that chemical and thermal freeze-out occur at the same time.

### 2.2.1 Cooper-Frye formalism

The result of the hydrodynamic calculations is the freeze-out hypersurface  $\Sigma^\mu$ . A three-dimensional element of the surface is defined as [4]

$$d\Sigma_\mu = \epsilon_{\mu\alpha\beta\gamma} \frac{\partial x^\alpha}{\partial \alpha} \frac{\partial x^\beta}{\partial \beta} \frac{\partial x^\gamma}{\partial \gamma} d\alpha d\beta d\gamma, \quad (2.5)$$

where  $\epsilon_{\mu\alpha\beta\gamma}$  is the Levi-Civita tensor and the variables  $\alpha, \beta, \gamma \in \{1, 2, 3\}$  are used to parametrize the three-dimensional freeze-out hypersurface in the Minkowski four-dimensional space. The Levi-Civita tensor is equal to 1 when the indices form an even permutation (eg.  $\epsilon_{0123}$ ), to -1 when the permutation is odd (e.g.  $\epsilon_{2134}$ ) and has a value of 0 if any index is repeated. Therefore [4],

$$d\Sigma_0 = \begin{vmatrix} \frac{\partial x}{\partial \alpha} & \frac{\partial x}{\partial \beta} & \frac{\partial x}{\partial \gamma} \\ \frac{\partial y}{\partial \alpha} & \frac{\partial y}{\partial \beta} & \frac{\partial y}{\partial \gamma} \\ \frac{\partial z}{\partial \alpha} & \frac{\partial z}{\partial \beta} & \frac{\partial z}{\partial \gamma} \end{vmatrix} d\alpha d\beta d\gamma \quad (2.6)$$

and the remaining components are obtained by cyclic permutations of  $t, x, y$  and  $z$ .

One can obtain the number of hadrons produced on the hypersurface  $\Sigma^\mu$  from the Cooper-Frye formalism. The following integral yields the total number of created particles [4]:

$$N = (2s + 1) \int \frac{d^3p}{(2\pi)^3 E_p} \int d\Sigma_\mu(x) p^\mu f(x, p), \quad (2.7)$$

where

$$f(p \cdot u) = \left\{ \exp \left[ \frac{p_\mu u^\mu - (B\mu_B + I_3\mu_{I_3} + S\mu_S + C\mu_C)}{T} \right] \pm 1 \right\}^{-1} \quad (2.8)$$

is the phase-space distribution for particles (for stable ones and resonances). For the Fermi-Dirac distribution in the 2.8 there is a plus sign and for Bose-Einstein statistics minus sign respectively. The thermodynamic quantities appearing in the  $f(\cdot)$  are  $T$  - temperature,  $\mu_B$  - baryon chemical potential,  $\mu_{I_3}$  - isospin chemical potential,  $\mu_S$  - strange chemical potential,  $\mu_C$  - charmed chemical potential and the  $s$  is a spin of a particle. One can simply derive from equation 2.7, the dependence of the momentum density [9]:

$$E \frac{dN}{d^3p} = \int f(x, p) p^\mu d\Sigma_\mu. \quad (2.9)$$

134 The equations presented above are directly used in the THERMINATOR to generate  
135 the hadrons with the Monte-Carlo method.

## **Chapter 3**

# **Particle interferometry**

### **3.1 HBT interferometry**

### **3.2 Intensity interferometry in heavy ion collisions**

#### **3.2.1 Theoretical approach**

**Two particle wave function**

**Source function**

**Theoretical correlation function**

**Spherical harmonics decomposition of correlation function**

#### **3.2.2 Experimental approach**

### **3.3 Scaling of femtoscopic radii**

## 147 **Chapter 4**

# 148 **Results**

### 149 **4.1 Identical particles correlations**

### 150 **4.2 Results of the fit**

### 151 **4.3 Discussion of results**

<sup>152</sup> **Chapter 5**

<sup>153</sup> **Summary**

# Bibliography

- [1] Standard Model of Elementary Particles - Wikipedia, the free encyclopedia  
[http://en.wikipedia.org/wiki/standard\\_model](http://en.wikipedia.org/wiki/standard_model).
- [2] R. Aaij et al. (LHCb Collaboration). Observation of the resonant character of the  $z(4430)^-$  state. *Phys. Rev. Lett.*, 112:222002, Jun 2014.
- [3] Adam Kisiel, Tomasz Taluc, Wojciech Broniowski, and Wojciech Florkowski. THERMINATOR: THERMal heavy-IoN generATOR. *Comput.Phys.Commun.*, 174:669–687, 2006.
- [4] Mikolaj Chojnacki, Adam Kisiel, Wojciech Florkowski, and Wojciech Broniowski. THERMINATOR 2: THERMal heavy IoN generATOR 2. *Comput.Phys.Commun.*, 183:746–773, 2012.
- [5] I. et al (BRAHMS Collaboration) Bearden. Charged meson rapidity distributions in central Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. *Phys. Rev. Lett.*, 94:162301, Apr 2005.
- [6] W. Israel and J.M. Stewart. Transient relativistic thermodynamics and kinetic theory. *Annals of Physics*, 118(2):341 – 372, 1979.
- [7] Piotr Bożek. Flow and interferometry in (3 + 1)-dimensional viscous hydrodynamics. *Phys. Rev. C*, 85:034901, Mar 2012.
- [8] K. Kovtun, P. D. T. Son, and A. O. Starinets. Viscosity in strongly interacting quantum field theories from black hole physics. *Phys. Rev. Lett.*, 94:111601, Mar 2005.
- [9] Fred Cooper and Graham Frye. Single-particle distribution in the hydrodynamic and statistical thermodynamic models of multiparticle production. *Phys. Rev. D*, 10:186–189, Jul 1974.



# 178 List of Figures

<small>179</small>	1.1 The Standard Model of elementary particles [1]. . . . .	2
--------------------	---	---