



# Calculation of predictions for non-identical particle correlations in AA collisions at LHC energies from hydrodynamics-inspired models

MASTER OF SCIENCE THESIS

Author:

**Mateusz Wojciech Gałążyn**

Supervisor:

**Prof. Adam Kisiel**

Warsaw, 31st August 2014



# Obliczenia teoretycznych przewidywań korelacji cząstek nieidentycznych w zderzeniach AA przy energiach LHC pochodzących z modeli hydrodynamicznych

PRACA MAGISTERSKA

Autor:

**Mateusz Wojciech Gałążyn**

Promotor:

**dr hab. inż. Adam Kisiel, prof. PW**

Warszawa, 31 sierpnia 2014

## **Abstract**

## **Streszczenie**

# Contents

4	<b>1 Theory of heavy ion collisions</b>	<b>2</b>
5	1.1 The Standard Model . . . . .	2
6	1.2 Quantum Chromodynamics . . . . .	3
7	1.2.1 Quarks and gluons . . . . .	3
8	1.2.2 Quantum Chromodynamics potential . . . . .	4
9	1.2.3 The quark-gluon plasma . . . . .	6
10	1.3 Relativistic heavy ion collisions . . . . .	7
11	1.3.1 Stages of heavy ion collision . . . . .	7
12	1.3.2 QGP signatures . . . . .	9
13	<b>2 Therminator model</b>	<b>17</b>
14	2.1 (3+1)-dimensional viscous hydrodynamics . . . . .	17
15	2.2 Statistical hadronization . . . . .	18
16	2.2.1 Cooper-Frye formalism . . . . .	19
17	<b>3 Particle interferometry</b>	<b>21</b>
18	3.1 HBT interferometry . . . . .	21
19	3.2 Theoretical approach . . . . .	22
20	3.2.1 Conventions used . . . . .	22
21	3.2.2 Two particle wave function . . . . .	23
22	3.2.3 Source emission function . . . . .	24
23	3.2.4 Theoretical correlation function . . . . .	26
24	3.2.5 Spherical harmonics decomposition of a correlation function	27
25	3.3 Experimental approach . . . . .	27
26	3.4 Scaling of femtoscopic radii . . . . .	27
27	<b>4 Results</b>	<b>28</b>
28	4.1 Identical particles correlations . . . . .	28
29	4.2 Results of the fit . . . . .	28
30	4.3 Discussion of results . . . . .	28
31	<b>5 Summary</b>	<b>29</b>

# <sup>32</sup> Introduction

# Chapter 1

## Theory of heavy ion collisions

### 1.1 The Standard Model

In the 1970s, a new theory of fundamental particles and their interaction emerged. A new concept, which concerns the electromagnetic, weak and strong nuclear interactions between know particles. This theory is called *The Standard Model*. There are seventeen named particles in the standard model, organized into the chart shown below (Fig. 1.1). Fundamental particles are divided into two families: *fermions* and *bosons*.



Figure 1.1: The Standard Model of elementary particles [1].

41 Fermions are the building blocks of matter. They are divided into two groups.  
 42 Six of them, which must bind together are called *quarks*. Quarks are known to  
 43 bind into doublets (*mesons*), triplets (*baryons*) and recently confirmed four-quark  
 44 states.<sup>1</sup> Two of baryons, with the longest lifetimes, are forming a nucleus: a pro-  
 45 ton and a neutron. A proton is build from two up quarks and one down, and  
 46 neutron consists of two down quarks and one up. A proton is found to be a stable  
 47 particle (at least it has a lifetime larger than  $10^{35}$  years) and a free neutron has a  
 48 mean lifetime about  $8.8 \times 10^2$  s. Fermions, that can exist independently are called  
 49 *leptons*. Neutrinos are a subgroup of leptons, which are only influenced by weak  
 50 interaction. Fermions can be divided into three generations (three columns in  
 51 the Figure 1.1). Generation I particles can combine into hadrons with the longest  
 52 life spans. Generation II and III consists of unstable particles which form also  
 53 unstable hadrons.

54 Bosons are force carriers. There are four fundamental forces: weak - respons-  
 55 ible for radioactive decay, strong - coupling quarks into hadrons, electromagnetic  
 56 - between charged particles and gravity - the weakest, which causes the attraction  
 57 between particles with a mass. The Standard Model describes the first three. The  
 58 weak force is mediated by  $W^\pm$  and  $Z^0$  bosons, electromagnetic force is carried by  
 59 photons  $\gamma$  and the carriers of a strong interaction are gluons  $g$ . The fifth boson is  
 60 a Higgs boson which is responsible for giving other particles mass.

## 61 1.2 Quantum Chromodynamics

### 62 1.2.1 Quarks and gluons

63 Quarks interact with each other through the strong interaction. The mediator  
 64 of this force is a *gluon* - a massless and chargeless particle. In the quantum chro-  
 65 modynamics (QCD) - theory describing strong interaction - there are six types of  
 66 "charges" (like electrical charges in the electrodynamics) called *colours*. The col-  
 67 ours were introduced because some of the observed particles, like  $\Delta^-$ ,  $\Delta^{++}$  and  
 68  $\Omega^-$  appeared to consist of three quarks with the same flavour (*ddd*, *uuu* and *sss*  
 69 respectively), which was in conflict with the Pauli principle. One quark can carry  
 70 one of the three colours (usually called *red*, *green* and *blue*) and antiquark one of  
 71 the three anti-colours respectively. Only colour-neutral (or white) particles could  
 72 exist. Mesons are assumed to be a colour-anticolour pair, while baryons are *red-*  
 73 *green-blue* triplets. Gluons also are colour-charged and there are 8 types of gluons.  
 74 Therefore they can interact with themselves [3].

---

<sup>1</sup>The LHCb experiment at CERN in Geneva confirmed recently existence of  $Z(4430)$  - a particle consisting of four quarks [2].



## 1.2.2 Quantum Chromodynamics potential

As a result of that gluons are massless, one can expect, that the static potential in the QCD will have the similar form like one in the electrodynamics e.g.  $\sim 1/r$  (through an analogy to photons). In reality the QCD potential is assumed to have the form of [3]

$$V_s = -\frac{4}{3} \frac{\alpha_s}{r} + kr, \quad (1.1)$$

where the  $\alpha_s$  is a coupling constant of the strong force and the  $kr$  part is related with the *confinement*. In comparison to the electromagnetic force, a value of the strong coupling constant is  $\alpha_s \approx 1$  and the electromagnetic one is  $\alpha = 1/137$ .

The fact that quarks does not exist separately, but they are always bound, is called a confinement. As two quarks are pulled apart, the linear part  $kr$  in the Eq. 1.1 becomes dominant and the potential becomes proportional to the distance. This situation resembles stretching of a string. At some point, when the string is so large it is energetically favourable to create a quark-antiquark pair. At this moment such pair (or pairs) is formed, the string breaks and the confinement is preserved (Fig. 1.2).




Figure 1.2: A string break and a creation of a pair quark-anti-quark [4].

On the other hand, for the small  $r$ , an interaction between the quarks and gluons is dominated by the Coulomb-like term  $-\frac{4}{3} \frac{\alpha_s}{r}$ . The coupling constant  $\alpha_s$  depends on the four-momentum  $Q^2$  transferred in the interaction. This dependence is presented in Fig. 1.3. The value  $\alpha_s$  decreases with increasing momentum transfer and the interaction becomes weak for large  $Q^2$ , i.e.  $\alpha_s(Q) \rightarrow 0$ . Because of weakening of coupling constant, quarks at large energies (or small distances) are starting to behave like free particles. This phenomenon is known as an *asymptotic freedom*. The QCD potential has also temperature dependence - the force strength “melts” with the temperature increase. Therefore the asymptotic freedom is expected to appear in either the case of high baryon densities (small distances between quarks) or very high temperatures. This temperature dependence is illustrated in the Fig. 1.4.

If the coupling constant  $\alpha_s$  is small, one can use perturbative methods to calculate physical observables. Perturbative QCD (pQCD) successfully describes hard processes (with large  $Q^2$ ), such as jet production in high energy proton-antiproton collisions. The applicability of pQCD is defined by the *scale parameter*



Figure 1.3: The coupling parameter  $\alpha_s$  dependence on four-momentum transfer  $Q^2$  [5].




Figure 1.4: The QCD potential for a pair quark-antiquark as a function of distance for different temperatures. A value of a potential decreases with the temperature [4].

106  $\Lambda_{QCD} \approx 200$  MeV. If  $Q \gg \Lambda_{QCD}$  then the process is in the perturbative domain  
 107 and can be described by pQCD. A description of soft processes (when  $Q < 1$  GeV)  
 108 is a problem in QCD - perturbative theory breaks down at this scale. Therefore,  
 109 to describe processes with low  $Q^2$ , one has to use alternative methods like Lattice  
 110 QCD. Lattice QCD (LQCD) is non-perturbative implementation of a field theory  
 111 in which QCD quantities are calculated on a discrete space-time grid. LQCD al-

lows to obtain properties of matter in equilibrium, but there are some limitations. Lattice QCD requires fine lattice spacing to obtain precise results - therefore large computational resources are necessary. With the constant growth of computing power this problem will become less important. The second problem is that lattice simulations are possible only for baryon density  $\mu_B = 0$ . At  $\mu_B \neq 0$ , Lattice QCD breaks down because of the sign problem [6].

### 1.2.3 The quark-gluon plasma

The new state of matter in which quarks are no longer confined is known as a *quark-gluon plasma* (QGP). The predictions coming from the discrete space-time Lattice QCD calculations reveal a phase transition from the hadronic matter to the quark-gluon plasma at the high temperatures and baryon densities. The res-




Figure 1.5: A number of degrees of freedom as a function of a temperature [7].

ults obtained from such calculations are shown on Fig. 1.5. The energy density  $\epsilon$  which is divided by  $T^4$  is a measure of number of degrees of freedom in the system. One can observe significant rise of this value, when the temperature increases past the critical value  $T_C$ . Such increase is signaling a phase transition - the formation of QGP [8]. The values of the energy densities plotted in Fig. 1.5 do not reach the Stefan-Boltzmann limit  $\epsilon_{SB}$  (marked with arrows), which corresponds to an ideal gas. This can indicate some residual interactions in the system. According to the results from the RHIC<sup>2</sup>, the new phase of matter behaves more like an ideal fluid, than like a gas [9].

One of the key questions, to which current heavy ion physics tries to find an answer is the value of a critical temperature  $T_C$  as a function of a baryon chemical potential  $\mu_B$  (baryon density), where the phase transition occur. The results coming from the Lattice QCD are presented in the Fig. 1.6. The phase of matter in which quarks and gluons are deconfined is expected to exist at large

<sup>2</sup>Relativistic Heavy Ion Collider at Brookhaven National Laboratory in Upton, New York



Figure 1.6: Phase diagram coming from the Lattice QCD calculations [8].

temperatures. In the region of small temperatures and high baryon densities, a different state is supposed to appear - a *colour superconductor*. The phase transition between hadronic matter and QGP is thought to be of 1<sup>st</sup> order at  $\mu_B \gg 0$ . However as  $\mu_B \rightarrow 0$  quarks' masses become significant and a sharp transition transforms into a rapid but smooth cross-over. It is believed that in Pb-Pb collisions observed at the LHC<sup>3</sup>, the created matter has high enough temperature to be in the quark-gluon plasma phase, then cools down and converts into hadrons, undergoing a smooth transition [8].

### 1.3 Relativistic heavy ion collisions

#### 1.3.1 Stages of heavy ion collision

To create the quark-gluon plasma one has to achieve high enough temperatures and baryon densities. Such conditions can be recreated in the heavy ion collisions at the high energies. The left side of the Figure 1.7 shows simplified picture of a central collision of two highly relativistic nuclei in the centre-of-mass reference frame. The colliding nuclei are presented as thin disks because of the Lorentz contraction. In the central region, where the energy density is the highest, a new state of matter - the quark-gluon plasma - is supposedly created. Afterwards, the plasma expands and cools down, quarks combine into hadrons and their mutual interactions cease when the system reaches the *freeze-out* temperature. Subsequently, produced free hadrons move towards the detectors.

On the right side of the Figure 1.7 there is presented a space-time evolution of a collision process, plotted in the light-cone variables ( $z, t$ ). The two highly

<sup>3</sup>Large Hadron Collider at CERN, Geneva




Figure 1.7: Left: stages of a heavy ion collision simulated in the UrQMD model. Right: schematic view of a heavy ion collision evolution [8].

relativistic nuclei are traveling basically along the light cone until they collide at the centre of diagram. Nuclear fragments emerge from the collision again along the (forward) light cone, while the matter between fragmentation zones populates the central region. This hot and dense matter is believed to be in the state of the quark-gluon plasma. There exist several frameworks to describe this transition to the QGP phase, for example: QCD string breaking, QCD parton cascades or colour glass condensate evolving into glasma and later into quark-gluon plasma [10].

**String breaking** – In the string picture, the nuclei pass through each other forming colour strings. This is analogous to the situation depicted in the Fig 1.2 - the colour string is created between quarks inside particular nucleons in nuclei. In the next step strings decay / fragment forming quarks and gluons or directly hadrons. This approach becomes invalid at very high energies, when the strings overlap and cannot be treated as independent objects.

**Parton cascade** – The parton<sup>4</sup> cascade model is based on the pQCD. The colliding nuclei are treated as clouds of quarks and which penetrate through each other. The key element of this method is the time evolution of the parton phase-space distributions, which is governed by a relativistic Boltzmann equation with a collision term that contains dominant perturbative QCD interactions. The bottleneck of the parton cascade model is the low energies regime, where the  $Q^2$  is too small to be described by the perturbative theory.

<sup>4</sup>A parton is a common name for a quark and a gluon.

180 **Colour glass condensate** – The colour glass condensate assumes, that the had-  
 181 ron can be viewed as a tightly packed system of interacting gluons. The sat-  
 182 uration of gluons increases with energy, hence the total number of gluons may  
 183 increase without the bound. Such a saturated and weakly coupled gluon system  
 184 is called a colour glass condensate. The fast gluons in the condensate are Lorentz  
 185 contracted and redistributed on the two very thin sheets representing two col-  
 186 liding nuclei. The sheets are perpendicular to the beam axis. The fast gluons  
 187 produce mutually orthogonal colour magnetic and electric fields, that only ex-  
 188 ist on the sheets. Immediately after the collision, i.e. just after the passage of  
 189 the two gluonic sheets after each other, the longitudinal electric and magnetic  
 190 fields are produced forming the *glasma*. The glasma fields decay through the  
 191 classical rearrangement of the fields into radiation of gluons. Also decays due to  
 192 the quantum pair creations are possible. In this way, the quark-gluon plasma is  
 193 produced.

194 Interactions within the created quark-gluon plasma bring the system into the  
 195 local statistical equilibrium, hence its further evolution can be described by the  
 196 relativistic hydrodynamics. The hydrodynamic expansion causes that the sys-  
 197 tem becomes more and more dilute. The phase transition from the quark-gluon  
 198 plasma to the hadronic gas occurs. Further expansion causes a transition from the  
 199 strongly interaction hadronic gas to weakly interacting system of hadrons which  
 200 move freely to the detectors. Such decoupling of hadrons is called the *freeze-out*.  
 201 The freeze-out can be divided into two phases: the chemical freeze-out and the  
 202 thermal one. The *chemical freeze-out* occurs when the inelastic collisions between  
 203 constituents of the hadron gas stop. As the system evolves from the chemical  
 204 freeze-out to the thermal freeze-out the dominant processes are elastic collisions  
 205 (such as, for example  $\pi + \pi \rightarrow \rho \rightarrow \pi + \pi$ ) and strong decays of heavier reson-  
 206 ances which populate the yield of stable hadrons. The *thermal freeze-out* is the  
 207 stage of the evolution of matter, when the strongly coupled system transforms  
 208 to a weakly coupled one (consisting of essentially free particles). In other words  
 209 this is the moment, where the hadrons practically stop to interact. Obviously, the  
 210 temperatures corresponding to the two freeze-outs satisfy the condition

$$T_{chem} > T_{therm} , \quad (1.2)$$

211 where  $T_{chem}$  (inferred from the ratios of hadron multiplicities) is the temperature  
 212 of the chemical freeze-out, and  $T_{therm}$  (obtained from the investigation of the  
 213 transverse-momentum spectra) is the temperature of the thermal freeze-out [10].

### 214 1.3.2 QGP signatures

215 The quark-gluon plasma is a very short living and unstable state of matter.  
 216 One cannot investigate the properties of a plasma and confirm its existence di-  
 217 rectly. Hence, the several experimental effects were proposed as QGP signatures,  
 218 some of them have been already observed in heavy ion experiments [8]. As mat-  
 219 ter created in the heavy ions collisions is supposed to behave like a fluid, one

220 should expect appearance of collective behaviour at small transverse momenta  
 221 - so called *elliptic flow* and *radial flow*. The next signal is the temperature range  
 222 obtained from the measurements of *direct photons*, which gives us information,  
 223 that the system created in heavy ion collisions is far above the critical temperat-  
 224 ure obtained from the LQCD calculations. The *puzzle in the di-lepton spectrum* can  
 225 be explained by the modification of spectral shape of vector mesons (mostly  $\rho$   
 226 meson) in the presence of a dense medium. This presence of a medium can also  
 227 shed light on the *jet quenching* phenomenon - the suppression occurrence in the  
 228 high  $p_T$  domain.

### 229 Elliptic flow

230 In a non-central heavy ion collisions, created region of matter has an almond  
 231 shape with its shorter axis in the *reaction plane* (Fig. 1.8). The pressure gradient  
 232 is much larger in-plane rather than out-of-plane. This causes larger acceleration  
 and transverse velocities in-plane rather than out-of-plane. Such differences can




Figure 1.8: Overlapping region which is created in heavy ion collisions has an almond shape. Visible x-z plane is a *reaction plane*. The x-y plane is a *transverse plane*. The z is a direction of the beam [11].

233 be investigated by studying the distribution of particles with respect to the reac-  
 234 tion plane orientation [12]:  
 235

$$E \frac{d^3 N}{dp^3} = \frac{1}{2\pi} \frac{d^2 N}{p_T dp_T dy} (1 + 2v_1 \cos(\phi) + 2v_2 \cos(2\phi) + \dots), \quad (1.3)$$

236 where  $\phi$  is the angle between particle transverse momentum  $p_T$  (a momentum  
 237 projection on a transverse plane) and the reaction plane,  $N$  is a number of  
 238 particles and  $E$  is an energy of a particle. The  $y$  variable is a *rapidity* defined as:

$$y = \frac{1}{2} \ln \left( \frac{E + p_L}{E - p_L} \right), \quad (1.4)$$

where  $p_L$  is a longitudinal component of a momentum (parallel to the beam direction). The  $v_n$  coefficients indicate the shape of a system. For the most central collisions ( $b = 0$  - see Fig. 1.9) all coefficients vanish  $\bigwedge_{n \in N_+} v_n = 0$  (the overlapping region has the spherical shape). The Fourier series elements in the parentheses




Figure 1.9: Cross-section of a heavy ion collision in a transverse plane.  $\Psi_R$  is an angle between transverse plane and the reaction plane. The  $b$  parameter is an *impact parameter* - a distance between centers of nuclei during a collision. An impact parameter is related with the centrality of a collision and a volume of the quark-gluon plasma [12].

in Eq. 1.3 represent different kinds of a flow. The first value: “1” represents the *radial flow* - an isotropic flow in every direction. Next coefficient  $v_1$  is responsible for *direct flow*. The  $v_2$  coefficient is a measure of elliptic anisotropy (*elliptic flow*). The  $v_2$  has to build up in the early stage of a collision - later the system becomes too dilute: space asymmetry and the pressure gradient vanish. Therefore the observation of elliptic flow means that the created matter was in fact a strongly interacting matter.

The  $v_2$  coefficient was measured already at CERN SPS, LHC and RHIC. For the first time hydrodynamics successfully described the collision dynamics as the measured  $v_2$  reached hydrodynamic limit (Fig. 1.10). As expected, there is a mass ordering of  $v_2$  as a function of  $p_T$  (lower plot in the Fig. 1.10) with pions having the largest and protons the smallest anisotropy. In the upper plots in the Fig. 1.10 there is a  $v_2$  as a function of transverse kinetic energy. The left plot shows the two universal trend lines for baryons and mesons. After the scaling of  $v_2$  and the kinetic energy by the number of valence quarks, all of the hadrons follow the same universal curve. Those plots show that strong collectivity is observed in heavy ion collisions.





Figure 1.10: *Lower:* The elliptic flow  $v_2$  follows the hydrodynamical predictions for an ideal fluid perfectly. Note that  $> 99\%$  of all final hadrons have  $p_T < 1.5$  GeV/c. *Upper left:* The  $v_2$  plotted versus transverse kinetic energy  $KE_T = m_T - m_0 = \sqrt{p_T^2 + m_0^2} - m_0$ . The  $v_2$  follows different universal curves for mesons and baryons. *Upper right:* When scaled by the number of valence quarks, the  $v_2$  follows the same universal curve for all hadrons and for all values of scaled transverse kinetic energy [13].

## 260 Transverse radial flow

261 Elliptic flow described previously is caused by the pressure gradients which  
 262 must also produce a more simple collective behaviour of matter - a movement  
 263 inside-out, called radial flow. Particles are pushed to higher momenta and they  
 264 move away from the center of the collision. A source not showing collective

behaviour, like pp collisions, produces particle spectra that can be fitted by a power-law [8]:

$$\frac{1}{2\pi p_T} \frac{d^2 N}{dp_T d\eta} = C \left( 1 + \frac{p_T}{p_0} \right)^{-n} . \quad (1.5)$$

The  $\eta$  variable is a *pseudorapidity* defined as follows:

$$\eta = \frac{1}{2} \ln \left( \frac{p + p_L}{p - p_L} \right) = -\ln \left( \frac{\theta}{2} \right) , \quad (1.6)$$

where  $\theta$  is an emission angle  $\cos \theta = p_L/p$ .




Figure 1.11: Invariant yield of particles versus transverse mass  $m_T = \sqrt{p_T^2 + m_0^2}$  for  $\pi^\pm$ ,  $K^\pm$ ,  $p$  and  $\bar{p}$  at mid-rapidity for p+p collisions (bottom) and Au+Au events from 70-80% (second bottom) to 0-5% (top) centrality [14].

The hydrodynamical expansion of a system gives the same flow velocity kick for different kind of particles - ones with bigger masses will gain larger  $p_T$  boost. This causes increase of the yield of particles with larger transverse momenta. In the invariant yield plots one can observe the decrease of the slope parameter, especially for the heavier hadrons. This is presented in the Fig. 1.11. The most affected spectra are ones of kaons (b) and protons (c). One can notice decrease of the slope parameter for heavy ion collisions (plots from second bottom to top) comparing to the proton-proton collisions (bottom ones), where no boost from radial flow should occur [8].

## Direct photons

The direct photons are photons, which are not coming from the final state hadrons decays. Their sources can be various interaction from charged particles

created in the collision, either at the partonic or at the hadronic level. Direct photons are considered to be an excellent probe of the early stage of the collision. This is because their mean free path is very large to the created system in the collision. Thus photons created at the early stage leave the system without suffering any interaction and retain information about this stage, in particular about its temperature.

One can distinguish two kinds of direct photons: *thermal* and *prompt*. Thermal photons can be emitted from the strong processes in the quark-gluon plasma involving quarks and gluons or hot hadronic matter (e.g. processes:  $\pi\pi \rightarrow \rho\gamma$ ,  $\pi\rho \rightarrow \pi\gamma$ ). Thermal photons can be observed in the low  $p_T$  region. Prompt photons are believed to come from “hard” collisions of initial state partons belonging to the colliding nuclei. The prompt photons can be described using the pQCD. They will dominate the high  $p_T$  region. The analysis of transverse momentum of spectra of direct photons revealed, that the temperature of the source of thermal photons produced in heavy ion collisions at RHIC is in the range 300–600 MeV (Fig. 1.12). Hence the direct photons had to come from a system whose temperature is far above from the critical temperature for QGP creation.




Figure 1.12: Thermal photons spectra for the central Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV at computed within different hydrodynamical models compared with the pQCD calculations (solid line) and experimental data from PHENIX (black dots) [15].

### Puzzle in di-lepton mass spectrum

The invariant mass spectra (Fig. 1.13) of lepton pairs reveal many peaks corresponding to direct decays of various mesons into a lepton pair. The continuous background in this plot is caused by the decays of hadrons into more than two leptons (including so-called *Dalitz decays* into a lepton pair and a photon). Particular hadron decay channels, which contribute to this spectrum are shown




Figure 1.13: Left: Invariant mass spectrum of  $e^+e^-$  pairs in Pb+Au collisions at 158A GeV compared to the sum coming from the hadron decays predictions. Right: The expectations coming from model calculations assuming a dropping of the  $\rho$  mass (blue) or a spread of the  $\rho$  width in the medium (red) [16].

in the Fig. 1.13 with the coloured lines and their sum with the black one. The sum (called *the hadronic cocktail*) of various components describes experimental spectra coming from the simple collisions (like p+p or p+A) quite well with the statistical and systematical uncertainties [9]. This situation is different considering more complicated systems i.e. A+A. Spectra coming from Pb+Au collisions are presented on the plots in the Fig. 1.13. The “hadronic cocktail” does not describe the data, in the mass range between the  $\pi$  and the  $\rho$  mesons a significant excess of electron pairs over the calculated sum is observed. Theoretical explanation of this phenomenon assumes modification of the spectral shape of vector mesons in a dense medium. Two different interpretations of this increase were proposed: a decrease of meson mass with the medium density and increase of the meson width in the dense medium. In principle, one could think of simultaneous occurrence of both effects: mass shift and resonance broadening. Experimental results coming from the CERES disfavour the mass shift hypothesis indicating only broadening of resonance peaks (Fig. 1.13b) [9].

### 319 Jet quenching

320 A jet is defined as a group of particles with close vector momenta and high en-  
 321 ergies. It has its beginning when the two partons are going in opposite directions  
 322 and have energy big enough to produce new quark-antiquark pair and then ra-  
 323 diate gluons. This process can be repeated many times and it results in two back-  
 324 to-back jets of hadrons. It has been found that jets in the opposite hemisphere  
 325 (*away-side jets*) show a very different pattern in d+Au and Au+Au collisions. This  
 326 is shown in the azimuthal correlations in the Fig. 1.14. In d+Au collisions, like in  
 327 p+p, a pronounced away-side jet appears around  $\Delta\phi = \pi$ , exactly opposite to the  
 328 trigger jet, what is typical for di-jet events. In central Au+Au collisions the away-  
 jet is suppressed. When the jet has its beginning near the surface of the quark-




Figure 1.14: Azimuthal angle difference  $\Delta\phi$  distributions for different colliding systems at  $\sqrt{s_{NN}} = 200$  GeV. Transverse momentum cut:  $p_T > 2$  GeV. For the Au+Au collisions the away-side jet is missing [17].

329 quon plasma, one of the jets (*near-side jet*) leaves the system almost without any  
 330 interactions. This jet is visible on the correlation plot as a high peak at  $\Delta\phi = 0$ .  
 331 However, the jet moving towards the opposite direction has to penetrate a dense  
 332 medium. The interaction with the plasma causes energy dissipation of particles  
 333 and is visible on an azimuthal correlation plot as disappearance of the away-side  
 334 jet [9].  
 335

## Chapter 2

# Therminator model

`THERMINATOR` [18] is a Monte Carlo event generator designed to investigate the particle production in the relativistic heavy ion collisions. The functionality of the code includes a generation of the stable particles and unstable resonances at the chosen hypersurface model. It performs the statistical hadronization which is followed by space-time evolution of particles and the decay of resonances. The key element of this method is an inclusion of a complete list of hadronic resonances, which contribute very significantly to the observables. The second version of `THERMINATOR` [19] comes with a possibility to incorporate any shape of freeze-out hypersurface and the expansion velocity field, especially those generated externally with various hydrodynamic codes.

### 2.1 (3+1)-dimensional viscous hydrodynamics

Most of the relativistic viscous hydrodynamic calculations are done in (2+1)-dimensions. Such simplification assumes boost-invariance of a matter created in a collision. Experimental data reveals that no boost-invariant region is formed in the collisions [20]. Hence, for the better description of created system a (3+1)-dimensional model is required.

In the four dimensional relativistic dynamics one can describe a system using a space-time four-vector  $x^\nu = (ct, x, y, z)$ , a velocity four-vector  $u^\nu = \gamma(c, v_x, v_y, v_z)$  and a energy-momentum tensor  $T^{\mu\nu}$ . The particular components of  $T^{\mu\nu}$  have a following meaning:

- $T^{00}$  - an energy density,
- $cT^{0\alpha}$  - an energy flux across a surface  $x^\alpha$ ,
- $T^{\alpha 0}$  - an  $\alpha$ -momentum flux across a surface  $x^\alpha$  multiplied by  $c$ ,
- $T^{\alpha\beta}$  - components of momentum flux density tensor,

where  $\gamma = (1 - v^2/c^2)^{-1/2}$  is Lorentz factor and  $\alpha, \beta \in \{1, 2, 3\}$ . Using  $u^\nu$  one can express  $T^{\mu\nu}$  as follows [21]:

$$T_0^{\mu\nu} = (e + p)u^\mu u^\nu - pg^{\mu\nu} \quad (2.1)$$

where  $e$  is an energy density,  $p$  is a pressure and  $g^{\mu\nu}$  is an inverse metric tensor:

$$g^{\mu\nu} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}. \quad (2.2)$$

The presented version of energy-momentum tensor (Eq. 2.1) can be used to describe dynamics of a perfect fluid. To take into account influence of viscosity, one has to apply the following corrections coming from shear  $\pi^{\mu\nu}$  and bulk  $\Pi$  viscosities [22]:

$$T^{\mu\nu} = T_0^{\mu\nu} + \pi^{\mu\nu} + \Pi(g^{\mu\nu} - u^\mu u^\nu). \quad (2.3)$$

The stress tensor  $\pi^{\mu\nu}$  and the bulk viscosity  $\Pi$  are solutions of dynamical equations in the second order viscous hydrodynamic framework [21]. The comparison of hydrodynamics calculations with the experimental results reveal, that the shear viscosity divided by entropy  $\eta/s$  has to be small and close to the AdS/CFT estimate  $\eta/s = 0.08$  [22, 23].

When using  $T^{\mu\nu}$  to describe system evolving close to local thermodynamic equilibrium, relativistic hydrodynamic equations in a form of:

$$\partial_\mu T^{\mu\nu} = 0 \quad (2.4)$$

can be used to describe the dynamics of the local energy density, pressure and flow velocity.

Hydrodynamic calculations are starting from the Glauber<sup>1</sup> model initial conditions. The collective expansion of a fluid ends at the freeze-out hypersurface. That surface is usually defined as a constant temperature surface, or equivalently as a cut-off in local energy density. The freeze-out is assumed to occur at the temperature  $T = 140$  MeV.

## 2.2 Statistical hadronization

Statistical description of heavy ion collision has been successfully used to describe quantitatively *soft* physics, i.e. the regime with the transverse momentum not exceeding 2 GeV. The basic assumption of the statistical approach of evolution of the quark-gluon plasma is that at some point of the space-time evolution of the fireball, the thermal equilibrium is reached. When

<sup>1</sup>The Glauber Model is used to calculate “geometrical” parameters of a collision like an impact parameter, number of participating nucleons or number of binary collisions.

the system is in the thermal equilibrium the local phase-space densities of particles follow the Fermi-Dirac or Bose-Einstein statistical distributions. At the end of the plasma expansion, the freeze-out occurs. The freeze-out model incorporated in the THERMINATOR model assumes, that chemical and thermal freeze-out occur at the same time.

### 2.2.1 Cooper-Frye formalism

The result of the hydrodynamic calculations is the freeze-out hypersurface  $\Sigma^\mu$ . A three-dimensional element of the surface is defined as [19]

$$d\Sigma_\mu = \epsilon_{\mu\alpha\beta\gamma} \frac{\partial x^\alpha}{\partial \alpha} \frac{\partial x^\beta}{\partial \beta} \frac{\partial x^\gamma}{\partial \gamma} d\alpha d\beta d\gamma, \quad (2.5)$$

where  $\epsilon_{\mu\alpha\beta\gamma}$  is the Levi-Civita tensor and the variables  $\alpha, \beta, \gamma \in \{1, 2, 3\}$  are used to parametrize the three-dimensional freeze-out hypersurface in the Minkowski four-dimensional space. The Levi-Civita tensor is equal to 1 when the indices form an even permutation (eg.  $\epsilon_{0123}$ ), to -1 when the permutation is odd (e.g.  $\epsilon_{2134}$ ) and has a value of 0 if any index is repeated. Therefore [19],

$$d\Sigma_0 = \begin{vmatrix} \frac{\partial x}{\partial \alpha} & \frac{\partial x}{\partial \beta} & \frac{\partial x}{\partial \gamma} \\ \frac{\partial y}{\partial \alpha} & \frac{\partial y}{\partial \beta} & \frac{\partial y}{\partial \gamma} \\ \frac{\partial z}{\partial \alpha} & \frac{\partial z}{\partial \beta} & \frac{\partial z}{\partial \gamma} \end{vmatrix} d\alpha d\beta d\gamma \quad (2.6)$$

and the remaining components are obtained by cyclic permutations of  $t, x, y$  and  $z$ .

One can obtain the number of hadrons produced on the hypersurface  $\Sigma^\mu$  from the Cooper-Frye formalism. The following integral yields the total number of created particles [19]:

$$N = (2s + 1) \int \frac{d^3p}{(2\pi)^3 E_p} \int d\Sigma_\mu(x) p^\mu f(x, p), \quad (2.7)$$

where

$$f(p \cdot u) = \left\{ \exp \left[ \frac{p_\mu u^\mu - (B\mu_B + I_3\mu_{I_3} + S\mu_S + C\mu_C)}{T} \right] \pm 1 \right\}^{-1} \quad (2.8)$$

is the phase-space distribution for particles (for stable ones and resonances). For the Fermi-Dirac distribution in the Eq. 2.8 there is a plus sign and for Bose-Einstein statistics minus sign respectively. The thermodynamic quantities appearing in the  $f(\cdot)$  are  $T$  - temperature,  $\mu_B$  - baryon chemical potential,  $\mu_{I_3}$  - isospin chemical potential,  $\mu_S$  - strange chemical potential,  $\mu_C$  - charmed chemical potential and the  $s$  is a spin of a particle. One can simply derive from Eq. 2.7, the dependence of the momentum density [24]:

$$E \frac{dN}{d^3p} = \int f(x, p) p^\mu d\Sigma_\mu. \quad (2.9)$$



416 The equations presented above are directly used in the THERMINATOR to generate  
417 the hadrons with the Monte-Carlo method.

## 418 Chapter 3

# 419 Particle interferometry

420 Two-particle interferometry (also called *femtoscopy*) gives a possibility to  
421 investigate space-time characteristics of the particle-emitting source created  
422 in heavy ion collisions. Through the study of particle correlations, their  
423 momentum distributions can be used to obtain information about the spatial  
424 extent of the created system. Using this method, one can measure sizes of the  
425 order of  $10^{-15}$  m and time of the order of  $10^{-23}$  s.

### 426 3.1 HBT interferometry

427 In the 1956 Robert Hanbury Brown and Richard Q. Twiss proposed a  
428 method which through analysis of interference between photons allowed to  
429 investigate angular dimensions of stars. The most important result from the  
430 Hanbury-Brown-Twiss experiments is that two indistinguishable particles can  
431 produce an interference effect. There is almost no difference between normal  
432 interferometry and HBT method, except that the latter one does not take into  
433 account information about phase shift of registered particles. At the beginning  
434 this method was used in astronomy for photon interference, but this effect can  
435 be used also to measure extent of any emitting source. This method was adapted  
436 to heavy ion collisions to investigate dimensions of a system created in those  
437 collisions by studying correlations of identical particles [25]. The main difference  
438 between HBT method in astronomy and femtoscopy is that the first one is based  
439 on space-time HBT correlations and the latter one uses momentum correlations.  
440 The momentum correlations yield the space-time picture of the source, whereas  
441 the space-time HBT correlations provide the characteristic relative momenta of  
442 emitted photons, which gives the angular size of the star without the knowledge  
443 of its radius and lifetime [10].

## 3.2 Theoretical approach

Intensity interferometry in heavy ion physics uses similar mathematical formalism as the astronomy HBT measurement. Through the measurement of correlation between particles as a function of their relative momentum one can deduce the average separation between emitting sources.

### 3.2.1 Conventions used

In heavy ion collisions to describe particular directions, components of momentum and location of particles, one uses naming convention called the Bertsch-Pratt coordinate system. This system is presented in the Fig. 3.1. The three dir-




Figure 3.1: Bertsch-Pratt direction naming convention used in heavy ion collision.

Directions are called *longitudinal*, *outward* and *sideward*. The longitudinal direction is parallel to the beam axis. The plane perpendicular to the beam axis is called a *transverse plane*. A projection of a particle pair momentum  $\mathbf{k} = (\mathbf{p}_a + \mathbf{p}_b)/2$  on a transverse plane (a *transverse momentum*  $\mathbf{k}_T$ ) determines *outward* direction:  $(\mathbf{k})_{out} = \mathbf{k}_T$ . A direction perpendicular to the longitudinal and outward is called *sideward*.

A particle pair is usually described using two coordinate systems. The first one, *Longitudinally Co-Moving System (LCMS)* is moving along the particle pair with the longitudinal direction, in other words, the pair longitudinal momentum vanishes:  $(\mathbf{p}_a)_{long} = -(\mathbf{p}_b)_{long}$ . The second system is called *Pair Rest Frame (PRF)*. In the PRF the centre of mass rests:  $\mathbf{p}_a = -\mathbf{p}_b$ . Variables which are expressed in the PRF are marked with a star (e.g.  $\mathbf{k}^*$ ).

The transition of space-time coordinates from LCMS to PRF is simply a boost along the outward direction, with the transverse velocity of the

pair  $\beta_t = (\mathbf{v}/c)_{out}$  [25]:

$$r_{out}^* = \gamma_t(r_{out} - \beta_t \Delta t) \quad (3.1)$$

$$r_{side}^* = r_{side} \quad (3.2)$$

$$r_{long}^* = r_{long} \quad (3.3)$$

$$\Delta t^* = \gamma_t(\Delta t - \beta_t r_{out}) , \quad (3.4)$$

where  $\gamma_t = (1 - \beta_t^2)^{-1/2}$  is the Lorentz factor. However, in calculations performed in this work the equal time approximation is used, which assumes that particles in a pair were produced at the same time in PRF - the  $\Delta t^*$  is neglected.

The most important variables used to describe particle pair are their total momentum  $\mathbf{P} = \mathbf{p}_a + \mathbf{p}_b$  and relative momentum  $\mathbf{q} = \mathbf{p}_a - \mathbf{p}_b$ . In the PRF one has  $\mathbf{q} = 2\mathbf{k}^*$ , where  $\mathbf{k}^*$  is a momentum of the first particle in PRF.

### 3.2.2 Two particle wave function

Let us consider two identical particles with momenta  $\mathbf{p}_1$  and  $\mathbf{p}_2$  emitted from space points  $\mathbf{x}_1$  and  $\mathbf{x}_2$ . Those emitted particles can be treated as two incoherent waves. If the particles are identical, they are also indistinguishable, therefore one




Figure 3.2: The pair wave function is a superposition of all possible states. In case of particle interferometry it includes two cases: particles with momenta  $p_1, p_2$  registered by detectors  $A, B$  and  $p_1, p_2$  registered by  $B, A$  respectively.

has also take into account the scenario, where the particle with momentum  $\mathbf{p}_1$  is emitted from  $\mathbf{x}_2$  and particle  $\mathbf{p}_2$  from  $\mathbf{x}_1$  (Fig. 3.2). In such case, the wave function describing behaviour of a pair has to contain both components [8]:

$$\Psi_{ab}(\mathbf{q}) = \frac{1}{\sqrt{2}} [\exp(-i\mathbf{p}_1\mathbf{x}_1 - i\mathbf{p}_2\mathbf{x}_2) \pm \exp(-i\mathbf{p}_2\mathbf{x}_1 - i\mathbf{p}_1\mathbf{x}_2)] . \quad (3.5)$$

A two particle wave function of identical bosons is symmetric ("+" sign in Eq. 3.5) and in case of identical fermions - antisymmetric ("-") sign). This anti-symmetrization or symmetrization implies the correlation effect coming from the Fermi-Dirac or Bose-Einstein statistics accordingly.

To provide full description of a system consisting of two charged hadrons, one has to include in the wave function besides quantum statistics also Coulomb and strong Final State Interactions. Considering identical particles systems, the quantum statistics is a main source of a correlation. Hence, in case of space-time analysis of particle emitting source, effects coming from the Coulomb and Strong interactions can be neglected.

### 3.2.3 Source emission function

To describe particle emitting source, one uses a single emission function [25]:

$$S_A(\mathbf{x}_1, \mathbf{p}_1) = \int S(\mathbf{x}_1, \mathbf{p}_1, \mathbf{x}_2, \mathbf{p}_2, \dots, \mathbf{x}_N, \mathbf{p}_N) d\mathbf{x}_2 d\mathbf{p}_2 \dots d\mathbf{x}_N d\mathbf{p}_N \quad (3.6)$$

and a two-particle one:

$$S_{AB}(\mathbf{x}_1, \mathbf{p}_1, \mathbf{x}_2, \mathbf{p}_2) = \int S(\mathbf{x}_1, \mathbf{p}_1, \mathbf{x}_2, \mathbf{p}_2, \dots, \mathbf{x}_N, \mathbf{p}_N) d\mathbf{x}_3 d\mathbf{p}_3 \dots d\mathbf{x}_N d\mathbf{p}_N. \quad (3.7)$$

Emission function  $S(\cdot)$  can be interpreted as a probability to emit a particle, or a pair of particles from a given space-time point with a given momentum. In principle, the source emission function should encode all physics aspects of the particle emission process i.e. the symmetrization for bosons and fermions, as well as the two-body and many body Final State Interactions. Instead of this, one assume that each particle's emission process is independent - the interaction between final-state particles after their creation is independent from their emission process. The assumption of this independence allows to construct two-particle emission function from single particle emission functions via a convolution [25]:

$$S(\mathbf{k}^*, \mathbf{r}^*) = \int S_A(\mathbf{p}_1, \mathbf{x}_1) S_B(\mathbf{p}_2, \mathbf{x}_2) \delta \left[ \mathbf{k}^* - \frac{\mathbf{p}_1 + \mathbf{p}_2}{2} \right] \delta [\mathbf{r}^* - (\mathbf{x}_1 + \mathbf{x}_2)] \times d^4 \mathbf{x}_1 d^4 \mathbf{x}_2 d^4 \mathbf{x}_1 d^3 \mathbf{p}_1 d^3 \mathbf{p}_2 \quad (3.8)$$

In case of identical particles, ( $S_A = S_B$ ) several simplifications can be made. A convolution of the two same Gaussian distributions is also a Gaussian distribution with  $\sigma$  multiplied by  $\sqrt{2}$ . Femtoscopy can give information only about two-particle emission function, but when considering Gaussian distribution as a source function in Eq. 3.8, one can obtain a  $\sigma$  of a single emission function from a two-particle emission function. The Eq. 3.8 is not reversible - an information about  $S_A(\cdot)$  cannot be derived from  $S_{AB}(\cdot)$ . An exception from this rule is a Gaussian source function, hence it is often used in femtoscopic calculations. Considering pairs of identical particles, an emission function is assumed to be described by the following equation in the Pair Rest Frame [25]:

$$S_{1D}^{PRF}(\mathbf{r}^*) = \exp \left( - \frac{r_{out}^{*2} + r_{side}^{*2} + r_{long}^{*2}}{4R_{inv}^2} \right). \quad (3.9)$$

To change from the three-dimensional variables to the one-dimensional variable one requires introduction of the proper Jacobian  $r^{*2}$ :

$$\boxed{S_{1D}^{PRF}(r^*) = r^{*2} \exp \left( - \frac{r^{*2}}{4R_{inv}^2} \right)}. \quad (3.10)$$

512 The “4” in the denominator before the “standard deviation”  $R_{inv}$  in the Gaus-  
 513 sian distribution comes from the convolution of the two Gaussian distributions,  
 514 which multiplies the  $R_{inv}$  by a factor of  $\sqrt{2}$ .

A more complex form of emission function was used by all RHIC and SPS experiments in identical pion femtoscopy:

$$S_{3D}^{LCMS}(\mathbf{r}) = \exp \left( -\frac{r_{out}^2}{4R_{out}^2} - \frac{r_{side}^2}{4R_{side}^2} - \frac{r_{long}^2}{4R_{long}^2} \right). \quad (3.11)$$

515 The main difference of this source function is that it has three different and inde-  
 516 pendent widths  $R_{out}$ ,  $R_{side}$ ,  $R_{long}$  and they are defined in the LCMS, not in PRF.  
 517 Unlike in PRF, in LCMS an equal-time approximation is not used. For identical  
 518 particles this is not a problem - only Coulomb interaction inside a wave function  
 519 depends on  $\Delta t$ .

#### 520 Relationship between one-dimensional and three-dimensional source sizes

521 Up to now, most of femtoscopic measurements were limited only to averaged  
 522 source size  $R_{av}^L$  (the letter “L” in superscript stands for LCMS):

$$S_{1D}^{LCMS}(r) = \exp \left( -\frac{r_{out}^2 + r_{side}^2 + r_{long}^2}{2R_{av}^L{}^2} \right). \quad (3.12)$$

523 The relationship between  $S_{1D}^{LCMS}(\cdot)$  and  $S_{3D}^{LCMS}(\cdot)$  is given by:

$$S_{3D}^{LCMS}(r) = \int \exp \left( -\frac{r_{out}^2}{2R_{out}^L{}^2} - \frac{r_{side}^2}{2R_{side}^L{}^2} - \frac{r_{long}^2}{2R_{long}^L{}^2} \right) \times \delta \left( r - \sqrt{r_{out}^2 + r_{side}^2 + r_{long}^2} \right) dr_{out} dr_{side} dr_{long}. \quad (3.13)$$

524 The one-dimensional source size corresponding to the three-dimensional one can  
 525 be approximated by the following form:

$$S_{1D}^{LCMS}(r) = r^2 \exp \left( -\frac{r^2}{2R_{av}^L{}^2} \right). \quad (3.14)$$

526 The equation above assumes that  $R_{out}^L = R_{side}^L = R_{long}^L$  hence  $R_{av}^L = R_{out}^L$ . If this  
 527 condition is not satisfied, one can not give explicit mathematical relation between  
 528 one-dimensional and three-dimensional source sizes. However, for realistic val-  
 529 ues of  $R$  (i.e. for similar values of  $R_{out}$ ,  $R_{side}$ ,  $R_{long}$ ), the  $S_{3D}^{LCMS}$  from Eq. 3.13 is  
 530 not very different from Gaussian distribution and can be well approximated by  
 531 Eq. 3.13.

532 A deformation of an averaged source function in case of big differences in  
 533  $R_{out}$ ,  $R_{side}$ ,  $R_{long}$  is presented in the Fig. 3.3. A three-dimensional Gaussian dis-  
 534 tribution with varying widths was averaged into one-dimensional function using




Figure 3.3: An averaged three-dimensional Gaussian source function with different widths was averaged into one-dimensional function. To illustrate deformations, one-dimensional Gaussian distribution was fitted.

the Eq. 3.13. Afterwards, an one-dimensional Gaussian distribution was fitted. One can notice a heavy tail of an averaged distribution in long  $r$  region, which makes an approximation using one-dimensional distribution in this case quite inaccurate.

Using Eq. 3.13 and Eq. 3.14 one can obtain a relation between one-dimensional width and the three-dimensional ones. Through numerical calculations one can find the following approximate relation [25]:

$$R_{av}^L = \sqrt{\left(R_{out}^L{}^2 + R_{side}^L{}^2 + R_{long}^L{}^2\right)/3} . \quad (3.15)$$

This equation does not depend on the pair velocity, hence it is valid in the LCMS and PRF.

### 3.2.4 Theoretical correlation function

The fundamental object in a particle interferometry is a correlation function. The correlation function is defined as:

$$C(\mathbf{p}_a, \mathbf{p}_b) = \frac{P_2(\mathbf{p}_a, \mathbf{p}_b)}{P_1(\mathbf{p}_a)P_1(\mathbf{p}_b)} , \quad (3.16)$$

where  $P_2$  is a conditional probability to observe a particle with momentum  $\mathbf{p}_b$  if particle with momentum  $\mathbf{p}_a$  was also observed. A  $P_1$  is a probability to observe a particle with a given momentum. The relationship between source emission function, pair wave function and the correlation function is described by the following equation:

$$C(\mathbf{p}_1, \mathbf{p}_2) = \int S_{AB}(\mathbf{p}_1, \mathbf{x}_1, \mathbf{p}_2, \mathbf{x}_2) |\Psi_{AB}|^2 d^4\mathbf{x}_1 d^4\mathbf{x}_2 \quad (3.17)$$

Substituting the one-dimensional emission function (Eq. 3.10) into the integral above yields the following form of correlation function in PRF:

$$C(q) = 1 + \lambda \exp(-R_{inv}^2 q^2) \quad (3.18)$$

where  $q$  is a momentum difference between two particles. When using the three-dimensional emission function (Eq. 3.11) one gets the following correlation function defined in the LCMS:

$$C(\mathbf{q}) = 1 + \lambda \exp(-R_{out}^2 q_O^2 - R_{side}^2 q_S^2 - R_{long}^2 q_L^2) \quad (3.19)$$

where  $q_O, q_S, q_L$  are  $\mathbf{q}$  components in the outward, sideward and longitudinal direction. The  $\lambda$  parameter in the equations above determines correlation strength. The lambda parameter has values in the range  $\lambda \in [-0.5, 1]$  and it depends on a pair type. In case of pairs of identical bosons (like  $\pi$ - $\pi$  or  $K$ - $K$ ) the lambda parameter  $\lambda \rightarrow 1$ . For identical fermions (e.g.  $p$ - $p$ )  $\lambda \rightarrow -0.5$ . Values of  $\lambda$  observed experimentally are lower than 1 (for bosons) and greater than -0.5 (for fermions). There are few explanations to this effect: detector efficiencies, inclusion of misidentified particles in a used sample or inclusion of non-correlated pairs (when one or both particles come from e.g. long-lived resonance). The analysis carried out in this work uses data from a model, therefore the detector efficiency and particle purity is not taken into account.

### 3.2.5 Spherical harmonics decomposition of a correlation function

Results coming from a analysis using three-dimensional correlation function in Cartesian coordinates are quite difficult to visualize. To do that one usually performs a projection into a one dimension in outward, sideward and longitudinal directions. Doing that, one loose important information about a correlation function

## 3.3 Experimental approach

## 3.4 Scaling of femtosopic radii



## 568 **Chapter 4**

# 569 **Results**

### 570 **4.1 Identical particles correlations**

### 571 **4.2 Results of the fit**

### 572 **4.3 Discussion of results**

573 **Chapter 5**

574 **Summary**

## Bibliography

- [1] Standard Model of Elementary Particles - Wikipedia, the free encyclopedia  
[http://en.wikipedia.org/wiki/standard\\_model](http://en.wikipedia.org/wiki/standard_model).
- [2] R. Aaij et al. (LHCb Collaboration). Observation of the resonant character of the  $\psi(4430)^-$  state. *Phys. Rev. Lett.*, 112:222002, Jun 2014.
- [3] Donald H. Perkins. *Introduction to High Energy Physics*. Cambridge University Press, fourth edition, 2000. Cambridge Books Online.
- [4] G. Odyniec. *Phase Diagram of Quantum Chromo-Dynamics* - course at Faculty of Physics, Warsaw University of Technology, Jun 2012.
- [5] J. Beringer et al. (Particle Data Group). The Review of Particle Physics. *Phys. Rev.*, D86:010001, 2012.
- [6] Z. Fodor and S.D. Katz. The Phase diagram of quantum chromodynamics. 2009.
- [7] F. Karsch. Lattice results on QCD thermodynamics. *Nuclear Physics A*, 698(1-4):199 – 208, 2002.
- [8] Adam Kisiel. *Studies of non-identical meson-meson correlations at low relative velocities in relativistic heavy-ion collisions registered in the STAR experiment*. PhD thesis, Warsaw University of Technology, Aug 2004.
- [9] J. Bartke. *Relativistic Heavy Ion Physics*. World Scientific Pub., 2009.
- [10] W. Florkowski. *Phenomenology of Ultra-Relativistic Heavy-Ion Collisions*. World Scientific, 2010.
- [11] Science Grid This Week, October 25, 2006 - Probing the Perfect Liquid with the STAR Grid  
[http://www.interactions.org/sgtw/2006/1025/star\\_grid\\_more.html](http://www.interactions.org/sgtw/2006/1025/star_grid_more.html).
- [12] K. Grebieszko. Fizyka zderzeń ciężkich jonów,  
<http://www.if.pw.edu.pl/~kperl/hip/hip.html>.
- [13] Ulrich W. Heinz. From SPS to RHIC: Maurice and the CERN heavy-ion programme. *Phys.Scripta*, 78:028005, 2008.

- [14] J. Adams et al. Identified particle distributions in pp and Au+Au collisions at  $\sqrt{s(NN)} = 200$  GeV. *Phys.Rev.Lett.*, 92:112301, 2004.
- [15] G. David, R. Rapp, and Z. Xu. Electromagnetic Probes at RHIC-II. *Phys.Rept.*, 462:176–217, 2008.
- [16] A. Marin et al. Dilepton measurements with CERES. *PoS*, CPOD07:034, 2007.
- [17] John Adams et al. Experimental and theoretical challenges in the search for the quark gluon plasma: The STAR Collaboration’s critical assessment of the evidence from RHIC collisions. *Nucl.Phys.*, A757:102–183, 2005.
- [18] Adam Kisiel, Tomasz Taluc, Wojciech Broniowski, and Wojciech Florkowski. THERMINATOR: THERMal heavy-IoN generATOR. *Comput.Phys.Commun.*, 174:669–687, 2006.
- [19] Mikolaj Chojnacki, Adam Kisiel, Wojciech Florkowski, and Wojciech Broniowski. THERMINATOR 2: THERMal heavy IoN generATOR 2. *Comput.Phys.Commun.*, 183:746–773, 2012.
- [20] I. et al (BRAHMS Collaboration) Bearden. Charged meson rapidity distributions in central Au + Au collisions at  $\sqrt{s_{NN}} = 200$  GeV. *Phys. Rev. Lett.*, 94:162301, Apr 2005.
- [21] W. Israel and J.M. Stewart. Transient relativistic thermodynamics and kinetic theory. *Annals of Physics*, 118(2):341 – 372, 1979.
- [22] Piotr Bożek. Flow and interferometry in (3 + 1)-dimensional viscous hydrodynamics. *Phys. Rev. C*, 85:034901, Mar 2012.
- [23] K. Kovtun, P. D. T. Son, and A. O. Starinets. Viscosity in strongly interacting quantum field theories from black hole physics. *Phys. Rev. Lett.*, 94:111601, Mar 2005.
- [24] Fred Cooper and Graham Frye. Single-particle distribution in the hydrodynamic and statistical thermodynamic models of multiparticle production. *Phys. Rev. D*, 10:186–189, Jul 1974.
- [25] Adam Kisiel. Nonidentical-particle femtoscopy at  $\sqrt{s_{NN}} = 200$  GeV in hydrodynamics with statistical hadronization. *Phys. Rev. C*, 81:064906, Jun 2010.

## List of Figures

635	1.1	The Standard Model of elementary particles [1]. . . . .	2
636	1.2	A string break and a creation of a pair quark-anti-quark [4]. . . . .	4
637	1.3	The coupling parameter $\alpha_s$ dependence on four-momentum transfer $Q^2$ [5]. . . . .	5
638			
639	1.4	The QCD potential for a pair quark-antiquark as a function of distance for different temperatures. A value of a potential decreases with the temperature [4]. . . . .	5
640			
641			
642	1.5	A number of degrees of freedom as a function of a temperature [7].	6
643	1.6	Phase diagram coming from the Lattice QCD calculations [8]. . . .	7
644	1.7	Left: stages of a heavy ion collision simulated in the UrQMD model. Right: schematic view of a heavy ion collision evolution [8].	8
645			
646	1.8	Overlapping region which is created in heavy ion collisions has an almond shape. Visible x-z plane is a <i>reaction plane</i> . The x-y plane is a <i>transverse plane</i> . The z is a direction of the beam [11]. . . . .	10
647			
648			
649	1.9	Cross-section of a heavy ion collision in a transverse plane. $\Psi_R$ is an angle between transverse plane and the reaction plane. The $b$ parameter is an <i>impact parameter</i> - a distance between centers of nuclei during a collision. An impact parameter is related with the centrality of a collision and a volume of the quark-gluon plasma [12].	11
650			
651			
652			
653			
654	1.10	<i>Lower:</i> The elliptic flow $v_2$ follows the hydrodynamical predictions for an ideal fluid perfectly. Note that $> 99\%$ of all final hadrons have $p_T < 1.5$ GeV/c. <i>Upper left:</i> The $v_2$ plotted versus transverse kinetic energy $KE_T = m_T - m_0 = \sqrt{p_T^2 + m_0^2} - m_0$ . The $v_2$ follows different universal curves for mesons and baryons. <i>Upper right:</i> When scaled by the number of valence quarks, the $v_2$ follows the same universal curve for all hadrons and for all values of scaled transverse kinetic energy [13]. . . . .	12
655			
656			
657			
658			
659			
660			
661			
662	1.11	Invariant yield of particles versus transverse mass $m_T = \sqrt{p_T^2 + m_0^2}$ for $\pi^\pm$ , $K^\pm$ , $p$ and $\bar{p}$ at mid-rapidity for p+p collisions (bottom) and Au+Au events from 70-80% (second bottom) to 0-5% (top) centrality [14]. . . . .	13
663			
664			
665			

666	1.12	Thermal photons spectra for the central Au+Au collisions at	
667		$\sqrt{s_{NN}} = 200$ GeV at computed within different hydrodynamical	
668		models compared with the pQCD calculations (solid line) and	
669		experimental data from PHENIX (black dots) [15]. . . . .	14
670	1.13	Left: Invariant mass spectrum of $e^+e^-$ pairs in Pb+Au collisions	
671		at 158A GeV compared to the sum coming from the hadron decays	
672		predictions. Right: The expectations coming from model calcula-	
673		tions assuming a dropping of the $\rho$ mass (blue) or a spread of the	
674		$\rho$ width in the medium (red) [16]. . . . .	15
675	1.14	Azimuthal angle difference $\Delta\phi$ distributions for different colliding	
676		systems at $\sqrt{s_{NN}} = 200$ GeV. Transverse momentum cut: $p_T > 2$	
677		GeV. For the Au+Au collisions the away-side jet is missing [17]. . .	16
678	3.1	Bertsch-Pratt direction naming convention used in heavy ion col-	
679		lision. . . . .	22
680	3.2	The pair wave function is a superposition of all possible states. In	
681		case of particle interferometry it includes two cases: particles with	
682		momenta $p_1, p_2$ registered by detectors $A, B$ and $p_1, p_2$ registered	
683		by $B, A$ respectively. . . . .	23
684	3.3	An averaged three-dimensional Gaussian source function with dif-	
685		ferent widths was averaged into one-dimensional function. To il-	
686		lustrate deformations, one-dimensional Gaussian distribution was	
687		fitted. . . . .	26