



Calculation of predictions for non-identical particle correlations in AA collisions at LHC energies from hydrodynamics-inspired models

MASTER OF SCIENCE THESIS

Author:

Mateusz Wojciech Gałążyn

Supervisor:

Prof. Adam Kisiel

Warsaw, 28th July 2014



Obliczenia teoretycznych przewidywań korelacji cząstek nieidentycznych w zderzeniach AA przy energiach LHC pochodzących z modeli hydrodynamicznych

PRACA MAGISTERSKA

Autor:

Mateusz Wojciech Gałążyn

Promotor:

dr hab. inż. Adam Kisiel, prof. PW

Warszawa, 28 lipca 2014

Abstract

Streszczenie

Contents

4	1 Theory of heavy ion collisions	2
5	1.1 The Standard Model	2
6	1.2 Quantum Chromodynamics	3
7	1.2.1 Quarks and gluons	3
8	1.2.2 Quantum Chromodynamics potential	4
9	1.2.3 The quark-gluon plasma	5
10	1.3 Relativistic heavy ion collisions	6
11	2 Terminator model	7
12	2.1 (3+1)-dimensional viscous hydrodynamics	7
13	2.2 Statistical hadronization	8
14	2.2.1 Cooper-Frye formalism	9
15	3 Particle interferometry	11
16	3.1 HBT interferometry	11
17	3.2 Intensity interferometry in heavy ion collisions	11
18	3.2.1 Theoretical approach	11
19	3.2.2 Experimental approach	11
20	3.3 Scaling of femtoscopic radii	11
21	4 Results	12
22	4.1 Identical particles correlations	12
23	4.2 Results of the fit	12
24	4.3 Discussion of results	12
25	5 Summary	13

26 Introduction

Chapter 1

Theory of heavy ion collisions

1.1 The Standard Model

In the 1970s, a new theory of fundamental particles and their interaction emerged. A new concept, which concerns the electromagnetic, weak and strong nuclear interactions between know particles. This theory is called *The Standard Model*. There are seventeen named particles in the standard model, organized into the chart shown below (Fig. 1.1). Fundamental particles are divided into two families: *fermions* and *bosons*.




Figure 1.1: The Standard Model of elementary particles [1].

Fermions are the building blocks of matter. They are divided into two groups. Six of them, which must bind together are called *quarks*. Quarks are known to bind into doublets (*mesons*), triplets (*baryons*) and recently confirmed four-quark states.¹ Two of baryons, with the longest lifetimes, are forming a nucleus: a proton and a neutron. A proton is build from two up quarks and one down, and neutron consists of two down quarks and one up. A proton is found to be a stable particle (at least it has a lifetime larger than 10^{35} years) and a free neutron has a mean lifetime about 8.8×10^2 s. Fermions, that can exist independently are called *leptons*. Neutrinos are a subgroup of leptons, which are only influenced by weak interaction. Fermions can be divided into three generations (three columns in the Figure 1.1). Generation I particles can combine into hadrons with the longest life spans. Generation II and III consists of unstable particles which form also unstable hadrons.

Bosons are force carriers. There are four fundamental forces: weak - responsible for radioactive decay, strong - coupling quarks into hadrons, electromagnetic - between charged particles and gravity - the weakest, which causes the attraction between particles with a mass. The Standard Model describes the first three. The weak force is mediated by W^\pm and Z^0 bosons, electromagnetic force is carried by photons γ and the carriers of a strong interaction are gluons g . The fifth boson is a Higgs boson which is responsible for giving other particles mass.

1.2 Quantum Chromodynamics

1.2.1 Quarks and gluons

Quarks interact with each other through the strong interaction. The mediator of this force is a *gluon* - a massless and chargeless particle. In the quantum chromodynamics (QCD) - theory describing strong interaction - there are six types of "charges" (like electrical charges in the electrodynamics) called *colours*. The colors were introduced because some of the observed particles, like Δ^- , Δ^{++} and Ω^- appeared to consist of three quarks with the same flavour (*ddd*, *uuu* and *sss* respectively), which was in conflict with the Pauli principle. One quark can carry one of the three colors (usually called *red*, *green* and *blue*) and antiquark one of the three anti-colors respectively. Only color-neutral (or white) particles could exist, mesons are assumed to be a color-anticolor pair, while baryons are *red-green-blue* triplets. Gluons also are color-charged and there are 8 types of gluons. Therefore they can interact with themselves [3].

¹The LHCb experiment at CERN in Geneva confirmed recently existence of $Z(4430)$ - a particle consisting of four quarks [2].

1.2.2 Quantum Chromodynamics potential

As a result of that gluons are massless, one can expect, that the static potential in the QCD will have the similar form like one in the electrodynamics e.g. $\sim 1/r$. In reality the QCD potential is assumed to have the form of [3]

$$V_s = -\frac{4}{3} \frac{\alpha_s}{r} + kr, \quad (1.1)$$

where the α_s is a coupling constant of the strong force and the kr part is related with the *confinement*. In comparison to the electromagnetic force, a value of the strong coupling constant is $\alpha_s \approx 1$ and the electromagnetic one is $\alpha = 1/137$.




Figure 1.2: A string break and a creation of a pair quark-anti-quark [4].

The fact that quarks does not exist separately, but they are always bound, is called a confinement. As two quarks are pulled apart, the distance and the potential grow. This situation resembles stretching of a string. At some point, when the string is so large it is energetically favourable to create a quark-antiquark pair. At this moment such pair (or pairs) is formed, the string breaks and the confinement is preserved (Fig. 1.2).




Figure 1.3: The QCD potential for a pair quark-antiquark as a function of distance for different temperatures. [4].

For the small distances, an interaction between the quarks and gluons is weakening and they are starting to behave like free particles. This phenomenon is known as an *asymptotic freedom*. The QCD potential has also temperature dependence - the force strength “melts” with the temperature increase. Therefore the asymptotic freedom is expected to appear in either the case of high baryon densities (small distances between quarks) or very high temperatures. This temperature dependence is illustrated in the Fig. 1.3.

1.2.3 The quark-gluon plasma

The new state of matter in which quarks are no longer confined is known as *quark-gluon plasma* (QGP). The predictions coming from the discrete space-time Lattice QCD calculations reveal a phase transition from the hadronic matter to the quark-gluon plasma at the high temperatures and baryon densities. The results




Figure 1.4: A number of degrees of freedom as a function of a temperature [5].

obtained from such calculations are shown on Fig. 1.4. The energy density ϵ which is divided by T^4 is a measure of number of degrees of freedom in the system. One can observe significant rise of this value, when the temperature increases past the critical value T_C . Such increase is signaling of a phase transition - the formation of QGP [6]. The values of the energy densities plotted in Fig. 1.4 do not reach the Stefan-Boltzmann limit ϵ_{SB} (marked with arrows), which corresponds to the ideal gas. This can indicate some residual interactions in the system. According to the results from the RHIC², the new phase of matter behaves more like an ideal fluid, than like a gas [7].

One of the key questions, to which current heavy ion physics tries to find an answer is the value of a critical temperature T_C as a function of a baryon chemical potential μ_B (baryon density), where the phase transition occurs. The

²Relativistic Heavy Ion Collider at Brookhaven National Laboratory in Upton, New York




Figure 1.5: Phase diagram coming from the Lattice QCD calculations [6].

106 results coming from the Lattice QCD are presented in the Fig. 1.5. The phase of
 107 matter in which quarks and gluons are deconfined is expected to exist at large
 108 temperatures. In the region of small temperatures and high baryon densities, a
 109 different state is supposed to appear - a *colour superconductor*. The phase trans-
 110 ition between hadronic matter and QGP is thought to be of 1st order at $\mu_B \gg 0$.
 111 However as $\mu_B \rightarrow 0$ quarks' masses become significant and a sharp transition
 112 transforms into a rapid but smooth cross-over. It is believed that in Pb-Pb colli-
 113 sions observed at the LHC³, the created matter has high enough temperature to
 114 be in the quark-gluon plasma phase, then cools down and converts into hadrons,
 115 undergoing a smooth transition [6].

116 1.3 Relativistic heavy ion collisions

³Large Hadron Collider at CERN, Geneva

117 Chapter 2

118 Terminator model

119 THERMINATOR [8] is a Monte Carlo event generator designed to investigate
120 the particle production in the relativistic heavy ion collisions. The functionality
121 of the code includes a generation of the stable particles and unstable resonances
122 at the chosen hypersurface model. It performs the statistical hadronization which
123 is followed by space-time evolution of particles and the decay of resonances. The
124 key element of this method is an inclusion of a complete list of hadronic reson-
125 ances, which contribute very significantly to the observables. The second version
126 of THERMINATOR [9] comes with a possibility to incorporate any shape of freeze-
127 out hypersurface and the expansion velocity field, especially those generated ex-
128 ternally with various hydrodynamic codes.

129 2.1 (3+1)-dimensional viscous hydrodynamics

130 Most of the relativistic viscous hydrodynamic calculations are done in
131 (2+1)-dimensions. Such simplification assumes boost-invariance of a matter
132 created in a collision. Experimental data reveals that no boost-invariant region is
133 formed in the collisions [10]. Hence, for the better description of created system
134 a (3+1)-dimensional model is required.

135 In the four dimensional relativistic dynamics one can describe a system
136 using a space-time four-vector $x^\nu = (ct, x, y, z)$, a velocity four-vector
137 $u^\nu = \gamma(c, v_x, v_y, v_z)$ and a energy-momentum tensor $T^{\mu\nu}$. The particular
138 components of $T^{\mu\nu}$ have a following meaning:

- 139 • T^{00} - an energy density,
- 140 • $cT^{0\alpha}$ - an energy flux across a surface x^α ,
- 141 • $T^{\alpha 0}$ - an α -momentum flux across a surface x^α multiplied by c ,
- 142 • $T^{\alpha\beta}$ - components of momentum flux density tensor,

where $\gamma = (1 - v^2/c^2)^{-1/2}$ is Lorentz factor and $\alpha, \beta \in \{1, 2, 3\}$. Using u^ν one can express $T^{\mu\nu}$ as follows [11]:

$$T_0^{\mu\nu} = (e + p)u^\mu u^\nu - pg^{\mu\nu} \quad (2.1)$$

where e is an energy density, p is a pressure and $g^{\mu\nu}$ is an inverse metric tensor:

$$g^{\mu\nu} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}. \quad (2.2)$$

The presented version of energy-momentum tensor (2.1) can be used to describe dynamics of a perfect fluid. To take into account influence of viscosity, one has to apply the following corrections coming from shear $\pi^{\mu\nu}$ and bulk Π viscosities [12]:

$$T^{\mu\nu} = T_0^{\mu\nu} + \pi^{\mu\nu} + \Pi(g^{\mu\nu} - u^\mu u^\nu). \quad (2.3)$$

The stress tensor $\pi^{\mu\nu}$ and the bulk viscosity Π are solutions of dynamical equations in the second order viscous hydrodynamic framework [11]. The comparison of hydrodynamics calculations with the experimental results reveal, that the shear viscosity divided by entropy η/s has to be small and close to the AdS/CFT estimate $\eta/s = 0.08$ [12, 13].

When using $T^{\mu\nu}$ to describe system evolving close to local thermodynamic equilibrium, relativistic hydrodynamic equations in a form of:

$$\partial_\mu T^{\mu\nu} = 0 \quad (2.4)$$

can be used to describe the dynamics of the local energy density, pressure and flow velocity.

Hydrodynamic calculations are starting from the Glauber¹ model initial conditions. The collective expansion of a fluid ends at the freeze-out hypersurface. That surface is usually defined as a constant temperature surface, or equivalently as a cut-off in local energy density. The freeze-out is assumed to occur at the temperature $T = 140$ MeV.

2.2 Statistical hadronization

Statistical description of heavy ion collision has been successfully used to describe quantitatively *soft* physics, i.e. the regime with the transverse momentum not exceeding 2 GeV. The basic assumption of the statistical approach of evolution of the quark-gluon plasma is that at some point of the space-time evolution of the fireball, the thermal equilibrium is reached. When

¹The Glauber Model is used to calculate “geometrical” parameters of a collision like an impact parameter, number of participating nucleons or number of binary collisions.

the system is in the thermal equilibrium the local phase-space densities of particles follow the Fermi-Dirac or Bose-Einstein statistical distributions. At the end of the plasma expansion, the freeze-out occurs. The freeze-out model incorporated in the THERMINATOR model assumes, that chemical and thermal freeze-out occur at the same time.

2.2.1 Cooper-Frye formalism

The result of the hydrodynamic calculations is the freeze-out hypersurface Σ^μ . A three-dimensional element of the surface is defined as [9]

$$d\Sigma_\mu = \epsilon_{\mu\alpha\beta\gamma} \frac{\partial x^\alpha}{\partial \alpha} \frac{\partial x^\beta}{\partial \beta} \frac{\partial x^\gamma}{\partial \gamma} d\alpha d\beta d\gamma, \quad (2.5)$$

where $\epsilon_{\mu\alpha\beta\gamma}$ is the Levi-Civita tensor and the variables $\alpha, \beta, \gamma \in \{1, 2, 3\}$ are used to parametrize the three-dimensional freeze-out hypersurface in the Minkowski four-dimensional space. The Levi-Civita tensor is equal to 1 when the indices form an even permutation (eg. ϵ_{0123}), to -1 when the permutation is odd (e.g. ϵ_{2134}) and has a value of 0 if any index is repeated. Therefore [9],

$$d\Sigma_0 = \begin{vmatrix} \frac{\partial x}{\partial \alpha} & \frac{\partial x}{\partial \beta} & \frac{\partial x}{\partial \gamma} \\ \frac{\partial y}{\partial \alpha} & \frac{\partial y}{\partial \beta} & \frac{\partial y}{\partial \gamma} \\ \frac{\partial z}{\partial \alpha} & \frac{\partial z}{\partial \beta} & \frac{\partial z}{\partial \gamma} \end{vmatrix} d\alpha d\beta d\gamma \quad (2.6)$$

and the remaining components are obtained by cyclic permutations of t, x, y and z .

One can obtain the number of hadrons produced on the hypersurface Σ^μ from the Cooper-Frye formalism. The following integral yields the total number of created particles [9]:

$$N = (2s + 1) \int \frac{d^3p}{(2\pi)^3 E_p} \int d\Sigma_\mu(x) p^\mu f(x, p), \quad (2.7)$$

where

$$f(p \cdot u) = \left\{ \exp \left[\frac{p_\mu u^\mu - (B\mu_B + I_3\mu_{I_3} + S\mu_S + C\mu_C)}{T} \right] \pm 1 \right\}^{-1} \quad (2.8)$$

is the phase-space distribution for particles (for stable ones and resonances). For the Fermi-Dirac distribution in the 2.8 there is a plus sign and for Bose-Einstein statistics minus sign respectively. The thermodynamic quantities appearing in the $f(\cdot)$ are T - temperature, μ_B - baryon chemical potential, μ_{I_3} - isospin chemical potential, μ_S - strange chemical potential, μ_C - charmed chemical potential and the s is a spin of a particle. One can simply derive from equation 2.7, the dependence of the momentum density [14]:

$$E \frac{dN}{d^3p} = \int f(x, p) p^\mu d\Sigma_\mu. \quad (2.9)$$

197 The equations presented above are directly used in the THERMINATOR to generate
198 the hadrons with the Monte-Carlo method.

199 **Chapter 3**

200 **Particle interferometry**

201 **3.1 HBT interferometry**

202 **3.2 Intensity interferometry in heavy ion collisions**

203 **3.2.1 Theoretical approach**

204 **Two particle wave function**

205 **Source function**

206 **Theoretical correlation function**

207 **Spherical harmonics decomposition of correlation function**

208 **3.2.2 Experimental approach**

209 **3.3 Scaling of femtoscopic radii**

210 **Chapter 4**

211 **Results**

212 **4.1 Identical particles correlations**

213 **4.2 Results of the fit**

214 **4.3 Discussion of results**

215 **Chapter 5**

216 **Summary**

Bibliography

- [1] Standard Model of Elementary Particles - Wikipedia, the free encyclopedia
http://en.wikipedia.org/wiki/standard_model.
- [2] R. Aaij et al. (LHCb Collaboration). Observation of the resonant character of the $z(4430)^-$ state. *Phys. Rev. Lett.*, 112:222002, Jun 2014.
- [3] Donald H. Perkins. *Introduction to High Energy Physics*. Cambridge University Press, fourth edition, 2000. Cambridge Books Online.
- [4] G. Odyniec. *Phase Diagram of Quantum Chromo-Dynamics* - course at Faculty of Physics, Warsaw University of Technology, Jun 2012.
- [5] F. Karsch. Lattice results on QCD thermodynamics. *Nuclear Physics A*, 698(1-4):199 – 208, 2002.
- [6] Adam Kisiel. *Studies of non-identical meson-meson correlations at low relative velocities in relativistic heavy-ion collisions registered in the STAR experiment*. PhD thesis, Warsaw University of Technology, Aug 2004.
- [7] J. Bartke. *Relativistic Heavy Ion Physics*. World Scientific Pub., 2009.
- [8] Adam Kisiel, Tomasz Taluc, Wojciech Broniowski, and Wojciech Florkowski. THERMINATOR: THERMal heavy-IoN generATOR. *Comput.Phys.Commun.*, 174:669–687, 2006.
- [9] Mikolaj Chojnacki, Adam Kisiel, Wojciech Florkowski, and Wojciech Broniowski. THERMINATOR 2: THERMal heavy IoN generATOR 2. *Comput.Phys.Commun.*, 183:746–773, 2012.
- [10] I. et al (BRAHMS Collaboration) Bearden. Charged meson rapidity distributions in central Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV. *Phys. Rev. Lett.*, 94:162301, Apr 2005.
- [11] W. Israel and J.M. Stewart. Transient relativistic thermodynamics and kinetic theory. *Annals of Physics*, 118(2):341 – 372, 1979.
- [12] Piotr Bozek. Flow and interferometry in (3 + 1)-dimensional viscous hydrodynamics. *Phys. Rev. C*, 85:034901, Mar 2012.

- 245 [13] K. Kovtun, P. D. T. Son, and A. O. Starinets. Viscosity in strongly interacting
246 quantum field theories from black hole physics. *Phys. Rev. Lett.*, 94:111601,
247 Mar 2005.
- 248 [14] Fred Cooper and Graham Frye. Single-particle distribution in the hydro-
249 dynamic and statistical thermodynamic models of multiparticle production.
250 *Phys. Rev. D*, 10:186–189, Jul 1974.

251 List of Figures

252	1.1	The Standard Model of elementary particles [1].	2
253	1.2	A string break and a creation of a pair quark-anti-quark [4].	4
254	1.3	The QCD potential for a pair quark-antiquark as a function of dis-	
255		tance for different temperatures. [4].	4
256	1.4	A number of degrees of freedom as a function of a temperature [5].	5
257	1.5	Phase diagram coming from the Lattice QCD calculations [6]. . . .	6