Type C1 (chirp plane streak 211), 3.56 kHz @ chirps 3.1 Hz at 25 °C

rows: positions along the line, cold and slow at the top, warm and fast at the bottom
columns: singers at that position, distinct recorder-nights, by air temperature




Figure S7-C1 (rate-filtered). Type C1 (chirp plane streak 211): 3.56 kHz @ chirps 3.1 Hz at 25 °C, 62 singer-nights. RATE-FILTERED VIEW of the grid: each cell is the period-synchronous average of the excerpt over 5 periods at the singer's own rate, so the singer's syllables stand out and neighbors at other rates smear. Rows are positions along the line from the cold, slow end at the top to the warm, fast end at the bottom; columns are singers at that position from distinct recorder-nights, ordered by air temperature. Each cell is 1 s of the loudest 30 s of the singer's deepest window on one shared log-frequency axis, the detected band in color (dB above the row's quiet level, 0 to 30) and the rest in grayscale; a cell whose singer had no syllable rate in that window is folded at its chirp rate. The cell head gives the night, recorder, site, air temperature and rate (c: chirp rate). Top right: the type on the plane, and the row positions on the streak, in order.

Type C2 (chirp plane streak 21), 3.94 kHz @ chirps 4.2 Hz at 25 °C

rows: positions along the line, cold and slow at the top, warm and fast at the bottom
columns: singers at that position, distinct recorder-nights, by air temperature




Figure S7-C2 (rate-filtered). Type C2 (chirp plane streak 21): 3.94 kHz @ chirps 4.2 Hz at 25 °C, 351 singer-nights. RATE-FILTERED VIEW of the grid: each cell is the period-synchronous average of the excerpt over 5 periods at the singer's own rate, so the singer's syllables stand out and neighbors at other rates smear. Rows are positions along the line from the cold, slow end at the top to the warm, fast end at the bottom; columns are singers at that position from distinct recorder-nights, ordered by air temperature. Each cell is 1 s of the loudest 30 s of the singer's deepest window on one shared log-frequency axis, the detected band in color (dB above the row's quiet level, 0 to 30) and the rest in grayscale; a cell whose singer had no syllable rate in that window is folded at its chirp rate. The cell head gives the night, recorder, site, air temperature and rate (c: chirp rate). Top right: the type on the plane, and the row positions on the streak, in order.




Figure S7-C3 (rate-filtered). Type C3 (chirp plane streak 10): 5.75 kHz @ chirps 3.4 Hz at 25 °C, 55 singer-nights. RATE-FILTERED VIEW of the grid: each cell is the period-synchronous average of the excerpt over 5 periods at the singer's own rate, so the singer's syllables stand out and neighbors at other rates smear. Rows are positions along the line from the cold, slow end at the top to the warm, fast end at the bottom; columns are singers at that position from distinct recorder-nights, ordered by air temperature. Each cell is 1 s of the loudest 30 s of the singer's deepest window on one shared log-frequency axis, the detected band in color (dB above the row's quiet level, 0 to 30) and the rest in grayscale; a cell whose singer had no syllable rate in that window is folded at its chirp rate. The cell head gives the night, recorder, site, air temperature and rate (c: chirp rate). Top right: the type on the plane, and the row positions on the streak, in order.

Type C4 (chirp plane streak 9), 6.16 kHz @ chirps 3.0 Hz at 25 °C

rows: positions along the line, cold and slow at the top, warm and fast at the bottom
columns: singers at that position, distinct recorder-nights, by air temperature




Figure S7-C4 (rate-filtered). Type C4 (chirp plane streak 9): 6.16 kHz @ chirps 3.0 Hz at 25 °C, 90 singer-nights. RATE-FILTERED VIEW of the grid: each cell is the period-synchronous average of the excerpt over 5 periods at the singer's own rate, so the singer's syllables stand out and neighbors at other rates smear. Rows are positions along the line from the cold, slow end at the top to the warm, fast end at the bottom; columns are singers at that position from distinct recorder-nights, ordered by air temperature. Each cell is 1 s of the loudest 30 s of the singer's deepest window on one shared log-frequency axis, the detected band in color (dB above the row's quiet level, 0 to 30) and the rest in grayscale; a cell whose singer had no syllable rate in that window is folded at its chirp rate. The cell head gives the night, recorder, site, air temperature and rate (c: chirp rate). Top right: the type on the plane, and the row positions on the streak, in order.

Type C5 (chirp plane streak 91), 6.44 kHz @ chirps 3.2 Hz at 25 °C

rows: positions along the line, cold and slow at the top, warm and fast at the bottom
columns: singers at that position, distinct recorder-nights, by air temperature




Figure S7-C5 (rate-filtered). Type C5 (chirp plane streak 91): 6.44 kHz @ chirps 3.2 Hz at 25 °C, 48 singer-nights. RATE-FILTERED VIEW of the grid: each cell is the period-synchronous average of the excerpt over 5 periods at the singer's own rate, so the singer's syllables stand out and neighbors at other rates smear. Rows are positions along the line from the cold, slow end at the top to the warm, fast end at the bottom; columns are singers at that position from distinct recorder-nights, ordered by air temperature. Each cell is 1 s of the loudest 30 s of the singer's deepest window on one shared log-frequency axis, the detected band in color (dB above the row's quiet level, 0 to 30) and the rest in grayscale; a cell whose singer had no syllable rate in that window is folded at its chirp rate. The cell head gives the night, recorder, site, air temperature and rate (c: chirp rate). Top right: the type on the plane, and the row positions on the streak, in order.

Type C6 (chirp plane streak 5), 6.84 kHz @ chirps 2.9 Hz at 25 °C

rows: positions along the line, cold and slow at the top, warm and fast at the bottom
columns: singers at that position, distinct recorder-nights, by air temperature




Figure S7-C6 (rate-filtered). Type C6 (chirp plane streak 5): 6.84 kHz @ chirps 2.9 Hz at 25 °C, 365 singer-nights. RATE-FILTERED VIEW of the grid: each cell is the period-synchronous average of the excerpt over 5 periods at the singer's own rate, so the singer's syllables stand out and neighbors at other rates smear. Rows are positions along the line from the cold, slow end at the top to the warm, fast end at the bottom; columns are singers at that position from distinct recorder-nights, ordered by air temperature. Each cell is 1 s of the loudest 30 s of the singer's deepest window on one shared log-frequency axis, the detected band in color (dB above the row's quiet level, 0 to 30) and the rest in grayscale; a cell whose singer had no syllable rate in that window is folded at its chirp rate. The cell head gives the night, recorder, site, air temperature and rate (c: chirp rate). Top right: the type on the plane, and the row positions on the streak, in order.

Type C7 (chirp plane streak 4), 8.07 kHz @ chirps 3.4 Hz at 25 °C

rows: positions along the line, cold and slow at the top, warm and fast at the bottom
columns: singers at that position, distinct recorder-nights, by air temperature




Figure S7-C7 (rate-filtered). Type C7 (chirp plane streak 4): 8.07 kHz @ chirps 3.4 Hz at 25 °C, 2659 singer-nights. RATE-FILTERED VIEW of the grid: each cell is the period-synchronous average of the excerpt over 5 periods at the singer's own rate, so the singer's syllables stand out and neighbors at other rates smear. Rows are positions along the line from the cold, slow end at the top to the warm, fast end at the bottom; columns are singers at that position from distinct recorder-nights, ordered by air temperature. Each cell is 1 s of the loudest 30 s of the singer's deepest window on one shared log-frequency axis, the detected band in color (dB above the row's quiet level, 0 to 30) and the rest in grayscale; a cell whose singer had no syllable rate in that window is folded at its chirp rate. The cell head gives the night, recorder, site, air temperature and rate (c: chirp rate). Top right: the type on the plane, and the row positions on the streak, in order.

Type C8 (chirp plane streak 7), 8.71 kHz @ chirps 2.7 Hz at 25 °C

rows: positions along the line, cold and slow at the top, warm and fast at the bottom
columns: singers at that position, distinct recorder-nights, by air temperature




Figure S7-C8 (rate-filtered). Type C8 (chirp plane streak 7): 8.71 kHz @ chirps 2.7 Hz at 25 °C, 101 singer-nights. RATE-FILTERED VIEW of the grid: each cell is the period-synchronous average of the excerpt over 5 periods at the singer's own rate, so the singer's syllables stand out and neighbors at other rates smear. Rows are positions along the line from the cold, slow end at the top to the warm, fast end at the bottom; columns are singers at that position from distinct recorder-nights, ordered by air temperature. Each cell is 1 s of the loudest 30 s of the singer's deepest window on one shared log-frequency axis, the detected band in color (dB above the row's quiet level, 0 to 30) and the rest in grayscale; a cell whose singer had no syllable rate in that window is folded at its chirp rate. The cell head gives the night, recorder, site, air temperature and rate (c: chirp rate). Top right: the type on the plane, and the row positions on the streak, in order.