

Methods for the Detection of the Epoch of Reionization  
by Interferometers Measuring the 21 cm Signal

By

Carina Cheng

A dissertation submitted in partial satisfaction of the

requirements for the degree of

Doctor of Philosophy

in

Astrophysics

in the

Graduate Division

of the

University of California, Berkeley

Committee in charge:

Professor Aaron Parsons, Chair  
Professor Stuart Bale  
Professor Mariska Kriek

Spring 2019

Methods for the Detection of the Epoch of Reionization  
by Interferometers Measuring the 21 cm Signal

Copyright 2019  
by  
Carina Cheng

## Abstract

### Methods for the Detection of the Epoch of Reionization by Interferometers Measuring the 21 cm Signal

by

Carina Cheng

Doctor of Philosophy in Astrophysics

University of California, Berkeley

Professor Aaron Parsons, Chair

The Epoch of Reionization is one of the last unexplored eras of our Universe's history. Beginning about a billion years after the Big Bang, this epoch is characterized by the births of the first stars and galaxies, whose light subsequently altered the nature of the gas surrounding them. There are several experiments aiming to detect the phase transition of this gas as it changes from neutral hydrogen to ionized hydrogen. Such a detection would open a wealth of information about our early Universe, revealing details about the nature of the first luminous sources and the evolution of structure formation.

Interferometers such as the Precision Array for Probing the Epoch of Reionization (PAPER) and the Hydrogen Epoch of Reionization Array (HERA) seek to measure the 21 cm signal from neutral hydrogen and map its evolution over spatial and temporal scales. A successful detection of reionization, however, is a difficult measurement. Though the 21 cm signal is a powerful topological probe of the intergalactic medium, it is easily buried underneath bright foreground signals and instrumental systematics. A clean detection of reionization is ambitious and requires analysis methods that maximize data sensitivity and increase confidence in results.

The work presented in this thesis addresses many of the key challenges that face the current field of 21 cm cosmology. This includes algorithms to locate contaminated data, methods to ensure accurate power spectrum measurements, and techniques for removing unwanted systematics while preserving the reionization signal. These developments serve as the foundation of the latest 21 cm measurements from the PAPER-64 and PAPER-128 arrays, whose results lie at the forefront of the field. These methods are also fundamental to HERA and future experiments, as they provide a strong foundation for the continued exploration of our cosmic dawn.

*to my parents*

# Contents

<b>List of Figures</b>	<b>iv</b>
<b>List of Tables</b>	<b>v</b>
<b>Acknowledgments</b>	<b>vi</b>
<b>1 Introduction</b>	<b>1</b>
1.1 The Epoch of Reionization . . . . .	1
1.1.1 Cosmic History . . . . .	1
1.1.2 CMB and Galaxy Measurements . . . . .	2
1.1.3 Measurements of HI . . . . .	4
1.1.4 This Thesis . . . . .	7
1.2 Interferometry . . . . .	7
1.2.1 The Visibility Equation . . . . .	7
1.2.2 Calibration . . . . .	7
1.2.3 Foreground and Filtering . . . . .	7
1.2.4 Fringe-rate Filtering . . . . .	7
1.3 Instruments . . . . .	7
1.3.1 The Precision Array for Probing the Epoch of Reionization . . . . .	7
1.3.2 The Hydrogen Epoch of Reionization Array . . . . .	7
1.3.3 Other Experiments . . . . .	7
<b>2 Power Spectrum Methods</b>	<b>8</b>
2.1 The 21 cm Power Spectrum . . . . .	8
2.2 Signal Loss . . . . .	8
2.3 Error Estimation . . . . .	8
2.4 Bias . . . . .	8
<b>3 PAPER-64</b>	<b>9</b>
3.1 Overview . . . . .	9
3.2 Signal Loss . . . . .	9
3.3 Error Estimation . . . . .	9
3.4 Bias . . . . .	9

---

3.5	Power Spectrum Limits . . . . .	9
<b>4</b>	<b>PAPER-128</b>	<b>10</b>
4.1	Overview . . . . .	10
4.2	Quality Assurance . . . . .	10
4.2.1	Flagging Julian Dates . . . . .	10
4.2.2	Flagging Antennas . . . . .	10
<b>5</b>	<b>HERA</b>	<b>11</b>
<b>6</b>	<b>Conclusion</b>	<b>12</b>
	<b>Bibliography</b>	<b>13</b>

# List of Figures

1.1	Timeline of the history of the Universe. The Epoch of Reionization marks the era when the first stars and galaxies formed and ionized the neutral hydrogen in the Universe. Image credit: NAOJ. . . . .	2
1.2	A cartoon diagram of the observable Universe centered on us. Close-by, galaxy observations have mapped out cosmic web structure in our nearby Universe (image credit: SDSS). Far-away, the cosmic microwave background is observed at a redshift of $z \sim 1100$ (image credit: WMAP). The Epoch of Reionization represents a mostly unexplored era between the two, and can be probed by measuring red-shifted 21 cm radiation from neutral hydrogen. . . . .	5
1.3	The evolution of the global 21 cm signal, starting with the Dark Ages, through galaxy formation and reionization (image credit: Pritchard & Loeb (2012)). The work in this thesis mainly focuses on a redshift range of $6 < z < 12$ as reionization is expected to progress and complete. . . . .	5

# List of Tables



# Acknowledgments

# Chapter 1

## Introduction

### 1.1 The Epoch of Reionization


Our Universe has a complex, rich history, of which enormous progress has been made in the past few decades to unravel its story. Much has been learned about the very beginnings of the Universe, from the Big Bang’s large explosion of energy to the relatively smooth and simple cosmic background radiation that was leftover. Additionally, observational feats have revealed the status of the present-day Universe and the intricate *cosmic web*, or large scale structure, of galaxies today.

The Epoch of Reionization (EoR) ties these two bookends together, occurring about a billion years after the Big Bang when the very first stars and galaxies formed. How did the tiny density fluctuations from the cosmic microwave background develop into the structure we see today? How did the first luminous structures form, and how did they evolve and influence the gas around them? Exploring the reionization era opens up a new chapter of our Universe’s story - a chapter that promises to connect the dots between our past and present.

#### 1.1.1 Cosmic History

As the Universe expanded and cooled after the Big Bang, electrons and protons eventually combined to form neutral hydrogen atoms. At the young age of  $\sim 380,000$  years, the Universe’s ordinary baryonic content was almost entirely neutral hydrogen, while most of its total matter was dark matter (Loeb & Furlanetto 2013). Then, for the next several hundred million years, the *Dark Ages* proceeded, with concentrations of dark matter setting the foundations for the formation of the first luminous structures. More specifically, the tiny, primordial density fluctuations that were established at the release of the CMB grew with inflation and the expansion of the Universe. The densest regions then collapsed to form dark matter halos, inside of which hydrogen gas could cool, condense, and fragment into stars.

The first luminous structures are thought to have formed at an age of  $\sim 200$  million years ( $z \sim 20$ ) and are predicted to have been massive stars with high luminosities and



*Figure 1.1:* Timeline of the history of the Universe. The Epoch of Reionization marks the era when the first stars and galaxies formed and ionized the neutral hydrogen in the Universe. Image credit: NAOJ.

large ionizing powers (Loeb & Furlanetto 2013). The ionizing photons from the first stars carved out pockets of ionized hydrogen gas, and as the number of sources grew, an increased number of ionized bubbles emerged. Eventually, the first generations of stars and galaxies are responsible for ionizing all the neutral hydrogen in the Universe, with reionization complete by about one billion years after the Big Bang ( $z \sim 6$ ) (Furlanetto et al. 2006).

The exact timescale and details of the reionization process, which are shown within the context of the history of the Universe in Figure 1.1, are current research questions in the field of cosmology. The physics of reionization depends on several factors, including the nature of the first stars (masses, luminosities, ionizing photons) and the surrounding gas (efficiency of the ionizing photons, feedback effects). There are several ways to approach the investigation of this era, with CMB measurements working to constrain the duration of reionization, galaxy measurements unveiling the end of reionization, and direct hydrogen measurements attempting to map out the changing nature of the gas over time. All of these probes serve to illuminate this watershed era between a Universe dominated by darkness and a Universe defined by light.

### 1.1.2 CMB and Galaxy Measurements

There are several observational probes of the reionization epoch, three of which we highlight in this section. The first is the study of CMB anisotropies, which carry with them an imprint of the early Universe. But that's not the only imprint it has - CMB photons can scatter off of free electrons after reionization, and these scatterings leave behind polarization and temperature imprints (Haiman & Knox 1999). For example, the amplitude of the CMB is sensitive to scatterings, as an increased number of scatterings is akin to mixing different parts of the CMB together as photons are scattered in all directions, or in other words, this

scattering washes out anisotropies in the CMB and lowers its overall amplitude.

A useful parameter to quantify the amount of electron scattering that occurs is the optical depth,  $\tau_{es}$ , defined as:

$$\tau_{es} = \int n_e \sigma_T dl, \quad (1.1)$$

where  $n_e$  is the number density of free electrons,  $\sigma_T$  is the Thompson cross-section, and the integral is taken over a proper length  $dl$ . Once reionization begins, the number of free electrons increases, contributing to increasingly higher values of  $\tau_{es}$ . Hence, an earlier start for reionization would yield higher optical depths than a late reionization scenario.

Observations of the CMB by WMAP and Planck have placed constraints on the optical depth parameter (Hinshaw et al. 2013; Planck Collaboration et al. 2016), with the more recent Planck result suggesting a lower value of  $\tau_{es} \sim 0.07$  than WMAP. This value suggests that reionization ends at a redshift of  $z \sim 6$ , with instantaneous reionization (“mean” reionization) at  $z \sim 8.8$  (Planck Collaboration et al. 2016).

Currently, the results from CMB measurements are in agreement with a second powerful probe of EoR — that is, observations of high-redshift galaxies. This probe comes in many flavors. For example, the spectra of distant quasars can illuminate the end of reionization. Quasars, being extraordinarily bright and energetic objects, are detectable at very far distances and their spectra reveal the amount of absorption their light has undergone due to neutral hydrogen. While nearby quasar spectra exhibit sharp absorption lines, distant ones show the Gunn-Peterson trough, implying that the quasar light was entirely suppressed by hydrogen absorption. Studying the absorption features of quasars at different redshifts implies that reionization has indeed ended by  $z \sim 6$  (Becker et al. 2001).

In addition to quasar observations, high-redshift galaxy observations can also reveal important characteristics about the state of the intergalactic medium (IGM). Namely, distant star-forming galaxies can be detected using a variety of techniques, such as narrow-band imaging to find Lyman- $\alpha$  emitters (produced by recombination near young stars) or broad-band observations to find Lyman-break galaxies (spectral breaks associated with absorption by neutral hydrogen). High-redshift galaxy observations can then be used to construct luminosity functions (number of stars per luminosity interval) and star formation histories, which in turn impact the evolution of the IGM.

More specifically, if star-forming galaxies dominated the reionization process, then the ionization rate can be related to star-formation parameters as:

$$\dot{n}_{\text{ion}} = f_{\text{esc}} \xi_{\text{ion}} \rho_{\text{SFR}}, \quad (1.2)$$

where  $\dot{n}_{\text{ion}}$  is the cosmic ionization rate,  $f_{\text{esc}}$  is the escape fraction of photons into the IGM,  $\xi_{\text{ion}}$  is the rate of production of ionizing photons for a stellar population, and  $\rho_{\text{SFR}}$  is the star formation rate density. All three parameters influence the rate at which the IGM is ionized, with the latter able to be constructed from galaxy luminosity functions. For example, Robertson et al. (2015) used data from the Hubble Space Telescope to construct

a star formation rate history out to high redshifts, backing out an optical depth parameter that is consistent with that of Planck.

While galaxy measurements can be used to constrain the EoR, they are ultimately doing so by unveiling the properties of old, distant stars and galaxies. A similar, new technique that also aims to reconstruct the histories of the first luminous structures is observing nearby, metal-poor Local Group galaxies. Called “galactic archaeology,” observations of nearby star-forming ancestors can be used to constrain the faint-end slope of the luminosity function. Determining the shape of this function has important implications on the number of galaxies needed to drive reionization and the types of sources dominating this epoch (Weisz & Boylan-Kolchin 2017).

CMB measurements and galaxy observations are both important in studying the EoR. However, they each have their limitations. For example, CMB measurements can only reveal the integrated quantity of  $\tau_{es}$ , therefore unable to provide insight into the evolution of reionization as it progresses over time. Similarly, galaxy observations are currently hovering only around the tail end of the reionization era, and greater sensitivities are needed to push past this limit. A different, but complimentary, probe is needed to unlock the entire window into the EoR.


### 1.1.3 Measurements of HI

A direct measurement of neutral hydrogen gas over time would provide a fundamental way to track the IGM over the reionization process. Such a probe, which is made possible by the spin-flip transition of hydrogen, is a powerful technique that allows the tracing of gas over time, and it is this technique that serves as the basis for the remainder of this thesis.


The spin-flip transition of neutral hydrogen occurs when a hydrogen atom changes energy state between two hyperfine levels. Namely, if a hydrogen atom moves from an aligned state to an anti-aligned state, the energy difference is released in the form of a photon with a wavelength of 21 cm.

Because this transition has a well-defined wavelength, the signal can be directly mapped to a distance, or redshift, by measuring its wavelength upon detection. For example, a 21 cm photon that was initially emitted at a redshift of  $z = 6$  would have expanded by a factor of  $(1 + z)$  due to the expansion of the Universe and be 1.5 m long when it arrives at our telescopes. Hence, observing longer wavelengths of the hydrogen signal means that it has traveled for a greater distance (and has stretched out more) and thus comes from farther away at a higher redshift. This means that the 21 cm signal is a powerful tracer of neutral hydrogen at any distance (i.e. as a function of time), as long as it exists. This technique is especially compelling because it allows the direct exploration of the EoR as it occurs, whereas CMB measurements and galaxy measurements surround this era from the beginning and end only, respectively (Figure 1.2).

In practice, the 21 cm signal is encapsulated by the quantity  $T_{\text{spin}}$ , which measures the relative number of hydrogen atoms in the excited (aligned) versus ground (anti-aligned) spin-flip state. A high spin temperature means that the hydrogen gas is more likely to emit 21 cm



*Figure 1.2:* A cartoon diagram of the observable Universe centered on us. Close-by, galaxy observations have mapped out cosmic web structure in our nearby Universe (image credit: SDSS). Far-away, the cosmic microwave background is observed at a redshift of  $z \sim 1100$  (image credit: WMAP). The Epoch of Reionization represents a mostly unexplored era between the two, and can be probed by measuring red-shifted 21 cm radiation from neutral hydrogen.



*Figure 1.3:* The evolution of the global 21 cm signal, starting with the Dark Ages, through galaxy formation and reionization (image credit: [Pritchard & Loeb \(2012\)](#)). The work in this thesis mainly focuses on a redshift range of  $6 < z < 12$  as reionization is expected to progress and complete.

photons, whereas a low  $T_{\text{spin}}$  implies that the gas is more likely to absorb 21 cm photons.

The spin temperature is always measured with respect to the temperature of the CMB, which serves as a backlight for our measurement. During different stages of our cosmic history,  $T_{\text{CMB}}$  and  $T_{\text{spin}}$  take turns in the spotlight, with the *differential brightness temperature*,  $\delta T_b$  describing their evolution:

$$\delta T_b \propto (1 + \delta_b) x_{\text{HI}} \left( 1 - \frac{T_{\text{CMB}}}{T_{\text{spin}}} \right). \quad (1.3)$$

Equation (1.3) captures the EoR signal that we would like to measure, where  $\delta_b$  is the fractional over-density of matter and  $x_{\text{HI}}$  is the fraction of neutral hydrogen (1 if all neutral, 0 if all ionized). The differential brightness temperature can be measured in multiple ways — in Chapter 1.2 we explain how interferometry (many telescopes) can be used to measure the correlations of  $\delta T_b$  on various spatial scales on the sky. Here we describe the evolution of the sky-averaged  $\delta T_b$ , called the *global signal*, in order to summarize how the signal is expected to behave during our *cosmic dawn* and through reionization.

A theoretical prediction for the evolution of  $\delta T_b$  is shown in Figure 1.3. At the very far left, a cool, neutral IGM remains after recombination and the release of the CMB. Residual electrons collide off of both CMB photons and the hydrogen gas, driving couplings between  $T_{\text{gas}}$  and  $T_{\text{CMB}}$ , and  $T_{\text{gas}}$  and  $T_{\text{spin}}$ , respectively. Hence, we expect to see no signal ( $\delta T_b = 0$ ) at this time.

During the Dark Ages, collisions still couple  $T_{\text{gas}}$  and  $T_{\text{spin}}$ , but Compton scattering becomes rarer as the CMB dilutes with the expansion of the Universe. While the CMB dilutes as  $T_{\text{CMB}} \propto 1/a$ , where  $a$  is the scale factor, the gas now follows an adiabatic expansion ( $T_{\text{gas}} \propto 1/a^2$ ). Thus, the gas cools quicker than the CMB, and since it is still coupled to the spin temperature,  $T_{\text{spin}} < T_{\text{CMB}}$  and the signal is expected to be seen in absorption. By the time the first galaxies begin forming, however, the gas is expected to be so dilute that it is no longer coupled to the spin temperature. The spin temperature therefore couples once again to the CMB, and no signal is produced.

As the first stars in the first galaxies begin emitting Lyman- $\alpha$  photons, the hydrogen gas is excited and de-excited, coupling  $T_{\text{spin}}$  and  $T_{\text{gas}}$ . The gas, still cool from adiabatic expansion, implies that  $T_{\text{spin}} < T_{\text{CMB}}$  and the signal is seen in absorption. Eventually, x-rays from the first sources begin heating the cooled gas, driving both the gas and spin temperatures above that of the CMB, where the signal is expected to be seen in emission for the first time.

Finally, even though the timing and details of reionization are unknown, UV photons from the first luminous structures are believed to eventually ionize all the neutral hydrogen, leaving no signal to be detected by a redshift of  $z \sim 6$ .

The shape of the global signal holds important science implications about our early Universe. For example, the position of the heating trough reveals the types of sources and energies of the x-rays doing the heating, as well as the sizes of the dark matter halos hosting those first sources (i.e. late heating and deep absorption troughs imply harder x-ray spectras, as shown in Fialkov et al. (2014)). The first tentative detection of this signature from the Experiment to Detect the Global Epoch of Reionization Signature (EDGES) even suggests



that known physics and commonly accepted scenarios may not be responsible for creating the absorption profile (Bowman et al. 2018). As the field continues to investigate our cosmic dawn through HI measurements — both the global signal and statistical fluctuations — we can expect to learn much about the constituents that make up the Universe and their complex interactions during this era.

### 1.1.4 This Thesis

Although 21 cm observations promises an uninterrupted window into the EoR, from which we can learn much about galaxy formation and the properties of the IGM, there are many challenges facing this field of cosmology. In general, the 21 cm signal is extremely faint, with bright foregrounds (mostly synchrotron radiation from our own Galaxy) and radio interference easily overshadowing the target signal. As a consequence, instruments need to be extremely well-understood, precisely calibrated, and capable of being sensitive enough for a successful detection. In addition, analysis techniques must be innovative and rigorously construed so as to be able to extract clean and accurate measurements.

In this thesis, I present work associated with data from radio interferometers seeking to measure 21 cm fluctuations during the EoR. While a confirmed detection by an interferometer remains elusive at this time, this work serves as a huge leap forward in working with large datasets and extracting measurements of the cosmological signal with confidence. The rest of this thesis thus focuses on the characterization of data from large telescope arrays in order to place accurate, stringent limits on the EoR signal. It is an exciting time and this field is young, and the work in this thesis serves as a foundation of what promises to be an eye-opening adventure to-come.

## 1.2 Interferometry

### 1.2.1 The Visibility Equation

### 1.2.2 Calibration

### 1.2.3 Foreground and Filtering

### 1.2.4 Fringe-rate Filtering

## 1.3 Instruments

### 1.3.1 The Precision Array for Probing the Epoch of Reionization

### 1.3.2 The Hydrogen Epoch of Reionization Array

### 1.3.3 Other Experiments



## Chapter 2

# Power Spectrum Methods

### 2.1 The 21 cm Power Spectrum

### 2.2 Signal Loss

### 2.3 Error Estimation

### 2.4 Bias

# Chapter 3

## PAPER-64

### 3.1 Overview

### 3.2 Signal Loss

### 3.3 Error Estimation

### 3.4 Bias

### 3.5 Power Spectrum Limits

# Chapter 4

## PAPER-128

### 4.1 Overview

### 4.2 Quality Assurance

#### 4.2.1 Flagging Julian Dates

#### 4.2.2 Flagging Antennas

## Chapter 5

### HERA

## Chapter 6

## Conclusion

# Bibliography

- Becker, R. H., Fan, X., White, R. L., et al. 2001, [AJ](#), **122**, 2850
- Bowman, J. D., Rogers, A. E. E., Monsalve, R. A., Mozdzen, T. J., & Mahesh, N. 2018, [Nature](#), **555**, 67
- Fialkov, A., Barkana, R., & Visbal, E. 2014, [Nature](#), **506**, 197
- Furlanetto, S. R., Oh, S. P., & Briggs, F. H. 2006, [Phys. Rep.](#), **433**, 181
- Haiman, Z., & Knox, L. 1999, in Astronomical Society of the Pacific Conference Series, Vol. 181, Microwave Foregrounds, ed. A. de Oliveira-Costa & M. Tegmark, 227
- Hinshaw, G., Larson, D., Komatsu, E., et al. 2013, [The Astrophysical Journal Supplement Series](#), **208**, 19
- Loeb, A., & Furlanetto, S. 2013, *The First Galaxies in the Universe* (Princeton University Press)
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, [A&A](#), **594**, A13
- Pritchard, J. R., & Loeb, A. 2012, [Reports on Progress in Physics](#), **75**, 086901
- Robertson, B. E., Ellis, R. S., Furlanetto, S. R., & Dunlop, J. S. 2015, [ApJ](#), **802**, L19
- Weisz, D. R., & Boylan-Kolchin, M. 2017, [MNRAS](#), **469**, L83