

Mixed and Longitudinal Models Group 11

final document

Group 11: Hannah Cooney, Carla Flore, Clara Domning

March 2026

```
load("Group_11.RData")
DF$sex <- factor(DF$sex, levels = c("male", "female"),
                labels = c("Male", "Female"))

DF$smoking_status <- factor(DF$smoking_status, levels = c(0, 1, 2),
                           labels = c("Never", "Former", "Current"))
```

Declaration

We declare that all students in the group have contributed sufficiently to the study and that the work is original and not copied from elsewhere.

Signed:

#Introduction (10-15 lines)

The aim of our analysis is to understand the evolution of motor symptom severity over time in patients diagnosed with Parkinson's Disease, and the influence of patient characteristics such as sex, age at diagnosis, and specific lifestyle factors on this evolution. In this study, Unified Parkinson's Disease Rating Scale (UPDRS) Part III is used to quantify motor symptom severity. This score ranges from 0 to 108, where the higher value indicates a more severe motor impairment.

500 diagnosed patients were enrolled in the study between 1995 and 2011. Assessments were conducted on patients at baseline, and then annually for 7 consecutive years. Several variables were recorded for each patient, including dates of each UPDRS assessment, the corresponding UPDRS Part III score at each visit, gender, and age of the patient when diagnosed. Their smoking history was documented at baseline. Our focus in this report is on the influence of smoking status on the disease trajectory.

A challenge often encountered in observational studies is the presence of missing data. This can be due to factors such as patient withdrawal, relocation, or inability to perform assessments as the disease progresses. We have also analysed the pattern of the missing data, and assessed whether the Missing At Random (MAR) assumption is reasonable in our case.

The structure of our report is as follows:

Section 2 presents a descriptive analysis of the data; Section 3 presents the statistical analysis, including the final model and analysis results; The Appendix details the procedure and the statistical techniques we have used when selecting our final model, and how we have dealt with missing data.

#Descriptive Analysis (1-2 pages)

##2.1 Sample Characteristics

###Age Distribution

##	n	Missing	Mean	SD	Median	Q1	Q3	IQR	Min	Max
## 1	500	0	64.6414	13.4897	65.1	52.575	76.325	23.75	41.1	88

The dataset has patients with baseline ages ranging from 41.1 years to 88 years, and no implausible values were observed. Overall, the age distribution reflects a typical Parkinson's disease cohort, with most patients diagnosed in their 50s to 70s and a moderate spread across older age groups.

###Sex Distribution

##	sex	n	prop
## 1	Male	250	0.5
## 2	Female	250	0.5

The dataset contains a balanced sample of male and female patients.

###Smoking Distribution

##	smoking_status	n	Percentage
## 1	Never	261	52.2
## 2	Former	141	28.2
## 3	Current	98	19.6

The majority of patients included in the dataset have never been smokers, slightly less than one third are former smokers, and approximately 20% are current smokers.


###Missing Data

At baseline, all 500 patients were observed. At subsequent visits, observation counts ranged from 303 to 342, reflecting intermittent missingness over the follow-up period.

2.2 Longitudinal Outcome

In order to explore the longitudinal behaviour of UPDRS scores, we examined the mean structure, the variance structure, and the correlation structure over time.

###Mean Structure:



The mean UPDRS Part III score increases steadily across follow-up visits, indicating a clear progression of motor symptom severity over time. The mean UPDRS score increases from 17.24 at baseline to 39.99 at Visit 7. The trajectory appears approximately linear, although the slope

increases slightly in later follow-up visits. This may indicate an slight acceleration of symptom progression overtime.

##Statistical Analysis (2-3 pages)

##Appendix (2-3 pages)