



UNIVERSITAT POLITÈCNICA DE CATALUNYA
BARCELONATECH

Facultat d'Informàtica de Barcelona



Curves 2A.13

Geometric Tools for Computer Graphics

November 13, 2024

Carles Matoses Gimenez *

*Email: carles.matoses@estudiantat@upc.edu

Contents

1	Exercise	2
2	Compute the distance to the origin from an arbitrary point of the curve. Which is the maximum value of these distances? Find the points of the curve achieving this maximum value.	2
3	Compute the tangent vector to the curve at an arbitrary point. Verify that the tangent vector at the point corresponding to $t = 0$ is linearly dependent from the tangent vector to the circle entered at the origin and of radius 5 at point $(5, 0)$.	3
4	Use change of reference systems to give a parametrization of a copy of the given curve placed in a disc of radius 5 centered at point $(-1, 3)$, so that both curves are tangent at point $(2, -1)$.	4
4.1	Change of basis for δ	4
5	Images	6

1 Exercise

Consider the curve parameterized by the function $\gamma : [0, \pi] \rightarrow \mathbb{R}^2$ defined as follows:

$$\gamma(t) = (x(t), y(t)) = (5 \cos(3t) \cos t, 5 \cos(3t) \sin t)$$

2 Compute the distance to the origin from an arbitrary point of the curve. Which is the maximum value of these distances? Find the points of the curve achieving this maximum value.

The distance $d(t)$ of an arbitrary point is given by:

$$d(t) = \sqrt{x(t)^2 + y(t)^2}$$

e.g.

$$d(2) = \sqrt{x(2)^2 + y(2)^2} = \sqrt{(5 \cos(2 \times 3) \cos(2))^2 + (5 \cos(2 \times 3) \sin(2))^2} = 4.79$$

Given the parametrization of the curve, the maximum distance is calculated as:

$$d(t)^2 = x(t)^2 + y(t)^2$$

$$d(t)^2 = (5 \cos(3t) \cos(t))^2 + (5 \cos(3t) \sin(t))^2$$

$$d(t)^2 = 25 \cos^2(3t) \cos^2(t) + 25 \cos^2(3t) \sin^2(t)$$

$$d(t)^2 = 25 \cos^2(3t)(\cos^2(t) + \sin^2(t))$$

With $\cos^2(t) + \sin^2(t) = 1$ and $25 \cos^2(3t) = 1$

$$d(t)^2 = 25 \cos^2(3t) = 25 \cdot 1 = 25$$

$$d(t) = \sqrt{25} = 5$$

The maximum points are achieved when $\cos(3t) = 1$:

$$3t = 0 \Rightarrow t = 0,$$

$$3t = \pi \Rightarrow t = \pi/3,$$

$$3t = 2\pi \Rightarrow t = 2\pi/3,$$

3 Compute the tangent vector to the curve at an arbitrary point. Verify that the tangent vector at the point corresponding to $t = 0$ is linearly dependent from the tangent vector to the circle entered at the origin and of radius 5 at point $(5, 0)$.

The tangent vector to the curve at an arbitrary point is calculated as the derivative $\gamma'(t)$.

$$\gamma(t) = \begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = \begin{pmatrix} 5 \cos(3t) \cos t \\ 5 \cos(3t) \sin t \end{pmatrix}$$

$$\gamma'(t) = \begin{pmatrix} x'(t) \\ y'(t) \end{pmatrix} = \begin{pmatrix} 5(-3 \sin(3t) \cos(t) + \cos(3t)(-\sin(t))) \\ 5(-3 \sin(3t) \sin(t) + \cos(3t) \cos(t)) \end{pmatrix}$$

Simplifying:

$$\gamma'(t) = \begin{pmatrix} x'(t) \\ y'(t) \end{pmatrix} = \begin{pmatrix} -15 \sin(3t) \cos(t) - 5 \cos(3t) \sin(t) \\ -15 \sin(3t) \sin(t) + 5 \cos(3t) \cos(t) \end{pmatrix}$$

For $t = 0$ in the curve we get:

$$\gamma'(0) = \begin{pmatrix} x'(0) \\ y'(0) \end{pmatrix} = \begin{pmatrix} -15 \sin(0) \cos(0) - 5 \cos(0) \sin(0) \\ -15 \sin(0) \sin(0) + 5 \cos(0) \cos(0) \end{pmatrix} = \begin{pmatrix} 0 \\ 5 \end{pmatrix}$$

For $t = 0$ in the tangent of a circle with *radius* = 5:

$$C(t) = \begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = \begin{pmatrix} R \cos(t) \\ R \sin(t) \end{pmatrix}$$

$$C'(0) = \begin{pmatrix} x'(0) \\ y'(0) \end{pmatrix} = \begin{pmatrix} 5(-\sin(0)) \\ 5(\cos(0)) \end{pmatrix} = \begin{pmatrix} 0 \\ 5 \end{pmatrix}$$

$\gamma'(0)$ and $C'(0)$ are linearly dependent since they are scalars of each other. More specifically, they are the same vector.

4 Use change of reference systems to give a parametrization of a copy of the given curve placed in a disc of radius 5 centered at point $(-1, 3)$, so that both curves are tangent at point $(2, -1)$.

The new curve δ will intersect the circle in the point:

$$p = (2, -1)$$

4.1 Change of basis for δ

$$s = \begin{cases} e_1 = (1, 0, 0) \\ e_2 = (0, 1, 0) \\ e_3 = (0, 0, 1) \end{cases}$$

$$circle_o = (-1, 3)$$

To generate the new basis we follow this steps:

1. Get the vector that goes from the center of the circle to the curve point p .

$$\delta_v = p - circle_o = (2, -1) - (-1, 3) = (3, -4)$$

2. Generate a tangent vector using δ_v and the perpendicular vector method $(-b, a)$.

$$\delta_t = (4, 3)$$

3. We now create the new vectors e'_1, e'_2, e'_3 using δ_t and creating the tangent vector:

$$s' = \begin{cases} e'_1 = \frac{1}{5}(3, -4, 0) \\ e'_2 = \frac{1}{5}(4, 3, 0) \\ e'_3 = (0, 0, 1) \end{cases}$$

4. Finally, we create the homogeneous matrix:

$$T = \begin{pmatrix} \frac{3}{5} & \frac{4}{5} & 0 & -1 \\ \frac{-4}{5} & \frac{3}{5} & 0 & 3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\delta(t) = T \cdot \gamma(t) = \begin{pmatrix} 3 \cos(3t) \cos t + 4 \cos(3t) \sin t - 1 \\ -4 \cos(3t) \cos t + 3 \cos(3t) \sin t + 3 \\ 0 \\ 1 \end{pmatrix}$$

To check the result, the easiest way is to multiply the matrix R by the known tangent vectors at point $(5,0)$.

$$R = \begin{pmatrix} \frac{3}{5} & \frac{4}{5} & 0 \\ \frac{-4}{5} & \frac{3}{5} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

The circle has a tangent vector of $(0,5)$ as well as the γ curve at $t = 0$. This means that rotating them by the same quantity will preserve the tangential property.

$$R \cdot (0, 5, 0) = (4, 3, 0)$$

The new tangent will be $(4, 3, 0)$ for both curves. Other solutions would require the derivation the $\delta(t)$ parametrization and getting the *sin* and *cos* of the circle at $(2,-1)$.

5 Images

To improve readability, I decided to place explanatory images after the operations to help on the visualization on some parts of the problem. The object we are working with looks like this. The parametrization presents a faster frequency on one component (cosine) than the other (sine) which makes the function to generate this smaller loops. The radius in front of the cosine will represent the amplitude of the final curve.




Figure 1: Base object




Figure 2: Tangent $\pi/3$

Changing the frequency will, of course increment or decrees the number of loops.




Figure 3: 1 loop




Figure 4: 5 loops




Figure 5: 11 loops

The following image 6 shows visually how the vector from the circle and the curve are the same:




Figure 6: Tangent vectors are equal.

Visualizing the tangent vectors 7 after the transformation allows us to observe how both tangents are 90° .




Figure 7: Tangent vectors

This link redirects to github where there is an intractable visualizer of the exercise done with blender. [https : //github.com/carlesmatoses/GTCG_2A_13](https://github.com/carlesmatoses/GTCG_2A_13)