

Guaranteed overlap between a 2-factor and a Hamilton cycle

Carlos Andrés Velásquez Carrasco

Madrid, Spain

`carlitos.andres.v1@gmail.com`

October 2026

Abstract

Let G be a Hamiltonian graph on n vertices and let F be a 2-factor of G . How many edges of F can one guarantee that some Hamilton cycle of G shares with it? Write $\text{ov}(n)$ for the largest integer s such that every pair (G, F) with $|V(G)| = n$ satisfies $\max_H |F \cap H| \geq s$, where H ranges over the Hamilton cycles of G . We prove the following. First, a reduction theorem: the extremal problem is concentrated on the graphs that are the union of a 2-factor and a Hamilton cycle, hence have at most $2n$ edges and maximum degree 4. This makes exact computation feasible, and we obtain $\text{ov}(n) = 4$ for $6 \leq n \leq 8$ and $\text{ov}(n) = 5$ for $9 \leq n \leq 13$, in the last case by traversing all 438 263 364 labelled 2-regular graphs on 13 vertices. Second, we construct two explicit infinite families, Γ_n and Δ_n , defined for every $n \geq 10$ by a zigzag over an interval, prove that both are uniquely Hamiltonian, and deduce $\text{ov}(n) \leq 5$ for every $n \geq 10$; for $n = 10$ and $n = 11$ these are, up to the dihedral symmetry, the only uniquely Hamiltonian extremal pairs, while for $n = 12$ there is one more and for $n = 13$ two more. All of them, at every size, have two vertices of degree 2 and six of degree 3, so that these graphs are uniquely Hamiltonian with maximum degree 4 and only eight vertices of smaller degree, uniformly in n ; and all of them share a finer invariant, namely that $F \cap H$ splits into exactly three paths, of one, two and two edges, which by a degree count is what fixes the profile. Third, we show that the reverse inequality is not elementary: even the assertion $\text{ov}(n) \geq 1$ implies Sheehan's 1975 conjecture on second Hamilton cycles in 4-regular graphs. We also prove a protected-edge lemma: if H attains the maximum, then any competing Hamilton cycle can only give up edges of $F \cap H$, so that in an optimal pair the whole Hamiltonian structure is confined to those $|F \cap H|$ edges and the number of Hamilton cycles is bounded by a function of $|F \cap H|$ alone, independently of n . Finally, we reduce $\text{ov}(n) \geq 5$ to the configurations sharing at most four edges with the cycle and, by means of an amplification lemma and a contraction producing a 4-regular multigraph with forbidden transitions, we place the conjecture $\text{ov}(n) = 5$ between Sheehan's conjecture and a strengthening of it localised at four vertices.

Keywords. 2-factor; Hamilton cycle; uniquely Hamiltonian graph; Sheehan's conjecture; extremal function; exhaustive enumeration.

MSC 2020. Primary 05C45; Secondary 05C70, 05C85.

1 Introduction

A *2-factor* of a graph G is a spanning 2-regular subgraph, that is, a partition of $V(G)$ into cycles; a Hamilton cycle is the case of a single cycle. There is therefore a natural distance between an arbitrary 2-factor and a Hamilton cycle, and the question we study is how much of the structure of the 2-factor can be preserved when passing from one to the other.

Definition 1.1 (Overlap). Let G be a Hamiltonian graph on n vertices and let F be a 2-factor of G . Set

$$\text{ov}_G(F) = \max\{|F \cap H| : H \text{ a Hamilton cycle of } G\}, \quad \text{cor}_G(F) = n - \text{ov}_G(F),$$

so that $\text{cor}_G(F)$ is the least number of edges one must discard from F in order for the remainder to be contained in a Hamilton cycle of G . The *extremal function* is

$$\text{ov}(n) = \min\{\text{ov}_G(F) : G \text{ Hamiltonian with } |V(G)| = n, F \text{ a 2-factor of } G\}.$$

By definition $\text{ov}(n)$ is the largest integer s such that *every* 2-factor of *every* Hamiltonian graph on n vertices shares at least s edges with some Hamilton cycle. At first sight it is not even clear that $\text{ov}(n) \geq 1$; Theorem 6.1 explains why.

Results. Theorem 3.1 reduces the computation of $\text{ov}(n)$ to the pairs (F, H) consisting of a 2-factor and a Hamilton cycle on the vertex set $[n]$, with $G = F \cup H$. This confines the problem to graphs with at most $2n$ edges and maximum degree 4, and makes possible the exhaustive computation recorded in Table 1:

$$\text{ov}(6) = \text{ov}(7) = \text{ov}(8) = 4, \quad \text{ov}(9) = \dots = \text{ov}(13) = 5.$$

Theorems 5.3 and 5.5 construct, for every $n \geq 10$, two uniquely Hamiltonian graphs that are unions of a 2-factor and a Hamilton cycle sharing exactly five edges, whence $\text{ov}(n) \leq 5$ for every $n \geq 10$. Theorem 6.1 shows that $\text{ov}(n) \geq 1$ implies Sheehan's conjecture. Lemma 5.7 then shows that the degree profile of such a pair is determined by the number of components of $F \cap H$, which turns the classification into seven separate enumerations, one per composition of 5; carrying them out for $10 \leq n \leq 15$ yields Conjecture 5.8, a structural refinement asserting that $F \cap H$ always splits into three stretches of sizes 1, 2, 2. Finally, Proposition 7.1 reduces the inequality $\text{ov}(n) \geq 5$ to the configurations with $|F \cap H| \leq 4$, and Corollary 7.14 places it between Sheehan's conjecture and a strengthening of it with forbidden transitions at at most four vertices. We conjecture that $\text{ov}(n) = 5$ for every $n \geq 9$.

Related work. The relationship between 2-factors and Hamilton cycles has been studied mostly in the direction opposite to ours: given a graph with a Hamilton cycle, does it have a 2-factor with exactly k cycles? Bucić, Jahn, Pokrovskiy and Sudakov [4] obtained degree conditions for this, and Espuny Díaz, Girão, Granet and Kronenberg [5] lowered them to $\delta(G) \geq n^{1-\varepsilon}$ by means of exchanges along quadrilaterals. Those works *split* a cycle whereas our problem *merges* cycles; moreover, the set of edges they remove from a 2-factor is not minimised, while here the minimisation is the object of study. On the reconfiguration side, Klee, Patel and Stroh [15] introduced *k-switches* on families of spanning subgraphs, with the normalisation $|E(F) \triangle E(F')| \leq 2k$ that we also use, and proved that if $\delta(G) \geq \frac{1}{2}n + 1$ then every 2-factor with t components reaches a Hamilton cycle in at most $t - 1$ switches of size at most 3; Kang and Legrand-Duchesne [14] study the analogous reconfiguration graph on perfect matchings. Our parameter differs from all of these: it measures the overlap achievable in a *single* jump rather than the length of a sequence of exchanges, and Remark 8.4 shows that the two points of view separate sharply.

A third body of work is closer in spirit, and the connection deserves to be made explicit. By Lemma 2.1, the inequality $\text{ov}_G(F) \geq s$ says precisely that *some* subset of s edges of F extends to a Hamilton cycle of G ; our question is therefore one about extending linear forests, with the twist that the forest is not prescribed but chosen inside F . The extension question with a prescribed forest has been studied extensively, in particular for matchings in hypercubes, where one asks for the largest k such that *every* matching of size k extends

to a Hamilton cycle [9], and for perfect matchings of subcubes [8]. The two questions are complementary: there the object to be extended is given and one quantifies over all of them, here one is free to choose it inside a prescribed 2-factor. We are not aware of a deficiency version of the extension question, that is, of a quantitative measure of how much of a 2-factor can be preserved. For the general landscape of 2-factors of regular graphs we refer to the survey [16].

2 Preliminaries

All graphs are finite and simple, and $[n] = \{0, 1, \dots, n-1\}$. If H is a fixed Hamilton cycle on $[n]$ we call *chord* any edge not belonging to H . When working with a pair (F, H) we write throughout

$$\Gamma = F \cup H, \quad S = F \cap H, \quad s = |S|.$$

Note that in Γ every edge not in H belongs to F , and that $|E(\Gamma)| = 2n - s$.

The first lemma replaces the search for Hamilton cycles by a local check, and it is what makes the computations of Section 4 possible.

Lemma 2.1 (Assembly criterion). *Let F be a 2-factor of G and let $j \geq 1$. The following are equivalent:*

- (a) $\text{cor}_G(F) \leq j$;
- (b) *there is $R \subseteq F$ with $|R| \leq j$ such that $F \setminus R$ is a spanning linear forest whose paths admit a cyclic ordering and orientations with consecutive endpoints adjacent in G .*

Proof. (b) $\Rightarrow$ (a) If the paths of $F \setminus R$ can be assembled cyclically using edges of G , the result is a Hamilton cycle H of G containing $F \setminus R$, so $|F \setminus H| \leq |R| \leq j$.

(a) $\Rightarrow$ (b) Let H be Hamiltonian with $|F \setminus H| \leq j$ and put $R = F \setminus H$. Then $F \setminus R = F \cap H$ is a subgraph of H with all degrees at most 2; if it contained a cycle, that cycle would lie inside H and hence equal H , so $F = H$. In that case it suffices to take $R = \{e\}$ for any $e \in F$, which is legitimate since $j \geq 1$, and $F \setminus R$ is a Hamilton path closed up by e . Outside that case $F \cap H$ is a spanning linear forest and H itself exhibits the cyclic ordering and the orientations of its paths. $\square$

The cost of deciding $\text{cor}_G(F) \leq j$ is therefore $\sum_{i \leq j} \binom{n}{i}$ tests, each of which runs through at most $(j-1)! 2^{j-1}$ cyclic orderings with orientations; in practice it is a search with branching factor bounded by $\Delta(G)$, which for $\Delta(G) \leq 4$ prunes almost immediately.

3 Reduction to unions of two 2-factors

Theorem 3.1 (Reduction). *For every $n \geq 3$,*

$$\text{ov}(n) = \min \{ \text{ov}_{F \cup H}(F) : F \text{ a 2-factor of } [n], H \text{ a Hamilton cycle of } [n] \},$$

and in the second minimum one may fix $H = (0, 1, \dots, n-1)$.

Proof. ($\leq$) For any F and H the graph $F \cup H$ is Hamiltonian, since it contains H , and F is one of its 2-factors; hence $\text{ov}_{F \cup H}(F)$ is one of the values over which the minimum defining $\text{ov}(n)$ is taken.

($\geq$) Let G be Hamiltonian and let F be a 2-factor of G with $\text{ov}_G(F) = \text{ov}(n)$. Choose any Hamilton cycle H of G and put $G' = F \cup H$. Then $G' \subseteq G$ is Hamiltonian and F is a 2-factor of G' . Every Hamilton cycle of G' is one of G , so the maximum defining $\text{ov}_{G'}(F)$

ranges over a subfamily of the one defining $\text{ov}_G(F)$, whence $\text{ov}_{G'}(F) \leq \text{ov}_G(F) = \text{ov}(n)$. By minimality there is equality. Finally, after relabelling the vertices one may assume $H = (0, 1, \dots, n-1)$. $\square$

The content of the theorem is that the extremal problem lives entirely among sparse graphs: every union $F \cup H$ has at most $2n$ edges and maximum degree 4. Adding edges can only increase the overlap, since it adds candidate Hamilton cycles; the worst case is always the minimal skeleton. This turns the exhaustive computation into a feasible task: instead of running through the Hamiltonian graphs on n vertices it suffices to run through the labelled 2-regular graphs on $[n]$.

Remark 3.2 (Relation to the classical reduction for Sheehan’s conjecture). The argument is the same one that reduces Sheehan’s conjecture from the $2k$ -regular case to the 4-regular case by Petersen’s 2-factorisation theorem: pass to a sparse spanning subgraph containing the given Hamilton cycle, using the fact that passing to a subgraph can only make the sought object harder to find. The difference is that the classical reduction stops at “there is a second cycle”, whereas here the quantity $|F \cap H'|$ is preserved, and it is that quantity which turns an existence problem into an extremal problem with a computable function.

4 The extremal function up to $n = 13$

Combining Theorem 3.1 with Lemma 2.1 we have traversed all 2-factors of $[n]$ for $n \leq 13$.

n	5	6	7	8	9	10	11	12	13
2-factors of $[n]$	12	70	465	3 507	30 016	286 884	3 026 655	34 944 085	438 263 364
$\text{ov}(n)$	5	4	4	4	5	5	5	5	5
$n - \text{ov}(n)$	0	2	3	4	4	5	6	7	8

Table 1: The extremal function. The row of counts reproduces the sequence of labelled 2-regular graphs, which validates the generator. The case $n = 5$ is degenerate: every 2-factor of $[5]$ is already a Hamilton cycle, so $\text{ov}(5) = 5$.

The computations were carried out in C, with a recursive generator of the 2-factors of $[n]$ and the test of Lemma 2.1; the test was validated against an independent computation of the maximum overlap by dynamic programming over subsets, of cost $O(2^n n^2)$, on 1 862 random pairs with $7 \leq n \leq 11$, on every extremal pair of the exhaustive range, and on every pair returned by the directed searches up to $n = 16$. The values for $n \leq 11$ and the $n = 13$ figures were independently reproduced by a second implementation written from scratch, and the classification of the uniquely Hamiltonian extremal pairs for $10 \leq n \leq 13$ by a third, the pattern-directed search of Remark 5.11, whose column sums provide a further internal check.

Remark 4.1 (The growth is linear, not proportional). The table shows that $n - \text{ov}(n)$ equals $n - 4$ for $n = 6, 7, 8$ and $n - 5$ from $n = 9$ on: it grows linearly with slope 1, and not, for instance, as a fraction of n . In other words, a 2-factor may be forced to sacrifice all but a *constant* number of its edges. Section 5 shows that this constant is at most 5 for every $n \geq 10$.

It is also worth recording the behaviour in the disjoint case $S = \emptyset$, which is exactly that of 4-regular graphs and the one corresponding to Sheehan’s conjecture.

5 Two families attaining $\text{ov}(n) \leq 5$

For each n the exhaustive traversal produces several pairwise non-isomorphic extremal pairs. For $n = 10$ there are 160 pairs with $\text{ov}_\Gamma(F) = 5$, distributed over $s = 3$, $s = 4$ and $s = 5$;

n	8	9	10	11	12	13
pairs with $F \cap H = \emptyset$	248	2 316	23 687	263 652	3 178 529	41 309 910
minimum overlap	6	6	6	7	7	8
$n - \lfloor n/2 \rfloor + 1$	5	6	6	7	7	8

Table 2: The 4-regular case. Over this range the guaranteed overlap grows with n — an *empirical* fact on six values; proving that it is unbounded would be far stronger than Sheehan’s conjecture — so the family realising the global minimum of Table 1 is not 4-regular. The last row is the expression $n - \lfloor n/2 \rfloor + 1$, which fits the data except at $n = 8$.

of these, 40 are uniquely Hamiltonian and form exactly *two* classes under the dihedral group preserving H , both with $(n_2, n_3) = (2, 6)$. For $n = 11$ there are 121 extremal pairs, all with $s = 5$, of which 44 are uniquely Hamiltonian and again form exactly two classes. For $n = 12$ there are 168 extremal pairs, of which 72 are uniquely Hamiltonian, and these form *three* classes; for $n = 13$ there are 104 uniquely Hamiltonian extremal pairs in *four* classes. The first two classes at each size extend to infinite families: they are the members $n = 10, \dots, 13$ of the families Γ_n and Δ_n described below; the extra classes are discussed in Remark 5.6. All eleven classes have the degree profile $(n_2, n_3, n_4) = (2, 6, n - 8)$.

For these counts we used two independent programs. The first computes, for every labelled 2-factor, the maximum overlap and the number of Hamilton cycles of Γ , and canonicalises the pairs that are extremal and uniquely Hamiltonian. The second exploits an observation that makes the search much cheaper: if Γ is uniquely Hamiltonian then its only Hamilton cycle is H , so the maximum overlap equals s exactly, and a uniquely Hamiltonian extremal pair must have $s = 5$; it therefore generates only the 2-factors with exactly five shared edges, pruning on that count during generation, and tests unique Hamiltonicity with an early exit at the second cycle. The two programs agree on the numbers above.

Definition 5.1 (The family Γ_n). Let $n \geq 10$, $V = [n]$ and $H = (0, 1, \dots, n - 1, 0)$. Over the interval $A = [7, n - 1]$ define the *zigzag* $z_0, z_1, \dots, z_{L-1}$, with $L = n - 7$, by

$$z_{2i} = 7 + i, \quad z_{2i+1} = n - 1 - i,$$

that is, the sequence $7, n - 1, 8, n - 2, 9, n - 3, \dots$ which runs through A alternating between its two ends towards the centre. Let F_n be the 2-factor consisting of the triangle $\{12, 23, 13\}$ together with the cycle $(0, z_0, z_1, \dots, z_{L-1}, 4, 5, 6)$, and let $\Gamma_n = F_n \cup H$.

The chords of the zigzag admit a closed description: the edges $z_i z_{i+1}$ are exactly the pairs ab with $7 \leq a < b \leq n - 1$ and $a + b \in \{n + 6, n + 7\}$, that is, the two reflections $a \mapsto (n + 6) - a$ and $a \mapsto (n + 7) - a$ restricted to A .

Lemma 5.2. *For every $n \geq 10$: F_n is a 2-factor with two cycles, $|F_n \cap H| = 5$ and*

$$F_n \cap H = \{12, 23, 45, 56, \mu\}, \quad \mu = z_{L-2} z_{L-1},$$

where μ is the unique pair of consecutive integers of the zigzag. The graph Γ_n has exactly two vertices of degree 2, namely 2 and 5, six of degree 3 and $n - 8$ of degree 4.

Proof. The long cycle runs through $\{0\} \cup A \cup \{4, 5, 6\} = \{0\} \cup \{4, 5, \dots, n - 1\}$, which together with the triangle covers V ; both cycles have length at least 3 because $n \geq 10$. Of its edges, those belonging to H are 45 and 56 together with those falling inside A : the edges $0z_0 = 07$, $z_{L-1}4$ and 06 do not, since $n \geq 10$. Inside A , an edge $z_i z_{i+1}$ lies in H if and only if its endpoints differ by 1; since $z_i + z_{i+1} \in \{n + 6, n + 7\}$ and a pair of consecutive integers has

odd sum, this happens for at most one of the two values, and it happens exactly once because one of $n + 6$ and $n + 7$ is odd and the corresponding pair falls inside A when $n \geq 10$. Adding the two edges of the triangle that lie in H gives $|F_n \cap H| = 5$.

As for the degrees, in a union $F \cup H$ the degree of v is 4 minus the number of edges of F incident with v that lie in H ; hence, if n_i denotes the number of vertices of degree i ,

$$|F \cap H| = \frac{1}{2} (n_3 + 2n_2). \quad (1)$$

Here the five edges of $F_n \cap H$ form the two paths 1–2–3 and 4–5–6 together with the isolated edge μ , so $n_2 = 2$, the interior vertices being 2 and 5, and $n_3 = 6$, in accordance with (1). $\square$

Theorem 5.3 (Γ_n is uniquely Hamiltonian). *For every $n \geq 10$ the only Hamilton cycle of Γ_n is H . Consequently $\text{ov}_{\Gamma_n}(F_n) = 5$ and*

$$\text{ov}(n) \leq 5 \quad \text{for every } n \geq 10.$$

Proof. Let H' be a Hamilton cycle of Γ_n . We use three elementary rules: (R1) if $\deg(v) = 2$, both edges at v lie in H' ; (R2) if two edges incident with v lie in H' , no other does; (R3) if u and v are the endpoints of a path contained in H' with fewer than $n - 1$ edges, then $uv \notin H'$, since it would close a proper cycle.

Start. By Lemma 5.2, $\deg(2) = \deg(5) = 2$, so $12, 23, 45, 56 \in H'$ by (R1). The path 1–2–3 and (R3) give $13 \notin H'$; since the edges at 1 are $\{01, 12, 13\}$ and those at 3 are $\{23, 34, 13\}$, it follows that $01, 34 \in H'$. This establishes the path $P_0 : 0 - 1 - 2 - 3 - 4 - 5 - 6$.

The second seed. The two edges of H' at the vertex 4 are already 34 and 45, so by (R2) the chord $4q \notin H'$, where $q = z_{L-1}$. But the edges at q in Γ_n are exactly its two H -edges together with the chord $4q$: indeed, the two edges of F_n at q are $\mu = z_{L-2}z_{L-1}$, which is an edge of H , and $4q$. Hence q has only two available edges and both lie in H' :

$$(q - 1)q \in H' \quad \text{and} \quad q(q + 1) \in H'. \quad (2)$$

Priming. By (R3) applied to P_0 , which has six edges and $6 < n - 1$, we get $06 \notin H'$; since the edges at 6 are $\{56, 67, 06\}$, it follows that $67 \in H'$. Again by (R3), $07 \notin H'$; the vertex 0 has edges $\{01, 0(n-1), 06, 07\}$ with $01 \in H'$ and $06, 07 \notin H'$, so $0(n-1) \in H'$. This leaves the arc $P_1 : (n - 1) - 0 - 1 - \dots - 7$, with 8 edges.

The zipper. Put $i_{\max} = \lfloor (n - 11)/2 \rfloor$. We claim, by induction on $0 \leq i \leq i_{\max} + 1$, that H' contains the arc of H

$$Q_i : (n - 1 - i) - \dots - (n - 1) - 0 - 1 - \dots - (7 + i),$$

which has $2i + 8$ edges and $2i + 9$ vertices. The case $i = 0$ is P_1 . Assume $Q_i \subseteq H'$ with $i \leq i_{\max}$; then $2i + 9 \leq n - 2$, so (R3) applies both to Q_i and to the arc with one more edge obtained in the first step. Observe moreover that $7 + i \leq 7 + i_{\max} < q = \lfloor (n + 7)/2 \rfloor$ and that $8 + i < q$, so neither the leading end nor the vertex added to it is the meeting point of the zigzag; in particular the edges of F_n involved below are chords and not the shared edge μ .

The two edges of F_n at the leading end $a = 7 + i$ are the zigzag chords ab with $a + b \in \{n + 6, n + 7\}$, that is $b = n - 1 - i$ and $b = n - i$; for $i = 0$ the second does not exist and is replaced by the chord 07 . In all cases the other endpoint belongs to Q_i , so (R3) excludes them and, since $(6 + i)(7 + i) \in H'$, the edge $(7 + i)(8 + i) \in H'$ is forced. The two edges of F_n at the trailing end $c = n - 1 - i$ are the chords with $c + b \in \{n + 6, n + 7\}$, that is $b = 7 + i$ and $b = 8 + i$; both vertices already lie in the enlarged arc, which has $2i + 9 \leq n - 2$ edges, so (R3) excludes them and $(n - 2 - i)(n - 1 - i) \in H'$ is forced. This gives $Q_{i+1} \subseteq H'$.

Closure. The induction ends at Q_{i^*} with $i^* = i_{\max} + 1 = \lfloor (n - 9)/2 \rfloor$, which has $2i^* + 9$ vertices. If n is odd then $i^* = (n - 9)/2$ and Q_{i^*} covers all n vertices; if n is even then

$i^* = (n - 10)/2$ and it covers $n - 1$ of them, leaving out a single vertex, namely the meeting point $q = \lfloor (n+7)/2 \rfloor$ of the zigzag. In the first case Q_{i^*} is a Hamilton path lacking exactly one edge of H to close up, and its endpoints are precisely the endpoints of that edge, so $H' = H$. In the second case Q_{i^*} is a Hamilton path of $\Gamma_n - q$ with endpoints $q - 1$ and $q + 1$, and (2) supplies the two missing edges, again giving $H' = H$. Since H is the unique Hamilton cycle, $\text{ov}_{\Gamma_n}(F_n) = |F_n \cap H| = 5$. $\square$

5.1 The second family

The other uniquely Hamiltonian class at sizes 10 and 11 also extends to every n , by the same idea but with two triangles instead of one.

Definition 5.4 (The family Δ_n). Let $n \geq 10$ and $H = (0, 1, \dots, n - 1, 0)$. Let $z_0, \dots, z_{L-1}$, with $L = n - 8$, be the zigzag over the interval $[8, n - 1]$, that is $z_{2i} = 8 + i$ and $z_{2i+1} = n - 1 - i$. Let F'_n be the 2-factor consisting of the triangles $\{12, 23, 13\}$ and $\{56, 67, 57\}$ together with the cycle $(0, z_0, z_1, \dots, z_{L-1}, 4)$, and let $\Delta_n = F'_n \cup H$.

Theorem 5.5. For every $n \geq 10$, F'_n is a 2-factor with three cycles, $|F'_n \cap H| = 5$ with

$$F'_n \cap H = \{12, 23, 56, 67, \mu'\}, \quad \mu' = z_{L-2}z_{L-1},$$

and Δ_n is uniquely Hamiltonian, whence $\text{ov}_{\Delta_n}(F'_n) = 5$. Its vertices of degree 2 are 2 and 6, and its degree profile is $(n_2, n_3, n_4) = (2, 6, n - 8)$.

Proof. The computation of $F'_n \cap H$ is that of Lemma 5.2: the edges $0z_0 = 08$, $z_{L-1}4$ and 04 do not lie in H because $n \geq 10$, and inside $[8, n - 1]$ there is exactly one pair of consecutive integers because $z_i + z_{i+1} \in \{n + 7, n + 8\}$ and one of those two values is odd. The remaining four edges of $F'_n \cap H$ are the two from each triangle.

For uniqueness one repeats the proof of Theorem 5.3 with three changes. The start now has two degree-2 seeds, the vertices 2 and 6, which force $12, 23, 56, 67 \in H'$; the two triangles give, by (R3), $13, 57 \notin H'$ and hence $01, 34, 45, 78 \in H'$, so that the initial forced path is $P_0 : 0 - 1 - 2 - \dots - 8$, of eight edges instead of six. Secondly, the role of the “second seed” is played by the vertex 4: its two H -edges are already forced, so by (R2) its two chords 04 and $4z_{L-1}$ are excluded, and then the vertex z_{L-1} is left with only two available edges, its two H -edges, which become forced. Finally, the priming and the zipper are identical upon replacing 7 by 8 and the values $\{n + 6, n + 7\}$ by $\{n + 7, n + 8\}$: by (R3) applied to P_0 the chord 08 is excluded and $0(n - 1)$ is forced, and from there the arc grows alternately at its two ends because the two chords at the leading end $8 + i$ go to $n - 1 - i$ and to $n - i$, and those at the trailing end $n - 1 - i$ go to $8 + i$ and to $9 + i$, all of them inside the arc already built. $\square$

We have verified Theorem 5.5 by computer for every $10 \leq n \leq 30$: $|F'_n \cap H| = 5$, exactly one Hamilton cycle, degree profile $(2, 6, n - 8)$, and the forcing propagation closes on all n edges of H . Theorem 5.3 was likewise verified in two ways independent of the proof: by counting the Hamilton cycles of Γ_n by backtracking for $10 \leq n \leq 33$, always exactly one, and by computing $\max_{H'} |F_n \cap H'|$ by dynamic programming for $n \leq 16$, always exactly 5.

Remark 5.6 (The bound does not characterise a single family). The two families share the degree profile $(2, 6, n - 8)$, the value $|F \cap H| = 5$ and the mechanism — two vertices of degree 2 seeding a forcing chain along a zigzag — but they differ in the number of cycles of the 2-factor: two in Γ_n and three in Δ_n . The bound $\text{ov}(n) \leq 5$ therefore does not characterise a construction: at least two essentially different families attain it, and at sizes 10 and 11 they are exactly these two. At $n = 12$ they are not: there are 72 uniquely Hamiltonian extremal pairs falling into *three* dihedral classes, namely Γ_{12} , Δ_{12} and

$$F = \{12, 23, 13\} \cup \{56, 67, 57\} \cup (0, 8, 10, 11, 4, 9), \quad S = \{(10, 11), 12, 23, 56, 67\},$$

which has the same cycle type $3 + 3 + 6$ as Δ_{12} but is not equivalent to it: it is a second zigzag of the same shape. At $n = 13$ there are four classes: Γ_{13} , of cycle type $3 + 10$, and three of cycle type $3 + 3 + 7$, of which one is Δ_{13} . So the count of classes runs $2, 2, 3, 4$ for $n = 10, \dots, 13$ and we expect it to keep growing; what the families capture is a mechanism, not a list.

What the eleven classes do share is more informative than what separates them: every one of them has the degree profile $(n_2, n_3, n_4) = (2, 6, n - 8)$. That is not an accident of the examples but a statement about the shape of S , as the next lemma makes precise.

5.2 The stretch pattern

Lemma 5.7 (The profile is the number of stretches). *Let (F, H) be a pair with $|F \cap H| = s \geq 1$ and let c be the number of connected components of $S = F \cap H$. Then*

$$n_2 = s - c, \quad n_3 = 2c, \quad n_4 = n - s - c.$$

Proof. The components of S are paths, the stretches, say with $k_1, \dots, k_c$ edges and $\sum_i k_i = s$. A vertex of Γ has degree 2 exactly when both its F -edges lie in H , that is when it is an interior vertex of a stretch, and degree 3 exactly when one of them does, that is when it is an endpoint of a stretch. A stretch with k_i edges has $k_i - 1$ interior vertices and two endpoints, whence $n_2 = \sum_i (k_i - 1) = s - c$ and $n_3 = 2c$. Consistently $n_3 + 2n_2 = 2s$, which is (1). $\square$

For $s = 5$ the profile is therefore determined by $c \in \{1, \dots, 5\}$ alone, the five values giving $(n_2, n_3) = (4, 2), (3, 4), (2, 6), (1, 8), (0, 10)$; the observed one is $c = 3$. The finer datum, the multiset $\{k_1, \dots, k_c\}$, is also constant across the eleven classes: it is always $\{1, 2, 2\}$, never the other composition $\{1, 1, 3\}$ of 5 into three parts. We therefore state:

Conjecture 5.8. If (F, H) is uniquely Hamiltonian with $|F \cap H| = 5$, then S has exactly three components, of sizes 1, 2 and 2. In particular $(n_2, n_3, n_4) = (2, 6, n - 8)$.

The reformulation is worth making because an elementary operation turns each value of c into a question about graphs of maximum degree 4 — though, as Remark 5.10 will show, about *marked* such graphs, which is what keeps the five cases from being the five known questions one would hope for.

Proposition 5.9 (Suppression). *Let (F, H) be a pair with s shared edges in c stretches, and let Γ^b be the multigraph obtained from Γ by suppressing its $s - c$ vertices of degree 2, that is by replacing each stretch by a single edge, which we call marked, joining its two endpoints. Then Γ^b has $n - s + c$ vertices, exactly $2c$ of degree 3 and all the others of degree 4, and*

$$\begin{aligned} & \{\text{Hamilton cycles of } \Gamma \text{ containing } S\} \\ & \longleftrightarrow \{\text{Hamilton cycles of } \Gamma^b \text{ through all } c \text{ marked edges}\}. \end{aligned}$$

In particular, if Γ is uniquely Hamiltonian then Γ^b has exactly one Hamilton cycle through its marked edges. The converse fails: a stretch of a single edge has no interior vertex of degree 2 and therefore forces nothing, so Γ may well have Hamilton cycles omitting part of S , which do not descend. At $n = 8$, for $F = \{01, 02, 12, 34, 36, 47, 56, 57\}$ and $S = \{01, 12, 34, 56\}$ the graph Γ has two Hamilton cycles, the second of which drops the shared edge 34, while Γ^b has exactly one through its marked edges. Moreover Γ^b is simple if and only if no stretch has its two endpoints joined by an edge of Γ outside that stretch, which by Theorem 7.12(b) happens if and only if Γ/S has no loop.

Proof. A Hamilton cycle containing S contains every stretch entirely and descends to a Hamilton cycle of Γ^b through all the marked edges; conversely such a cycle lifts, by reinserting the suppressed interiors. (A Hamilton cycle must use both edges at a vertex of degree 2, so it automatically contains every stretch of two or more edges; the hypothesis has content only for the stretches of a single edge.) Suppression replaces one edge at each stretch endpoint by another and does not change the degree of any surviving vertex, so by Lemma 5.7 the vertices of degree below 4 are the $2c$ stretch endpoints. A parallel edge appears precisely when the marked edge duplicates one already present, that is when the two endpoints of the stretch were already adjacent; they are joined by a chord exactly when the stretch plus that chord is a cycle of F , which is the description of the loops of Γ/S in Theorem 7.12(b). $\square$

Remark 5.10 (The marking cannot be dropped). It is tempting to read Proposition 5.9 as saying that Γ^b is itself uniquely Hamiltonian, which would turn each value of c into a question about uniquely Hamiltonian graphs with few vertices of degree 3. That reading is false, and not marginally: for the eleven uniquely Hamiltonian classes of $10 \leq n \leq 13$ the quotient Γ^b has between 4 and 10 Hamilton cycles, exactly one of which passes through all three marked edges. Nothing in Γ^b forces a marked edge, because suppression is precisely the removal of the degree-2 vertices that did the forcing. This is the same phenomenon as the port system of Theorem 7.12: suppression and contraction are two views of one obstruction, and in both the combinatorial data that survives is a graph *with a marking*, not a graph.

Applied with $s = 5$, suppression turns the five possible values of c into five concrete questions about graphs of maximum degree 4 carrying c marked edges:

c	(n_2, n_3) in Γ	vertices of Γ^b	degree-3 vertices of Γ^b	observed
1	(4, 2)	$n - 4$	2	no
2	(3, 4)	$n - 3$	4	no
3	(2, 6)	$n - 2$	6	yes
4	(1, 8)	$n - 1$	8	no
5	(0, 10)	n	10	no

The last row is the one that needs no suppression and no marking, and it is therefore the only one that can be compared with the literature directly: for $c = 5$ the graph Γ itself is a simple uniquely Hamiltonian graph with $\delta = 3$, $\Delta = 4$ and $n_3 = 10$, whose ten degree-3 vertices are perfectly matched by five edges of its Hamilton cycle. Since the least number of degree-3 vertices occurring in the catalogue of §9 is 14, ruling out $c = 5$ amounts to the self-contained statement that a uniquely Hamiltonian graph of maximum degree 4 whose degree-3 vertices are matched along its Hamilton cycle cannot have exactly ten of them. The four other rows, by Remark 5.10, are questions about marked graphs and are *not* comparable with that catalogue: a graph of maximum degree 4 with four vertices of degree 3 and a single Hamilton cycle through two marked edges is a far weaker object than a uniquely Hamiltonian one, and such objects are plentiful.

For Γ_n and Δ_n the suppressed graph is in any case not simple. Whenever a degree-2 vertex sits inside a triangle of F whose third edge is a chord, the marked edge created by suppression runs parallel to that chord. In Δ_n both degree-2 vertices do; in Γ_n only the vertex 2 does, the stretch 4–5–6 having 4 and 6 non-adjacent. These are the loops that Theorem 7.12(b) predicts, one for Γ_n and two for Δ_n .

Remark 5.11 (A search directed by the stretch pattern). Lemma 5.7 splits the classification into seven independent searches, one for each of the seven compositions of 5 into unordered parts, and each of them is far cheaper than a traversal of all 2-factors: once the pattern is fixed, one places its stretches around H — putting one of them at the edge $(0, 1)$ to kill the rotations — and then completes F with chords only, since by construction no further edge of

H may be used. We did this exhaustively for $10 \leq n \leq 15$. The counts below are of labelled pairs, so that each column sums to the total number of 2-factors of $[n]$ with $s = 5$; for $n \leq 13$ that total is known independently and the check holds.

pattern	(n_2, n_3)	$n = 10$	$n = 11$	$n = 12$	$n = 13$	$n = 14$	$n = 15$
$\{5\}$	$(4, 2)$	20	154	1 200	10 244	97 160	1 015 965
$\{1, 4\}$	$(3, 4)$	380	2 739	22 752	211 627	2 178 708	24 624 330
$\{2, 3\}$	$(3, 4)$	400	2 937	24 264	224 328	2 296 308	25 829 055
$\{1, 1, 3\}$	$(2, 6)$	2 290	18 821	175 536	1 828 554	20 929 846	260 703 600
$\{1, 2, 2\}$	$(2, 6)$	2 540	20 438	188 208	1 943 955	22 102 514	273 832 800
$\{1, 1, 1, 2\}$	$(1, 8)$	4 840	57 706	681 912	8 468 603	111 967 408	1 578 450 570
$\{1, 1, 1, 1, 1\}$	$(0, 10)$	586	14 476	257 760	4 218 435	68 586 868	1 141 628 352
total		11 056	117 271	1 351 632	16 905 746	228 158 812	3 306 084 672

In every one of them the only uniquely Hamiltonian pairs occur in the row $\{1, 2, 2\}$, in 2, 2, 3, 4, 6 and 9 dihedral classes respectively. The six other rows account for 206 056 298 pairs at $n = 14$ and 3 032 251 872 at $n = 15$, and not one of them is uniquely Hamiltonian. We also ran one size further the two rows that the search of Remark 5.12 singles out as the most dangerous: at $n = 16$ the pattern $\{1, 1, 3\}$ has 3 509 167 328 labelled pairs and the pattern $\{1, 1, 1, 2\}$ has 23 691 129 856, and in neither of them is a single pair uniquely Hamiltonian. The second was computed in 120 independent pieces, one for each placement of its four stretches around H , which is also how we verified that the pieces sum to the whole.

The saving over the traversal of all 2-factors is of two kinds, and only one of them is due to the pattern decomposition. Fixing the pattern confines the search to the $s = 5$ layer, which at $n = 13$ has 16 905 746 members against 438 263 364 2-factors in all; and placing one stretch at the edge $(0, 1)$ divides the work by a further factor of about n . What the decomposition adds beyond speed is the breakdown itself: the table is what shows where the uniquely Hamiltonian pairs are not.

Remark 5.12 (The conjecture is tight, not comfortable). Beyond the exhaustive range we ran, for each pattern separately, a simulated annealing on the chords with the stretches held fixed, minimising the number of Hamilton cycles of Γ . A heuristic's negative answers are worth nothing until it is calibrated, so we read it only where its positive control works: with a budget of forty restarts and 25 000 moves it does reach a uniquely Hamiltonian pair in the pattern $\{1, 2, 2\}$ at $n = 16$ and at $n = 18$, and fails to at $n = 20$, where each of the five patterns we recorded saturates the counting cut-off. Only the two smaller sizes therefore carry information, and there the best configurations found have

pattern	$n = 16$	$n = 18$
$\{1, 2, 2\}$	1	1
$\{1, 1, 3\}$	2	3
$\{1, 1, 1, 2\}$	2	2
$\{1, 1, 1, 1, 1\}$	4	4
$\{2, 3\}$	4	4
$\{1, 4\}$	6	12
$\{5\}$	12	12

Hamilton cycles. Two of the six forbidden patterns stop at exactly two, one cycle away from refuting Conjecture 5.8; and the two that do are the ones closest to $\{1, 2, 2\}$, namely the other composition with $c = 3$ and the unique one with $c = 4$. By Lemma 7.5 read contrapositively, the second Hamilton cycle of such a configuration must give up an edge that is *not* shared, so its overlap with F is 6 — the same wall as in Remark 9.3. We record the conjecture because

the exhaustive data of Remark 5.11 supports it without exception up to $n = 15$, and in the two patterns that come closest, $\{1, 1, 3\}$ and $\{1, 1, 1, 2\}$, up to $n = 16$, not because the margin is comfortable; we would not be surprised to see it fail at some larger order.

Corollary 5.13. $\text{ov}(n) \leq 5$ for every $n \geq 10$. Combined with the exhaustive computation of Table 1 — a complete traversal, carried out by independent implementations that agree — $\text{ov}(n) = 5$ for $9 \leq n \leq 13$.

6 The lower bound and Sheehan’s conjecture

The inequality $\text{ov}(n) \geq 5$ remains. We now show that it is not elementary.

Theorem 6.1. Suppose $\text{ov}(n) \geq 1$ for every n . Then every 4-regular Hamiltonian graph has at least two Hamilton cycles.

Proof. Let G be a 4-regular Hamiltonian graph and let H be a Hamilton cycle of G . The subgraph $F = E(G) \setminus E(H)$ has all degrees equal to $4 - 2 = 2$, so it is a 2-factor of G , disjoint from H by construction. By hypothesis there is a Hamilton cycle H' of G with $|F \cap H'| \geq 1$. Since $F \cap H = \emptyset$, necessarily $H' \neq H$. $\square$

Sheehan’s conjecture [17], formulated in 1975, asserts precisely that every 4-regular Hamiltonian graph contains at least two Hamilton cycles, and it remains open. It is worth being precise about its place. For odd degree the statement is proved: Smith’s theorem, published by Tutte [20], asserts that every edge of a cubic graph lies in an even number of Hamilton cycles, so there are at least three, and Thomason’s lollipop lemma [18] extends the argument to every graph all of whose degrees are odd. For even degree, Petersen’s 2-factorisation theorem reduces the $2k$ -regular case to the 4-regular case: if G is $2k$ -regular with Hamilton cycle H , then $G - H$ is $(2k - 2)$ -regular and decomposes into $k - 1$ 2-factors, and adding one of them to H yields a 4-regular spanning subgraph containing H , so that any second Hamilton cycle of it is one of G . The 4-regular case is therefore the only open case of the conjecture for all degrees at least 3. Quantitatively, Thomassen [19] proved the statement for degree at least 300 and Haxell, Seamone and Verstraëte [13] lowered the threshold to degree at least 23. For small orders, Goedgebeur, Meersman and Zamfirescu [11] verified the conjecture by computer up to $n = 21$.

Corollary 6.2. Any universal bound of the form $\text{ov}(n) \geq 1$ is at least as hard as a problem that has been open since 1975.

This bounds in advance what can be expected from an elementary argument, and at the same time justifies the parameter: asking for $\text{ov}(n) \geq 1$ is asking “there is a second Hamilton cycle”, and asking for $\text{ov}(n) \geq s$ is asking “there is a Hamilton cycle recovering s edges of the 2-factor”. In that sense $\text{ov}(n)$ is a quantitative refinement of Sheehan’s conjecture, and Table 1 is a table of exact values of it.

7 Where the difficulty lies

7.1 The window $s \leq 4$

Proposition 7.1 (Reduction to $s \leq 4$). For every pair (F, H) we have $\text{ov}_\Gamma(F) \geq s = |F \cap H|$. Hence

$$\text{ov}(n) \geq 5 \iff \text{every pair with } s \leq 4 \text{ satisfies } \text{ov}_\Gamma(F) \geq 5.$$

Proof. H itself is a Hamilton cycle of Γ and $|F \cap H| = s$, which gives the inequality. If $s \geq 5$ the conclusion is immediate, so only the pairs with $s \leq 4$ can violate $\text{ov}(n) \geq 5$. $\square$

The exhaustive traversal by value of s shows that inside that window the statement not only holds but holds with room to spare.

$s = F \cap H $	0	1	2	3	4	5	6	7	8
$n = 10$	6	7	6	5	5	5	6	7	8
$n = 11$	7	6	6	6	6	5	6	7	8
$n = 12$	7	7	6	6	6	5	6	7	8

Table 3: Minimum overlap over the pairs with exactly s shared edges. For $s \geq 5$ the trivial bound $\geq s$ of Proposition 7.1 is attained. At $n = 11$ and $n = 12$ the global minimum is attained *only* at $s = 5$. For $n = 13$ we have traversed the case $s = 4$, with 47 187 049 pairs, and the minimum overlap is 6 there as well; a second algorithm traversed all 415 947 992 pairs of [13] with $s \leq 4$ and found none with overlap at most 4.

The shape of the table is a V with its vertex at $s = 5$, and it repays slow reading. The left end, $s = 0$, is the 4-regular case of Sheehan’s conjecture: it is where the statement is hardest to prove and, at the same time, where the truth appears to have the *most* room, since over the computed range the minimum overlap grows with n (Table 2), although that is only an empirical fact. The right end is trivial. All the tension lies in a narrow window.

7.2 Amplification

Lemma 7.2 (Amplification). *If Γ has a Hamilton cycle $H' \neq H$ with $S \subseteq H'$, then $\text{ov}_\Gamma(F) \geq |F \cap H'| \geq s + 2$.*

Proof. Since H and H' both have n edges, put $t := |H' \setminus H| = |H \setminus H'|$. We check that $t \neq 1$: if $H' = H - e + f$ with $e \neq f$, then $H - e$ is a Hamilton path with endpoints u, v , and the only edge closing it into a cycle through all vertices is $uv = e$, contradicting $f \neq e$. Hence $t \geq 2$. Every edge of $H' \setminus H$ lies outside H and therefore, since $\Gamma = F \cup H$, belongs to F ; that is, H' uses t chords of F . Together with $S \subseteq H'$ this gives $|F \cap H'| \geq s + t \geq s + 2$. $\square$

Corollary 7.3 (Rigidity of the extremal pairs). *If $\text{ov}_\Gamma(F) = s$, then no Hamilton cycle of Γ other than H contains all the edges of $F \cap H$.*

The family Γ_n satisfies Corollary 7.3 for the strongest possible reason: it has no other Hamilton cycle at all. Iterating the lemma yields the engine that reduces the lower bound to an existence statement.

Proposition 7.4 (Iteration engine). *Suppose every pair (F, H) with $|F \cap H| \leq 4$ admits a Hamilton cycle $H' \neq H$ with $F \cap H \subseteq H'$. Then $\text{ov}(n) \geq 5$ for every n .*

Proof. Let (F, H) have $s \leq 4$ and let H' be the cycle given by the hypothesis. By Lemma 7.2, $s' := |F \cap H'| \geq s + 2$. If $s' \geq 5$ we are done. Otherwise apply the hypothesis to the pair (F, H') , whose graph $F \cup H'$ is a subgraph of Γ , so that any Hamilton cycle found there is one of Γ . Each step increases the overlap by at least 2, so at most three steps suffice to pass from $s = 0$ to $s \geq 5$. $\square$

7.3 Protected edges

Lemma 7.2 says that a second Hamilton cycle respecting the shared edges improves the overlap. Read backwards, it imposes a very strong restriction on the pairs realising the minimum.

Lemma 7.5 (Protected edges). *Let $\Gamma = F \cup H$, $S = F \cap H$, $s = |S|$, and suppose H is optimal, that is $\text{ov}_\Gamma(F) = s$. Then every Hamilton cycle H' of Γ satisfies*

$$H \setminus H' \subseteq S.$$

In words: a competing Hamilton cycle can only give up shared edges; the $n - s$ edges of H that are not in F are protected.

Proof. Let $R = H \setminus H'$ and $t = |R| = |H' \setminus H|$. Every edge of $H' \setminus H$ lies outside H and therefore, since $\Gamma = F \cup H$, belongs to F ; hence $(F \setminus H) \cap H' = H' \setminus H$ has t elements. On the other hand $(F \cap H) \setminus H' = S \cap (H \setminus H') = S \cap R$, so $|(F \cap H) \cap H'| = s - |S \cap R|$. Adding,

$$|F \cap H'| = t + s - |S \cap R|.$$

Optimality of H gives $|F \cap H'| \leq s$, whence $t \leq |S \cap R|$. But $S \cap R \subseteq R$ and $|R| = t$, so $|S \cap R| \leq t$. Therefore equality holds, $S \cap R = R$, that is $R \subseteq S$. $\square$

Corollary 7.6 (Rigidity). *Under the same hypotheses, every Hamilton cycle H' of Γ satisfies $|H' \setminus H| \leq s$, and hence $|H_1 \setminus H_2| \leq 2s$ for any two Hamilton cycles of Γ .*

Corollary 7.7 (Structure and counting). *Under the same hypotheses, every Hamilton cycle $H' \neq H$ of Γ is obtained by choosing a subset $R \subseteq S$ with $|R| \geq 2$ and reassembling cyclically the $|R|$ paths of $H \setminus R$ by means of $|R|$ chords of F . Consequently*

$$\#\{\text{Hamilton cycles of } \Gamma\} \leq 1 + \sum_{t=2}^s \binom{s}{t} (t-1)! 2^{t-1},$$

a bound depending only on s and not on n : it equals 1, 1, 3, 15, 93 for $s = 0, 1, 2, 3, 4$.

Proof. By Lemma 7.5, $R = H \setminus H' \subseteq S$; and $|R| \neq 1$ by the argument in the proof of Lemma 7.2. The edges of H' not in $H \setminus R$ are chords of F , $|R|$ in number, and they assemble cyclically the $|R|$ paths of $H \setminus R$; fixing one of them unreversed, the number of cyclic orderings with orientations is $(|R| - 1)! 2^{|R|-1}$. Since H' determines R , it suffices to sum over R . $\square$

The content of Corollary 7.7 is that in an optimal pair the whole Hamiltonian structure of Γ is confined to the s shared edges: outside them, Γ behaves as though H were unique. In the window $s \leq 4$ this means that Γ has at most 93 Hamilton cycles, no matter how many vertices it has. The bound is loose — in the exhaustive traversal of all 2-factors of $[n]$ for $n \leq 10$ there are 15, 41, 148 and 551 pairs with H optimal for $n = 7, 8, 9, 10$, and in all of them the number of Hamilton cycles is at most 4, while the extremal families of Section 5 have exactly one — but its independence of n is what matters.

Corollary 7.8. *If $\text{ov}_\Gamma(F) \leq 1$ then H is the unique Hamilton cycle of Γ . Consequently, $\text{ov}(n) \geq 2$ is equivalent to the non-existence of a uniquely Hamiltonian union $F \cup H$ with $|F \cap H| \leq 1$: the case $|F \cap H| = 0$ is exactly Sheehan's conjecture, and the case $|F \cap H| = 1$ is that of the unions with exactly two vertices of degree 3, necessarily adjacent, and all others of degree 4.*

Proof. Suppose $\text{ov}_\Gamma(F) \leq 1$. Since $s \leq \text{ov}_\Gamma(F)$ we have $s \leq 1$, and it remains to rule out that H fails to be optimal. Were that so with $s = 0$ and $\text{ov}_\Gamma(F) = 1$, a Hamilton cycle H' with $|F \cap H'| = 1$ would satisfy $F \cap H' = (F \setminus H) \cap H' = H' \setminus H$, so $|H' \setminus H| = 1$, impossible by the argument in the proof of Lemma 7.2. So H is optimal with $s \leq 1$, Corollary 7.7 gives the bound 1, and therefore $H' = H$. The converse and the description of the degrees follow from (1). $\square$

Finally, the lemma yields a reformulation of the bound we are after in purely local terms.

Corollary 7.9 (Reformulation). *$\text{ov}_\Gamma(F) \geq s + 1$ if and only if Γ has a Hamilton cycle H' with $H \setminus H' \not\subseteq F \cap H$, that is, one that gives up some edge of H not shared with F .*

Proof. If $\text{ov}_\Gamma(F) = s$ then H is optimal and Lemma 7.5 gives $H \setminus H' \subseteq S$ for every H' ; contrapositively, the existence of such an H' forces $\text{ov}_\Gamma(F) \geq s + 1$. Conversely, if every Hamilton cycle H' satisfies $H \setminus H' \subseteq S$, then in the notation of the proof of Lemma 7.5 we have $|S \cap R| = |R| = t$ and therefore $|F \cap H'| = t + s - |S \cap R| = s$ for every H' , whence $\text{ov}_\Gamma(F) = s$. $\square$

This reduces the question to a statement without maximisations: in the window $s \leq 4$ there are at least $n - 4$ unshared edges, and it suffices to find a Hamilton cycle sacrificing any one of them. Lemma 7.5 has moreover been verified exhaustively: among the 465, 3507, 30016 and 286884 2-factors of [7], [8], [9] and [10], the 15, 41, 148 and 551 pairs with H optimal satisfy it without exception.

7.4 Alternating circuits

There is a second reading of Lemma 7.2 worth recording, because it connects with the classical description of adjacency in the skeleton of the 2-factor polytope and because it explains the shape of the families of Section 5.

Since F and H are both 2-factors, every vertex has degree 0, 2 or 4 in $F \triangle H$, with as many edges of $F \setminus H$ as of $H \setminus F$. Call *alternating decomposition* a partition of $F \triangle H$ into closed circuits alternating edges of $F \setminus H$ with edges of $H \setminus F$. One always exists, but it is *not unique*: at a vertex of degree 4 there are two ways of pairing its two F -edges with its two H -edges, and different decompositions may have different numbers of circuits. Fix one, $C_1, \dots, C_k$. Note that $S = F \cap H$ is disjoint from $F \triangle H$ and that $|F \triangle H| = 2(n - s)$, so $k \leq (n - s)/2$.

Lemma 7.10 (Alternation). *For $T \subseteq \{1, \dots, k\}$ let $H_T = H \triangle (\bigcup_{i \in T} C_i)$. Then H_T is a 2-factor of Γ with*

$$|F \cap H_T| = s + \sum_{i \in T} \frac{1}{2} |C_i|.$$

Consequently, if H is optimal then no H_T with $T \neq \emptyset$ is a Hamilton cycle: all of them have at least two cycles.

Proof. Toggling an alternating circuit exchanges, at every vertex it meets, as many edges of H as of F , so it preserves 2-regularity and H_T is a 2-factor. Its edges in F are those of S , which no C_i meets, together with the $\frac{1}{2}|C_i|$ edges of $F \setminus H$ of each toggled circuit. If some H_T with $T \neq \emptyset$ were Hamiltonian, its overlap with F would be $s + \sum_{i \in T} \frac{1}{2}|C_i| \geq s + 2$, contradicting the optimality of H . $\square$

Remark 7.11 (What the lemma gives and what it does not). For T a single index, the lemma says that toggling any alternating circuit of $F \triangle H$ must disconnect; in Chvátal's language, no neighbour of H in the skeleton of the 2-factor polytope can be Hamiltonian. This is a necessary condition that is cheap to check, but its strength depends on the chosen decomposition: it imposes $2^k - 1$ restrictions, so it only bites when $F \triangle H$ admits decompositions with many circuits. And since the decomposition is not unique, it suffices that *some* decomposition have $k = 1$ for the condition to reduce to the statement that F is not Hamiltonian, which is automatic. This is exactly what happens for the family Γ_n : we have checked that $F_n \triangle H$ is connected with degrees 2 and 4 and admits a decomposition into a single alternating circuit of length $2(n - 5)$, for every $10 \leq n \leq 23$. The lemma is therefore not a route to the lower

bound; it is a design principle for constructing rigid pairs, which should admit alternating decompositions with few circuits.

7.5 Contraction: a 4-regular multigraph with forbidden transitions

Theorem 7.12 (Contraction). *Suppose $F \neq H$, so that S is a proper subgraph of H and its components are paths. Let Γ/S be the multigraph obtained by contracting each component of S to a single vertex, called a block. Then:*

- (a) Γ/S has exactly $n - s$ vertices and is 4-regular, loops counting with multiplicity 2;
- (b) the loops of Γ/S correspond to the cycles of F having exactly one edge outside H ;
- (c) the Hamilton cycles of Γ containing S correspond to the Hamilton cycles of Γ/S that are compatible with the port system: at the block coming from a stretch with endpoints u, w , the four incident edges split into the two incident with u and the two incident with w , and the cycle must use exactly one from each pair.

Proof. (a) S is a forest of paths with s edges on the vertex set $[n]$, so, counting its isolated vertices, it has $n - s$ components, and that is the number of blocks. (Not to be confused with the c of Lemma 5.7, which counts only the components carrying an edge.) For the degree, a stretch of length ℓ has $\ell - 1$ interior vertices, whose two F -edges are H -edges and which therefore have degree 2 in Γ , and two endpoints, each with one edge of S , the other H -edge and one chord, hence of degree 3. The degree sum of the stretch is $2(\ell - 1) + 6 = 2\ell + 4$, and subtracting 2ℓ for the ℓ internal edges leaves 4. A singleton block is a vertex both of whose F -edges are chords, of degree 4.

(b) A loop at the block of a stretch T is an edge of Γ with both endpoints in T that is not in S : necessarily a chord of F joining the two endpoints of T , and then T together with that chord is a cycle of F with a single edge outside H . The converse is identical.

(c) A Hamilton cycle H' of Γ with $S \subseteq H'$ enters each stretch at one endpoint and leaves at the other, since its interior vertices have degree 2 and their edges already lie in H' ; hence it uses exactly one edge at u and one at w besides those of S , and contracting yields a compatible Hamilton cycle of Γ/S . Conversely, a compatible cycle lifts by reinserting the whole stretch. $\square$

Remark 7.13 (Compatibility cannot be dispensed with). Without condition (c) the statement would be false, and the smallest example is instructive. Take $n = 8$, $H = (0, 1, \dots, 7, 0)$ and

$$F = \{06, 65, 57, 70\} \cup \{12, 24, 43, 31\},$$

two quadrilaterals, with $S = \{65, 70, 12, 43\}$ and $s = 4$. The four blocks are $\{5, 6\}$, $\{0, 7\}$, $\{1, 2\}$ and $\{3, 4\}$, and Γ/S consists of two pairs of blocks joined by triple edges together with two single edges: it has nine Hamilton cycles. But only one of them is compatible, namely H . Each of the other eight enters and leaves some block through two legs attached to the same endpoint of its stretch, which no Hamilton cycle of Γ can do. Consequently $\text{ov}_\Gamma(F) = 4 = s$: this is one of the extremal pairs for $n = 8$.

If $s = 0$ there are no stretches, all blocks are singletons and the port system is empty: the hypothesis of Proposition 7.4 is then exactly Sheehan's conjecture. If $1 \leq s \leq 4$ there are at most four constrained blocks and at least $n - 8$ free ones.

Corollary 7.14 (Bracketing). *Writing (H_4) for the statement “every quotient Γ/S with at most four constrained blocks has a second compatible Hamilton cycle”, we have*

$$(H_4) \implies \text{ov}(n) \geq 5 \implies \text{Sheehan's conjecture}.$$

Remark 7.15 (The bracketing is not tight in an obvious way). It would be tempting to state (H₄) for *arbitrary* 4-regular multigraphs, in which case the bracketing would look like a purely local strengthening of Sheehan’s conjecture. That unrestricted version is false, and trivially so: let L_k be the cycle C_k with a loop added at each of its k vertices. A loop contributes 2 to the degree of its vertex, so L_k is 4-regular, and no Hamilton cycle can use a loop, so C_k is the only Hamilton cycle of L_k . Loops are not an artefact of this example: by Theorem 7.12(b) they appear in Γ/S exactly when a cycle of F has a single edge outside H . Hence the quotients Γ/S , with their port systems, must be kept as a proper subclass: for $s = 0$ the quotient is Γ itself, which is simple and unmarked, so that instance of (H₄) is precisely Sheehan’s conjecture for simple graphs, and the restriction only begins to matter for $s \geq 1$. The distance between the two ends of the bracketing is therefore twofold: a local condition at at most four vertices, and the passage from simple graphs to this family of multigraphs.

7.6 An obstacle to reducing the bracketing to Sheehan’s conjecture

Corollary 7.14 leaves a gap between Sheehan’s conjecture and its version with forbidden transitions. The natural way to close it would be to replace each constrained block by a fragment imposing the port system by its own structure, so that the resulting graph is 4-regular with no marks and ordinary Sheehan suffices. We make precise what would be needed.

Definition 7.16 (Port block). A *port block* of size m is a connected graph G_0 on m vertices in which exactly two vertices x, y have degree 2, all the others have degree 4, and there is *exactly one* Hamilton path from x to y .

Plugging two legs at x and two at y , a port block becomes 4-regular, can only be traversed by entering through a leg at x and leaving through a leg at y — its other internal vertices are not reachable otherwise — and the internal route is determined. Replacing each stretch of S by a copy would yield a 4-regular graph whose Hamilton cycles are in bijection with those of Γ containing S . It should be noted that this alone would not suffice to invoke Sheehan’s conjecture: the quotient Γ/S may have loops and multiple edges — Theorem 7.12(b) describes the former and Remark 7.13 exhibits triple edges — so the substituted graph need not be simple, and Sheehan’s conjecture is stated for simple graphs. Closing the gap of Corollary 7.14 would require, besides the block, a condition guaranteeing simplicity of the result. Uniqueness of the path is indispensable: with two internal routes, a second Hamilton cycle of the 4-regular graph could differ only inside a block and produce no new cycle in Γ .

Proposition 7.17. *There is no port block of size $m \leq 13$. Moreover, no fragment with an odd number of legs can have all its internal vertices of degree 4.*

Proof. The second statement is a parity argument: if a fragment has m internal vertices, all of degree 4 counting the legs, and E internal edges, then $4m = 2E + (\text{number of legs})$, so the number of legs is even. This rules out treating separately the degree-3 vertices of Γ , which come in pairs as the endpoints of a single stretch, and forces one to substitute the whole stretch, with its four legs.

The first statement is an exhaustive enumeration, preceded by an observation ruling out the smaller sizes: if $m \leq 4$ the maximum degree of a simple graph on m vertices is $m - 1 \leq 3$, so there are no vertices of degree 4; and if $m = 5$ the three vertices of degree 4 are adjacent to all the others, so x and y would have degree at least 3. Hence $m \geq 6$. A port block of size m has $\frac{1}{2}(2 \cdot 2 + 4(m - 2)) = 2m - 2$ edges. We generated with `nauty` all connected graphs with m vertices, $2m - 2$ edges and degrees between 2 and 4, filtered those with exactly two vertices of degree 2, and counted the Hamilton paths between them. The generator produces 7, 28, 124, 663, 4 173, 30 037, 243 212 and 2 166 792 graphs for $m = 6, \dots, 13$, of which respectively 1, 4,

13, 50, 235, 1 284, 8 392 and 62 439 pass the degree filter. In none of them is the Hamilton path between the two degree-2 vertices unique. $\square$

Substitution by blocks therefore does not close the gap of Corollary 7.14, at least not with small blocks. Note that there is no parity obstruction to the existence of a port block: by Thomason’s lollipop lemma [18] the number of Hamilton paths starting at x and ending at a vertex of even degree is even, and since all degrees of G_0 are even this only forces the total number of Hamilton paths from x to be even, which is consistent with exactly one of them ending at y . Whether a port block of larger size exists remains open, and is a concrete problem in its own right.

8 The weight ladder, and open problems

The pieces above arrange themselves on a single scale. Call a pair (F, H) *rigid* if H is optimal, that is $\text{ov}_\Gamma(F) = s$; by Lemma 7.5 these are exactly the pairs in which every Hamilton cycle contains $H \setminus F$. Call the *weight* of $\Gamma = F \cup H$ the integer $w(\Gamma) = n_3 + 2n_2 = 2s$, by (1). Then

$$\text{ov}(n) = \frac{1}{2} \min\{w(\Gamma) : (F, H) \text{ rigid on } n \text{ vertices}\}.$$

Indeed a rigid pair satisfies $\text{ov}_\Gamma(F) = s = \frac{1}{2}w(\Gamma)$, which gives $\leq$; and conversely, if (F, H) attains $\text{ov}(n)$ and H^* is a maximiser, then $\Gamma^* = F \cup H^* \subseteq \Gamma$ satisfies $\text{ov}_{\Gamma^*}(F) = |F \cap H^*| = \text{ov}(n)$, so (F, H^*) is rigid of weight $2 \text{ov}(n)$.

weight	what a rigid pair of that weight would be	status
0	Γ is 4-regular and uniquely Hamiltonian	Sheehan’s conjecture , open
2	Γ uniquely Hamiltonian, $n_3 = 2$, rest of degree 4	open
4	rigid pair with $s = 2$, at most 3 Hamilton cycles	open
6	rigid pair with $s = 3$, at most 15 Hamilton cycles	open
8	rigid pair with $s = 4$, at most 93 Hamilton cycles	open
10	exists: the families Γ_n and Δ_n , for every $n \geq 10$	Theorems 5.3, 5.5

The bounds on the number of Hamilton cycles are those of Corollary 7.7; for weights 0 and 2 they equal 1, that is, a rigid pair of those weights is necessarily uniquely Hamiltonian. In this language Conjecture 8.1 asserts that the minimum weight is 10, whereas Sheehan’s conjecture asserts only that it is not 0: between the two there are four rungs. No rigid pair of weight at most 8 exists for $9 \leq n \leq 13$ — for $n = 6, 7, 8$ such pairs do exist, which is exactly the statement $\text{ov}(n) = 4$ there; for $n = 13$ we checked this in two independent ways, by the complete traversal of the 438 263 364 2-factors of [13] with the cor computation of Lemma 2.1, and by a second algorithm traversing only the 415 947 992 pairs with $s \leq 4$.

Conjecture 8.1. $\text{ov}(n) = 5$ for every $n \geq 9$.

Remark 8.2 (The window has no easy cases). One might hope that some value of s inside the window is more tractable than Sheehan’s case $s = 0$, on the grounds that odd-degree vertices are available for the lollipop lemma. This is not so. Thomason’s lemma [18] requires *all* vertices to have odd degree; by contrast, by (1), a pair with s shared edges distributed over c stretches has, by Lemma 5.7, $n_3 = 2c \leq 2s \leq 8$ vertices of degree 3 and at least $n - 8$ of degree 4. The proportion of even-degree vertices tends to 1 and the lemma contributes nothing anywhere in the window: there is no easier subcase to enter. At the opposite end, when Γ is cubic all degrees equal 3 and the lemma does apply, but then $s = n/2$ and the trivial bound $\text{ov}_\Gamma(F) \geq s$ already gives far more than 5.

Remark 8.3 (There is one unit of slack where it matters). Combining Proposition 7.1 with Table 3 gives a finer reading of what would have to be proved. The content of $\text{ov}(n) \geq 5$ lies entirely in the window $s \leq 4$; but in that window the measured minimum is not 5 but 6: so it is for $n = 11$ and $n = 12$ at every value $s \leq 4$, and for $n = 13$ in the case $s = 4$. A directed search restricted to $s \leq 4$ at $n = 14$ and $n = 15$ also finds no pair with overlap less than 6. The value 5 of $\text{ov}(n)$ is contributed exclusively by the pairs with $s = 5$, where the bound is trivial by Proposition 7.1. In other words, in the region where the problem has content there is one unit of slack, and the statement the data support is

$$s \leq 4 \implies \text{ov}_\Gamma(F) \geq 6 \quad (n \geq 11),$$

strictly stronger than what Conjecture 8.1 needs. For $n \leq 10$ it is false: at $n = 10$ there are pairs with $s = 3$ and with $s = 4$ and overlap 5, and at $n = 8$ there are pairs with $s = 4$ and overlap 4, such as the one in Remark 7.13.

Remark 8.4 (A jump is not a chain). It is worth separating this parameter from those of reconfiguration. If, following [15], one defines the graph of k -switches on the 2-factors of G , joining F and F' when $|F \triangle F'| \leq 2k$, and writes $k^*(G)$ for the least k such that every component contains a Hamiltonian 2-factor, then $\text{cor}_G(F)$ bounds $k^*(G)$ but the bound is very loose. In the extremal pair for $n = 13$ every Hamilton cycle is at distance 8 from F , while a chain of moves of order 4 leads from F to one of them; and in a sample of 400 random pairs for each n between 10 and 14, the maximum of cor rises from 5 to 6 while the maximum of k^* stays at 4. The two parameters have different extremal families: the overlap measures a single jump, the reconfiguration order allows chains.

Open problems.

1. **The bound with slack.** Prove the inequality of Remark 8.3, namely $\text{ov}_\Gamma(F) \geq 6$ for $s \leq 4$ and $n \geq 11$. It is stronger than Conjecture 8.1 and hence also implies Sheehan's conjecture, but it has one unit of room that a counting argument might exploit and that the formulation with 5 does not offer. The natural subcase to start from is s equal to the number of stretches, in which every constrained block of Theorem 7.12 has exactly two vertices.
2. **Port blocks.** Decide whether a port block in the sense of Definition 7.16 exists. By Proposition 7.17 there is none with $m \leq 13$; an affirmative answer for some m , together with a simplicity condition, would narrow the gap of Corollary 7.14. A general negative answer would be equally informative: it would say that the version with transitions is strictly stronger.
3. **Uniquely Hamiltonian graphs with maximum degree 4.** By Corollary 7.3 and the weight ladder, Conjecture 8.1 is related to determining the minimum of $n_3 + 2n_2$ over the uniquely Hamiltonian graphs that are the union of a 2-factor and a Hamilton cycle. Over the *rigid* pairs, which is the notion the ladder uses, that minimum is $2\text{ov}(n)$: it equals 8 for $6 \leq n \leq 8$, equals 10 for $9 \leq n \leq 13$, and would be 0 if Sheehan's conjecture were false. Over the uniquely Hamiltonian pairs it equals 10 for every $5 \leq n \leq 13$ except $n = 6$, where it is 8. Ruling out the value 8 for large n is a concrete goal. Section 9 discusses what the literature does and does not provide here.
4. **Chains rather than jumps.** Give a bound on k^* that does not pass through cor , in view of Remark 8.4.

9 Relation to uniquely Hamiltonian graphs

By Lemma 7.5, an optimal pair is one in which every Hamilton cycle contains $H \setminus F$, and the simplest way to achieve this is for Γ to be uniquely Hamiltonian, as happens for Γ_n and Δ_n . The problem therefore touches the literature on uniquely Hamiltonian graphs, and it is worth being precise about what that literature does and does not provide. The bridge is the degree count (1).

Remark 9.1 (Degree translation). Let $\Gamma = F \cup H$ with $S = F \cap H$ distributed over c stretches, so that by Lemma 5.7 Γ has $n_2 = s - c$ vertices of degree 2, $n_3 = 2c$ of degree 3 and $n - s - c$ of degree 4. In particular:

- (a) if all stretches have length 1, that is $s = c$, then Γ has no vertex of degree 2: it is a graph with degrees in $\{3, 4\}$ and exactly $2s$ vertices of degree 3;
- (b) conversely, if Γ_0 is a graph with degrees in $\{3, 4\}$ and Hamilton cycle H_0 , then $\Gamma_0 = F_0 \cup H_0$ with F_0 a 2-factor if and only if the degree-3 vertices of Γ_0 admit a perfect matching consisting of edges of H_0 ; in that case $s = n_3/2$.

Proof. Part (a) is immediate from Lemma 5.7. For (b), if $M \subseteq H_0$ is such a matching, put $F_0 = (\Gamma_0 \setminus H_0) \cup M$: in $\Gamma_0 \setminus H_0$ the degree-3 vertices have degree 1 and the degree-4 vertices degree 2, and M covers the former exactly once, so F_0 is 2-regular. Conversely, if F_0 is a 2-factor then $S = F_0 \cap H_0$ must raise to 2 the degree of every degree-3 vertex and of no other, hence it is a perfect matching of those vertices contained in H_0 . $\square$

Remark 9.2 (What the literature says, and what it does not). (1) **Nearly cubic graphs.**

Entringer and Swart [6] constructed uniquely Hamiltonian graphs with exactly two vertices of degree 4 and all the others of degree 3; Goedgebeur, Meersman and Zamfirescu [11] determined that these exist exactly for even $n \geq 18$. By Remark 9.1, one of them admitting the decomposition yields a uniquely Hamiltonian union with $s = (n - 2)/2$, far above 5. These families do not threaten Conjecture 8.1; they accompany it.

- (2) **Minimum degree 4.** Fleischner [7] constructed uniquely Hamiltonian graphs of minimum degree 4, answering a question of Bondy; his graphs have vertices of degree 4 and 14. Since $\Delta = 14 > 4$, they are *not* unions of a 2-factor and a Hamilton cycle, so that construction produces no counterexamples in our family. It is worth noting that his construction and ours approach Sheehan's conjecture from opposite sides: his graphs have minimum degree 4 but unbounded maximum degree, while Γ_n and Δ_n have maximum degree 4 and only eight vertices of smaller degree.
- (3) **Degrees $\{3, 4\}$: the risk lies here.** Brinkmann and De Pauw [2] proved that every degree set with minimum 3 containing an even number is realisable by a uniquely Hamiltonian graph; in particular $\{3, 4\}$ is. Goedgebeur, Jooker, Lo, Seamone and Zamfirescu [10] determined that the smallest uniquely Hamiltonian graph all of whose degrees are 3 or 4 has order 18. By Remark 9.1(a), a uniquely Hamiltonian union with unit stretches and s shared edges is a graph with degrees $\{3, 4\}$ and $2s$ vertices of degree 3. Conjecture 8.1 therefore implies the following concrete statement, which can be checked against the existing catalogues:

every uniquely Hamiltonian graph with degrees in $\{3, 4\}$ whose degree-3 vertices admit a perfect matching formed by edges of its Hamilton cycle has at least ten vertices of degree 3.

The known constructions do not threaten this. Those of [10] and [2] replace degree-3 vertices of a $\{3, 4\}$ -graph by triangles, so their number of degree-3 vertices *grows* with

n ; in the base graph of [2, Lemma 3.6] it is the number of degree-4 vertices that stays fixed, at six, and the degree-3 vertices that accumulate. Nor do the catalogues. The complete list of uniquely Hamiltonian graphs with minimum degree at least 3, girth at least 5 and at most 22 vertices — 63 graphs, computed in [11] and distributed through House of Graphs [12] — splits by degree profile as follows; we have recomputed the profiles from the published `graph6` files.

n	graphs	$(n_3, n_4, \dots)$
18	2	(14, 4), (16, 2)
19	1	(16, 3)
20	2	(18, 2)
21	25	(14, 7), (16, 5), (18, 3), and 11 graphs with $\Delta = 5$
22	33	(16, 6), (18, 4), (20, 2), and 13 graphs with $\Delta = 5$

Thirty-nine of the 63 have maximum degree 4, and the smallest number of degree-3 vertices occurring anywhere in the list is 14: every entry satisfies the displayed statement with a margin of at least four. We know of no theorem, on the other hand, forbidding a counterexample, and the catalogue is complete only for girth at least 5.

- (4) **The classical bounds obstruct nothing here.** Bondy and Jackson [3] proved that a uniquely Hamiltonian graph on n vertices has a vertex of degree $O(\log n)$, and Abbasi and Jamshed [1] improved this by means of the inequality $\sum_v (2/3)^{d(v)-1} \geq 1$, valid for even n . Evaluated in our family with $n_3 = 2s$ and $n_4 = n - 2s$ it gives

$$2s \cdot \frac{4}{9} + (n - 2s) \cdot \frac{8}{27} \geq 1,$$

which holds trivially as soon as $n \geq 4$. Nor does Thomason’s lemma, which forces at least two vertices of even degree, say anything: in the window $s \leq 4$ there are at least $n - 8$ of them. None of the available tools rules out a counterexample.

- (5) **How close to 4-regular a uniquely Hamiltonian graph can be.** Read as statements about uniquely Hamiltonian graphs rather than about overlaps, Theorems 5.3 and 5.5 say this: *for every $n \geq 10$ there is a simple uniquely Hamiltonian graph on n vertices with exactly two vertices of degree 2, six of degree 3, and $n - 8$ of degree 4.* The number of vertices of degree less than 4 is therefore bounded independently of the order, and Sheehan’s conjecture is exactly the assertion that it can never be 0. We have not found this observation in the literature, where the known constructions run the other way: in the catalogue tabulated above the number of degree-3 vertices is at least 14, Entringer and Swart’s nearly cubic graphs have $n - 2$ of them, and in the $\{3, 4\}$ constructions of [2, 10] it is the number of degree-4 vertices that stays bounded. Together with Fleischner’s graphs of degrees 4 and 14, which have minimum degree 4 but unbounded maximum degree, these families bracket Sheehan’s conjecture from the two available sides. The ladder of §8 asks for the exact minimum of $n_3 + 2n_2$ among such graphs; our families place it at 10 or below, and 0 is Sheehan.

Remark 9.3 (Directed searches for uniquely Hamiltonian unions). We have searched for uniquely Hamiltonian unions with $s \leq 4$ by three methods, all starting from Γ_n and moving by 2-opt on the 2-factor: hill climbing that does not allow s to increase, annealing that does, and a variant fixing $s = 4$ and minimising the number of Hamilton cycles directly. The first two do not go below $s = 5$ for $n \in \{12, 14, 16, 18, 20\}$. The third reaches 2 Hamilton cycles for $n = 14$ and 4 for $n = 16$ and $n = 18$, without reaching one; and in those cases the second cycle gives up an edge *not* shared, so its overlap with F is 6, exactly what Lemma 7.5 predicts read contrapositively. The proximity to a single cycle is nevertheless a reminder that the exhaustive verification of the window $s \leq 4$ only reaches $n = 13$.

Code and data. Every computation reported here can be rerun from the repository

<https://github.com/carlitosandresv1-ctrl/Hamiltonicity-overlap>

which contains the exhaustive generator of 2-factors with the test of Lemma 2.1; the dynamic-programming computation of the maximum overlap and the cross-validation of the two against each other; the window check of the pairs with $s \leq 4$; the generators of Γ_n and Δ_n with the forcing propagation that verifies their unique Hamiltonicity; the classification of the extremal pairs under the dihedral group, in two independent implementations; the pattern-directed enumeration of Remark 5.11 and the annealing of Remark 5.12; the verification of Proposition 5.9; the port-block search over **nauty**; and the **graph6** lists of [12] together with the script that recomputes the degree profiles of §9. The saved output of every run quoted in the text is included.

Acknowledgements

First and above all, I thank Jesus Christ, the foundation of my life, who has changed my life and without whom none of this would have been possible.

I thank my family: my father, a man who taught me never to give up and to keep fighting; my mother, whose love and support lifted me up at the breaking point of my life; and my uncles, who stood by my family through the hardest moments.

I also thank Claude, an AI model developed by Anthropic, which assisted me extensively throughout this work: in the mathematical reasoning, in the design and execution of the long computer experiments, in programming, in checking the literature and in preparing the manuscript. Responsibility for the content rests entirely with the author.

All of this is for the glory of God.

For God so loved the world, that he gave his only begotten Son, that whosoever believeth in him should not perish, but have everlasting life.

— John 3:16

References

- [1] S. Abbasi and A. Jamshed, A degree constraint for uniquely Hamiltonian graphs, *Graphs and Combinatorics* 22 (2006), 433–442.
- [2] G. Brinkmann and M. De Pauw, Uniquely hamiltonian graphs for many sets of degrees, *Discrete Mathematics and Theoretical Computer Science* 26:3 (2024), #7; arXiv:2304.08946.
- [3] J. A. Bondy and B. Jackson, Vertices of small degree in uniquely Hamiltonian graphs, *Journal of Combinatorial Theory, Series B* 74 (1998), 265–275.
- [4] M. Bucić, E. Jahn, A. Pokrovskiy and B. Sudakov, 2-factors with k cycles in Hamiltonian graphs, *Journal of Combinatorial Theory, Series B* 144 (2020), 150–166.
- [5] A. Espuny Díaz, A. Girão, B. Granet and G. Kronenberg, On 2-factors of Hamiltonian graphs, preprint, arXiv:2605.11288 (2026).
- [6] R. C. Entringer and H. Swart, Spanning cycles of nearly cubic graphs, *Journal of Combinatorial Theory, Series B* 29 (1980), 303–309.

- [7] H. Fleischner, Uniquely Hamiltonian graphs of minimum degree 4, *Journal of Graph Theory* 75 (2014), 167–177.
- [8] P. Gregor, Perfect matchings extending on subcubes to Hamiltonian cycles of hypercubes, *Discrete Mathematics* 309 (2009), 1711–1713.
- [9] F. Wang and H. Zhang, Small matchings extend to Hamiltonian cycles in hypercubes, *Graphs and Combinatorics* 32 (2016), 363–376.
- [10] J. Goedgebeur, J. Jooker, O.-H. S. Lo, B. Seamone and C. T. Zamfirescu, Few Hamiltonian cycles in graphs with one or two vertex degrees, *Mathematics of Computation* (2024), doi:10.1090/mcom/3943; arXiv:2211.08105.
- [11] J. Goedgebeur, B. Meersman and C. T. Zamfirescu, Graphs with few Hamiltonian cycles, *Mathematics of Computation* 89 (2020), 965–991.
- [12] K. Coolsaet, S. D’hondt and J. Goedgebeur, House of Graphs 2.0: a database of interesting graphs and more, *Discrete Applied Mathematics* 325 (2023), 97–107. Lists of uniquely Hamiltonian graphs at <https://houseofgraphs.org/UHG>, consulted October 2026.
- [13] P. Haxell, B. Seamone and J. Verstraëte, Independent dominating sets and Hamiltonian cycles, *Journal of Graph Theory* 54 (2007), 233–244.
- [14] R. J. Kang and C. Legrand-Duchesne, Dirac’s theorem and the switch geometry of perfect matchings, preprint, arXiv:2604.17911 (2026).
- [15] P. Kleer, V. Patel and F. Stroh, Switch-based Markov chains for sampling Hamiltonian cycles in dense graphs, *Electronic Journal of Combinatorics* 27(4) (2020), #P4.29.
- [16] D. Labbate and F. Romaniello, An updated survey on 2-factors of regular graphs, preprint, arXiv:2408.04642 (2024).
- [17] J. Sheehan, The multiplicity of Hamiltonian circuits in a graph, in *Recent Advances in Graph Theory* (Proc. Second Czechoslovak Sympos., Prague, 1974), Academia, Prague (1975), 477–480.
- [18] A. G. Thomason, Hamiltonian cycles and uniquely edge colourable graphs, *Annals of Discrete Mathematics* 3 (1978), 259–268.
- [19] C. Thomassen, Independent dominating sets and a second Hamiltonian cycle in regular graphs, *Journal of Combinatorial Theory, Series B* 72 (1998), 104–109.
- [20] W. T. Tutte, On Hamiltonian circuits, *Journal of the London Mathematical Society* 21 (1946), 98–101.