Blood Glucose Target Ranges

Typical adult targets used in diabetes education. Your care team may set different goals based on age, pregnancy, hypoglycemia risk, or other conditions.

80–130	<180	90–150	<7%
BEFORE MEALS MG/DL	1–2 HR AFTER MEALS MG/DL	BEDTIME MG/DL	A1C GOAL (MANY ADULTS)

When to check & what "in range" often means

TIMING	TYPICAL TARGET	NOTES
Fasting / before meals	**80–130 mg/dL**	ADA general adult guidance
1–2 hours after starting a meal	**< 180 mg/dL**	Peak post-meal reading
Bedtime	**90–150 mg/dL**	Ask your clinician for your range
Before exercise	**100–250 mg/dL***	*Avoid intense exercise if high with ketones
During illness	**Check more often**	See Sick Day Rules chart

Low — treat now

Below 70 mg/dL HYPOGLYCEMIA

- Treat with 15g fast-acting carbs
- Recheck in 15 minutes
- Repeat if still under 70

High — take action

Above 250 mg/dL CHECK KETONES

- Drink water; follow care plan
- Test for ketones if instructed
- Call clinician if rising or ill

Personalize your numbers

Older adults, young children, people with frequent lows, or those who are pregnant often have different targets. Write your clinician's goals here:

Fasting _____ · After meals _____ · Bedtime _____ · A1C _____

Educational reference only — not medical advice. Follow the targets set by your healthcare provider. Based on commonly cited adult ADA education ranges.