



UNIVERSITÀ DEGLI STUDI DI MILANO

Structural Bioinformatics

Carlo Camilloni

Aims and Background

- **Have a general understanding of what you can do with computer modelling (what kind of problems can be addressed and how?)**
- **Have a general understanding of the theoretical concepts that are the foundation of the field (why things work?)**
- **Know what are the strengths and weaknesses of different methodologies (What is the right method for a given problem and what are the limitations?)**
- **Have the tools to read and understand the scientific literature of the field**



Program

- 1. Statistical Picture of Biomolecules: theoretical framework**
- 2. Structural Biology and Visualisation**
- 3. Classical Molecular Dynamics**
- 4. Enhanced Sampling Techniques**
- 5. Markov State Models**
- 6. Quantum Chemistry**
- 7. Simplified Models**
- 8. Structure Prediction and Molecular Docking**
- 9. Protein Design**

Topics will be both introduced theoretically and tried by yourself by following tutorials



Practicals

Install VMD if you will use
your own laptop

We are going to use COLAB:
colab.research.google.com



<https://www.ks.uiuc.edu/Research/vmd/>

To use them (essentially to save the
results of your work)
You need a google Drive account,
please do one



Course Material

- Slides
- Review papers provided
- Books (only if you really want to learn much more)



UNIVERSITÀ
DEGLI STUDI
DI MILANO



Time table

**Tuesday 14.30-17.30: Computer room at the Department of Biosciences,
if you have never been there, go before the next week to register an
account (if you want to use the local machines).**

Friday 13.30-15.30: B8 (Bioscience)



Exam

Produce a report of the lab experience, one of them will be discussed at the exam

Make a presentation (<15 min) and discuss a scientific paper from a list I will share at the end of the course. Additional questions will be asked on the techniques employed in the paper and presented in the lectures.





UNIVERSITÀ DEGLI STUDI DI MILANO

A mechano-statistical perspective on biomolecules in motion


Carlo Camilloni

Observing the microscopic world is hard:

**Long exposure,
good signal vs noise
ratio, but things
move...**



**Short exposure,
things don't move...
but**




UNIVERSITÀ
DEGLI STUDI
DI MILANO




Observing the microscopic world is hard:


- We do not have a light microscope, so we don't see directly the molecules, we need to interpret other kind of signals (shorter wavelength, electrons, magnetic fields, ...)
- We cannot do movies, so we cannot see molecules in motion, but we take picture with more or less long exposure times
- We do not look at a single molecule at a time but at many (~fraction of Avogadro number)



Light microscope cannot look at molecular spatial scales




Molecules motion can span many time scales




Spatio-temporal resolution

- Different experimental techniques can explore processes at different spatial and time resolution, there is a region that can only be covered by computational methods, only NMR can get close to it, but its results are hard to interpret



A stochastic picture for molecules in motion

Stochastic means randomly determined, events happen by chance, by observing many events one can learn **what is the probability of the different outcomes** that is one can learn their **probability distribution**. If some outcome is more likely than another one could think in terms of energy that that outcome is favourable in **energy**.



$$p(j) = 1/6 \text{ with } j=1 \text{ to } 6$$

Discrete

$$\sum_{j=1}^6 p(j) = 1$$


Normalised
When all possibilities are accounted for the probability is 100%



A stochastic picture for molecules in motion

If your dice is biased towards a given number, let's say 1, then your histogram will be higher at 1. In terms of energy you can think that the state 1 is energetically more favourable.


$$\sum_{j=1}^6 p(j) = 1 \quad \begin{array}{l} \text{Normalised} \\ \text{When all possibilities are accounted} \\ \text{for the probability is 100\%} \end{array}$$



A stochastic picture for molecules in motion

Stochastic events are estimated by histograms, where events are counted, these can deviate from the ideal expected behaviour due to random and systematic errors.

Absolute probability needs to have informations on all outcomes, relative probabilities are just frequency ratios:




$$p(i) = \text{count}(i)/(\sum_k \text{count}(k))$$
$$p(j)/p(i) = \text{count}(j)/\text{count}(i)$$



Ligand binding: a molecular example

Here energy and probability are intuitively linked, furthermore, events happen on some **time scale**, so one would like to find a relationship between probability, energy and time scales of events.




In ligand-receptor binding process we can picture the process, at least in some cases, as a probability for R and L to be unbound vs a probability to be bound (RL) and consider the time scale of the process in terms of rates of binding (on) and unbinding (off).

The k_{off} rate is the frequency of unbinding (s^{-1}) so that $k_{off}[RL]$ is the number of unbinding events per second.

The k_{on} represent instead two processes (1) collision and (2) complexation. So this is the collision frequency times the complexation probability, this means that binding depends on concentration, indeed k_{on} is measured in ($s^{-1} M^{-1}$). $k_{on}[R][L]$ is the number of binding events per second, while $k_{on}[L]$ is the frequency of binding for a given concentration of ligand (excess of ligand).



A stochastic picture: ligand binding



If we start from a solution of free [R] and [L] we start forming RL with rate k_{on} , once we have some [RL] this can also dissociated with rate k_{off} , after some time we will be at equilibrium that is that the concentration of [R], [L] and [RL] will not change anymore

$$K_d \equiv \frac{k_{off}}{k_{on}} = \frac{[R][L]}{[RL]} = C^0 \frac{p_{unbound}}{p_{bound}}$$

The **dissociation constant** is defined as the ratio between k_{off} and k_{on} , it is measured in (M), where lower values means stronger binding (i.e. lower concentrations of ligand are enough to bind)

$$C^0 = 1 \text{ M}$$

$$\begin{aligned}\Delta G_{\text{binding}} &= k_B T \ln \left(\frac{K_d}{C^0} \right) = k_B T \ln \left(\frac{k_{off}}{k_{on} C^0} \right) \\ &= k_B T \ln \left(\frac{p_{unbound}}{p_{bound}} \right)\end{aligned}$$

This is the binding free-energy that is the amount of work that can be extracted or that must be provided for the binding reaction to happen, and you see is related to the rates as well as to the probabilities



From experiments: Energy perspective: Isothermal Titration Calorimetry


Temperature is kept constant in some volume for:

- buffer alone
- buffer with protein and increased concentration of ligand


The amount of heat needed (or produced) to keep the temperature constant is measured

$$Q = [RL] \Delta H V$$

From the variation of Q and the concentration [LR] it is possible to recover ΔH and K_d



Binding is more complex than that not? Towards linking microscopic configurations to macroscopic processes



The macroscopic “bound” and “unbound” states can be realised by a very large number of microscopic configuration, of the receptor, of the ligand, but also of the solvent around them, if we could see them we should be then able to link them.

Free Energy

$$\Delta G = \Delta H - T\Delta S$$


We have seen that the free energy is the effective energy that gives the population of a state (e.g. bound/unbound). Macroscopically is the available work the system can perform, i.e. the maximum work you can extract from the system.

The enthalpy is the average energy of a state including the average fluctuations of the volume occupied at a given pressure. Macroscopically is the amount of heat given or absorbed by a reaction.

The entropy is the difference between the two and corresponds to the number of microscopic realisation of a macrostate (for example we can think of odds/even numbers in a dice and 1,3,5-2,4,6 being the corresponding microstates, in this case the entropy of the two macrostates is the same)



A stochastic picture: protein folding



Their ratio define the $K_D = k_u/k_f$. This can used to calculate the difference in free energy:


$$\Delta G_{Eq} = RT \ln K_D$$

Protein in high-conc of denaturant is mixed with buffer to a target final low concentration of denaturant

The k_u rate is the frequency of unfolding (s^{-1}) so that $k_u[P]$ is the number of unfolding events per second.

The k_f rate is the frequency of folding (s^{-1}) so that $k_f[P]$ is the number of folding events per second.

The two rates can be measured by changing the concentration of denaturants by a stopped-flow and monitoring fluorescence.



Protein in buffer is mixed with denaturant to a target final high concentration of denaturant

At high C all the protein unfold
 $k_{obs}=k_u$ at low C all the protein fold so
 $k_{obs}=k_f$


Energy perspective: Differential Scanning Calorimetry

Temperature is increased linearly in time:

- To buffer alone
- To buffer with protein

The amount of heat needed is different, by subtracting one gets the heat needed to increase the temperature of the protein.


Curve Cp vs temperature




From the integral of C_p and it is possible to derive the Enthalpy difference.



Single Molecule FRET: kinetics and equilibrium



Where the balance between the two major macro states is determined by temperature



At room temperature the folded state can be ~99% populated, essentially suggesting


$$\langle O \rangle = \sum_i p_i O_i \simeq O_f \approx ?O_{structure}$$

But one should not forget that a macro-state is still a collection of many micro-states

Continuous Probability Distributions

$$p_G(x) \propto e^{-\frac{(x - \langle x \rangle)^2}{2\sigma^2}}$$

Gaussian




Density, they are in units^{-1}

Even if we know only proportionality this can be enough to know make comparisons: the probability of $p(0)/p(1)$. This is clear if you think about histograms.

$$\int_{-\infty}^{+\infty} e^{-\frac{(x - \langle x \rangle)^2}{2\sigma^2}} dx = \sqrt{2\pi}\sigma$$
$$\rho_G(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x - \langle x \rangle)^2}{2\sigma^2}}$$

Probability distributions are normalised



To calculate a probability one multiples for small interval:




If you want a continuous probability distribution from a discrete sampling you still need to calculate the integral (i.e. the area)



Cumulative distribution function (CDF)

$$cdf(x) = \int_{-\infty}^x pdf(x')dx'$$

Is the amount of probability from the lower limit to x



CDF are easier to describe the data:

- up to -1 there is only 10% of the data
- Between -1 and 1 there the 80% of the data

Sampling is the action of estimating a probability distributions. One takes elements and make histograms.

In an experiment data are distributed following the unknown distribution and you just count them. How do you generate numbers from a probability distribution?

Using the CDF:

- take a random number y between 0 and 1
- take the x value corresponding to such y from the CDF

Probability distributions can be multimodal




Multidimensional distribution function: projection and correlation

Probability distributions can be more than 1D. For example the probability distribution of a molecule around a receptor may be studied in terms of (x,y,z) coordinates of the center-of-mass, in this case would be 3D.

$$\int \rho(x, y) dx dy = 1 \text{ Normalisation}$$

How can we show a multi dimensional distribution in less dimensions? By projecting it:

$$\rho(x) = \int \rho(x, y) dy \text{ Projection/Marginalisation}$$



$$\langle x \rangle = \int dx dy x \rho(x, y)$$


$$\langle y \rangle = \int dx dy y \rho(x, y)$$

$$\langle xy \rangle = \int dx dy x y \rho(x, y)$$

By looking at the data can you think of a projection that is more important to understand the data?



Multidimensional distribution function: correlation



$$\langle x \rangle = \int dx dy x \rho(x, y)$$

$$\langle y \rangle = \int dx dy y \rho(x, y)$$

$$\langle xy \rangle = \int dx dy xy \rho(x, y)$$

If two variables are independent than

$$\rho(x, y) = \rho(x)\rho(y) \quad \text{So}$$

$$\langle xy \rangle = \langle x \rangle \langle y \rangle$$

So we have correlation if

$$\langle xy \rangle - \langle x \rangle \langle y \rangle \neq 0$$

$$r = \frac{\langle xy \rangle - \langle x \rangle \langle y \rangle}{\sigma_x \sigma_y}$$



Statistical Mechanics: conformations and velocities

What is the connection between statistics and mechanics?

That is the Boltzmann equation: $pdf(x, v) \equiv \rho(x, v) \propto \exp\left[\frac{-E(x, v)}{k_B T}\right] = \exp\left[\frac{-U(x)}{k_B T}\right] \exp\left[\frac{-K(v)}{k_B T}\right]$

Let's look at this pdf in more detail:

$$pdf(x) \equiv \rho(x) \propto \exp\left[\frac{-U(x)}{k_B T}\right]$$
$$pdf(v) \equiv \rho(v) \propto \exp\left[\frac{-K(v)}{k_B T}\right] = \exp\left[\frac{-\frac{1}{2}mv^2}{k_B T}\right]$$

Conformations and velocities are independent. This means that we can study them separately. The distribution of the velocities does not affect the distribution of the configurations. Furthermore the distribution of the velocity is Gaussian, so it is easy to integrate it.

So for all practical purposes we can just ignore the velocity: $pdf(x) \equiv \rho(x) \propto \exp\left[\frac{-U(x)}{k_B T}\right]$

This gives a link between the energy of a conformation and the probability of observing it



Statistical Mechanics: Ergodicity

- 1. A single molecule (at constant temperature) will move through all the possible conformations at random due to the collisions with the solvent. Each conformation is gonna be populated accordingly to the energy and temperature (the interaction with environment).**
- 2. If at a given time one stops the movie of many many copies of the same system in the same condition the picture taken will display conformations according to the same statistics.**


The collection of conformations in both cases (that is the sampling) is usually called an ensemble of conformations. This ensemble is characterised first of all by the conditions in which it has been acquired (constant temperature, pressure or volume, constant number of particles, ecc)



States, Probabilities and Free-Energies

State: is a collection of configurations (microstate), ideally belonging to the same potential energy basin.

A simple 1D potential energy



Here we can visually define two states A and B for the two basins, e.g. all the conformations belonging to A and to B

$$p_A = \int_{V_A} \rho(x)dx \propto \int_{V_A} \exp[-U(x)/k_B T]dx$$

$$p_B = \int_{V_B} \rho(x)dx \propto \int_{V_B} \exp[-U(x)/k_B T]dx$$


The Free-Energy is the “effective” energy of a state, i.e. the energy that will give the same probability

$$\frac{p_A}{p_B} = \frac{\int_{V_A} \exp[-U(x)/k_B T]dx}{\int_{V_B} \exp[-U(x)/k_B T]dx} \equiv \frac{\exp(-F_A/k_B T)}{\exp(-F_B/k_B T)}$$

$$F_i \propto -k_B T \ln \left(\int_{V_i} \exp[-U(x)/k_B T]dx \right)$$



Summary: A stochastic picture of molecules in solution



Macrostate populations are associated with free energy differences:

$$p_i \propto \exp\left(-\frac{G_i}{RT}\right)$$

Exchange rates are associated with free energy barriers:

$$k_{ij} = k_0 \exp\left(-\frac{G_{ij}^\ddagger}{RT}\right)$$

Energy for these processes is provided by thermal energy, so essentially by collision with water molecules

An equilibrium bulk experiment ideally capture the contribution from all the states:

$$\langle O \rangle = \sum_i p_i O_i = \frac{\sum_i O_i \exp\left(\frac{-G_i}{RT}\right)}{\sum_i \exp\left(\frac{-G_i}{RT}\right)}$$

Microscopically:

$$G_i = -RT \ln \int \exp\left(-\frac{U(r)}{RT}\right) \delta(i - i(r)) dr$$



Ideally, what should a computational microscope do?

- Observe the time evolution of molecules at high spatial and time resolution
- Observe them for very long time scales
- Be aware of their energy
- Be able to set different experimental conditions (Temperature, Pressure, solution conditions)
- Be accurate
- Be interpretable, that is find suitable macrostates to compare with experiments
- ...

