

A Lean-verified offset construction for the Erdős–Straus equation

Abstract

We give a self-contained formulation of an offset construction for the Erdős–Straus equation

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}.$$

The construction packages the classical divisor split into a congruence condition indexed by a prime $p \equiv 3 \pmod{4}$. Its full positive-integer statement is formalised in Lean 4 with Mathlib: the development constructs the denominators by natural-number division, proves the divisions exact and the denominators positive, and concludes the equality in $\mathbb{Q}$. We also formalise three groups of explicit infinite families, including the progressions arising from $p = 3$ and $d_1 = 2, 5$. We make no claim toward resolving the Erdős–Straus conjecture and no priority claim for the underlying construction.

1 The divisor split

The Erdős–Straus conjecture asks whether, for every integer $n \geq 2$, there exist $x, y, z \in \mathbb{N}_{>0}$ satisfying

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}. \quad (1)$$

The conjecture remains open. We use the following classical algebraic reduction; see, for example, Mordell [1].

Fix $x \in \mathbb{N}_{>0}$ with $4x > n$, and reduce

$$\frac{4}{n} - \frac{1}{x} = \frac{4x - n}{nx} = \frac{a}{b}, \quad \gcd(a, b) = 1.$$

Multiplication and rearrangement give

$$(ay - b)(az - b) = b^2. \quad (2)$$

Lemma 1.1 (positive divisor split). *Let $n, x, a, b, d_1, d_2, y, z \in \mathbb{N}_{>0}$. Suppose*

$$a = 4x - n, \quad b = nx, \quad d_1 d_2 = b^2,$$

and

$$ay = b + d_1, \quad az = b + d_2.$$

Then (x, y, z) satisfies (1).

Proof. The last three equations are precisely the factorisation (2). Expanding it and using $a = 4x - n$ and $b = nx$ gives

$$4xyz = n(xy + xz + yz).$$

All four variables n, x, y, z are nonzero, so division by $nxyz$ gives (1). □

The Lean development separates these two steps. It first proves the cross-multiplied identity over $\mathbb{Z}$, then proves that a positive instance over $\mathbb{N}$ gives the rational equality in $\mathbb{Q}$.

2 A positive certificate

The offset form avoids subtraction and is convenient both mathematically and formally.

Theorem 2.1 (positive offset certificate). *Let $n, x, p, b, d_1, d_2, y, z \in \mathbb{N}$, with $n, x, p, y, z > 0$. If*

$$4x = n + p, \quad b = nx, \quad (3)$$

$$d_1 d_2 = b^2, \quad py = b + d_1, \quad pz = b + d_2, \quad (4)$$

then (x, y, z) satisfies (1).

Proof. The equations are the positive divisor split with $a = p = 4x - n$. Equivalently, substitution in the cross-multiplied target leaves the multiple

$$p(d_1 d_2 - b^2),$$

which vanishes by (4). Positivity then permits passage to the rational identity. $\square$

In Lean, `IsES n x y z` contains both the four positivity assertions and the equality (1) in $\mathbb{Q}$. Theorem 2.1 is the declaration `isES_of_offset_certificate`.

3 The constructive offset theorem

Theorem 3.1 (constructive offset theorem). *Let $n, p, d_1 \in \mathbb{N}$ satisfy*

$$n \geq 2, \quad p \text{ prime}, \quad p \equiv 3 \pmod{4}, \quad n \equiv 1 \pmod{4}, \quad p \nmid n.$$

Set

$$x = \frac{n+p}{4}, \quad b = nx.$$

Suppose $d_1 > 0$, $d_1 \mid b$, and

$$p \mid b + d_1, \quad (5)$$

equivalently $d_1 \equiv -b \pmod{p}$. Define

$$d_2 = \frac{b^2}{d_1}, \quad y = \frac{b + d_1}{p}, \quad z = \frac{b + d_2}{p}. \quad (6)$$

Then $x, y, z \in \mathbb{N}_{>0}$, all divisions in (6) are exact, and (x, y, z) satisfies (1).

Proof. The congruences on n and p show that $4 \mid n + p$, so $x \in \mathbb{N}$ and $4x = n + p$. Moreover $x > 0$. If $p \mid x$, then $p \mid 4x = n + p$, hence $p \mid n$, a contradiction. Thus $p \nmid b = nx$.

Since $d_1 \mid b$, the quotient $d_2 = b^2/d_1$ is exact and positive. It remains to prove $p \mid b + d_2$. Work in the field $\mathbb{Z}/p\mathbb{Z}$. Condition (5) gives

$$d_1 = -b.$$

The element b is nonzero, hence so is d_1 . From $d_1 d_2 = b^2$, we may cancel d_1 to obtain

$$d_2 = -b,$$

and therefore $p \mid b + d_2$. Both definitions of y, z in (6) are consequently exact. Their numerators are positive, so $y, z > 0$. Theorem 2.1 now applies. $\square$

Remark. The primality assumption is used only for cancellation modulo p . The Lean proof implements this step in `ZMod p`; the remainder of the construction uses natural numbers, with the final equality stated over the rationals.

4 Explicit infinite families

We record the families formalised in the accompanying Lean file. In every corollary, $k \in \mathbb{N}$.

Corollary 4.1 (Family I). *Let*

$$n = 12k + 9, \quad x = 3k + 3, \quad b = n(k + 1).$$

Then

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{b+1} + \frac{1}{b(b+1)}.$$

Proof. Here $4x - n = 3$ and $\gcd(3, nx) = 3$, so the reduced numerator is 1. Taking $d_1 = 1$ and $d_2 = b^2$ in the divisor split gives the displayed denominators. The Lean proof is a direct positive polynomial identity. $\square$

Corollary 4.2 ($p = 3, d_1 = 2$). *Let*

$$n = 24k + 5 \quad \text{or} \quad n = 24k + 13.$$

Put $x = (n + 3)/4$ and $b = nx$. Then

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{(b+2)/3} + \frac{1}{(b+b^2/2)/3}.$$

Proof. For both progressions, $n \equiv 1 \pmod{4}$, $3 \nmid n$, $2 \mid b$, and $3 \mid b + 2$. Theorem 3.1 applies with $p = 3$ and $d_1 = 2$. $\square$

Corollary 4.3 ($p = 3, d_1 = 5$). *Let*

$$n \in \{60k + 5, 60k + 17, 60k + 25, 60k + 37\}.$$

Put $x = (n + 3)/4$ and $b = nx$. Then

$$\frac{4}{n} = \frac{1}{x} + \frac{1}{(b+5)/3} + \frac{1}{(b+b^2/5)/3}.$$

Proof. Each progression satisfies $n \equiv 1 \pmod{4}$ and $3 \nmid n$. Direct factorisation of either n or $x = (n + 3)/4$ gives $5 \mid b$, while $3 \mid b + 5$. Apply Theorem 3.1 with $p = 3$ and $d_1 = 5$. $\square$

The two progressions in Corollary 4.2 are the nonmultiples of 3 in the classical Mordell region $n \equiv 5 \pmod{8}$; the remaining progression $n = 24k + 21$ is contained in Family I. The purpose of including them is to exhibit the general construction and its formal specialisation, not to claim a new residue family.

5 Lean formalisation

The formalisation is contained in `lean/ESTheorem.lean` and uses Lean 4 with Mathlib. Its public positive interface consists of:

- `IsES` and `isES_of_polynomial`;
- `isES_of_offset_certificate`;
- `second_divisibility`;
- `constructive_offset`;

- `family_I_positive`, `family_d2_24k5`, `family_d2_24k13`, and the four `family_d5_60k` declarations.

For every explicit progression, the file also checks the instances $k = 0$ and $k = 1$. The project builds with no `sorry`. The theorem statement includes the positive denominators and the rational unit-fraction equality; the formalisation is therefore not merely a check of a polynomial identity.

6 Scope

The divisor split is classical, and offset choices of x belong to the standard toolkit for Egyptian-fraction problems. The contribution of this note is a compact theorem interface, an end-to-end Lean proof of the positive construction, and a small collection of formally checked infinite specialisations. No computational coverage, density statement, or progress toward the full Erdős–Straus conjecture is asserted.

References

- [1] L. J. Mordell, *Diophantine Equations*, Academic Press, 1969.
- [2] R. C. Vaughan, *On a problem of Erdős, Straus and Schinzel*, J. London Math. Soc. (2) 2 (1970), 391–396.
- [3] The Mathlib Community, *Mathlib: the mathematical library of Lean 4*, [project repository](#).