

An inequality between determinant and signature for L -space knots

Machine-generated conjecture, its checks, a partial proof, and the open case

2026-09-19, revised 2026-10-10

Abstract

We consider the inequality $\det(K) \leq 1 + |\sigma(K)|$ for L -space knots. It was produced by a program that fits linear inequalities to the KnotInfo database and discards what its automatic checks refute. It survived those checks and the computations of Section 3. We prove it for every L -space knot that is an iterated torus knot, in particular for all algebraic knots. The hyperbolic case is open, and it is where the inequality is sharp: equality holds on the Baker–Kegel knots K_k , an infinite family of hyperbolic L -space knots, for which we prove $\det(K_k) = 4k + 5$ and compute $\sigma(K_k) = -(4k + 4)$ for $k \leq 60$. Each K_k has the Alexander polynomial, hence the knot Floer complex, of an L -space cable that attains equality as well. Among the 632 census L -space knots equality holds only on K_1 . Sections 2 and 3 give the provenance and the computations, so that the proof in Sections 5 and 6 can be checked without our files; the one lemma we prove from scratch rather than cite is additionally formalised in Lean 4 with no `sorry`.

1 Statement and conventions

Signs follow KnotInfo: the right-handed trefoil has $\sigma = -2$, $s = 2$, $\tau = 1$. A knot $K \subset S^3$ is an *L -space knot* if some positive Dehn surgery on K yields an L -space. Positive torus knots then have $\sigma < 0$. We do not assume a sign for a general L -space knot: $\sigma \leq 0$ is carried through the induction of Theorem 1 for the knots treated there, and it held on every L -space knot we computed (all 632 census knots and all 68 knots of the families of Section 3). Write $\det(K) = |\Delta_K(-1)|$ and $g = g(K)$ for the Seifert genus.

Conjecture 1 (K33). *For every L -space knot K ,*

$$\det(K) \leq 1 + |\sigma(K)|.$$

We use the following standard facts. An L -space knot is fibered, strongly quasipositive, and satisfies $g = g_4 = \tau$ (Ozsváth–Szabó, Ni, Hedden); its Alexander polynomial has the form $\Delta_K(t) = \sum_{k=0}^{2n} (-1)^k t^{a_k}$ with $a_0 = g > a_1 > \dots > a_{2n} = -g$ (Ozsváth–Szabó). Consequently $|\sigma(K)| \leq 2g$ and $\det(K) \leq 2g + 1$. The second is the determinant–genus inequality for L -space knots [18, Lemma 5], [3, §3.1]; Conjecture 1 would sharpen it.

2 Provenance

The inequality was not conjectured by a person. It is candidate K33 of 42 statements emitted by `ragusa`, an engine that enumerates bounds of the form (property) $\Rightarrow$ (target $\leq c \cdot$ feature $+k$) over a table of invariants, here KnotInfo [14] restricted to prime knots with 3 to 13 crossings, and reports

what survives its filters together with the evidence behind each survivor. K33 was found on the `l_space` column, where the table contains 11 knots; those knots were the whole of its initial evidence.

The other 41 candidates were triaged in the same session: 18 were refuted by explicit knots, 13 turned out to be known theorems or true by definition, 3 had one clause refuted and one left open, and 6 remain open but rest on weak evidence. Two remained open with evidence beyond the table: K33, and $u = g$ for L -space knots.

3 Verification history

All computations use SnapPy (knot Floer homology, for the L -space property and τ), Sage (Seifert matrices, signatures, determinants) and exact formula implementations; scripts are listed in Section 9.

family of L -space knots	tested	viol.	how the invariants are known
iterated torus knots	11728	0	exact: cabled Alexander polynomial, Litherland's formula
SnapPy census L -space knots (all)	632	0	Seifert matrix of a positive braid word
census knots, independent route	271	0	knot Floer homology, Sage signature
Baker–Kegel, Himeno, $(2, q)$ -cables	68	0	Seifert matrix of the braid word
Baker–Kegel K_k , $k \leq 60$	60	0	Seifert matrix of the braid word, certified
$H(n, m)$ (Section 7), outside the census	9	0	Seifert matrix of the braid word, certified
one-bridge braids $B(w, b, t)$, $w \leq 15$ (parameter triples)	274	0	Seifert matrix of the braid word
twisted torus knots $K(p, q; r, s)$ (distinct knots)	87	0	knot Floer homology, then exact Seifert signature
random braid words (distinct knots)	46	0	knot Floer homology, then exact Seifert signature

The census braid words are those published by Baker and Kegel (arXiv:2203.12013, Appendix), which cover all 632 census L -space knots. Every closure was checked to be the named census knot: 631 of 632 by SnapPy's isometry test directly, the remaining one after retriangulation and at higher precision. All 632 signatures are exact, computed by congruence diagonalisation over $\mathbb{Q}$ from the integral matrix $V + V^T$; they agree with an earlier floating-point computation on all 632 knots and with Sage's own exact signature on the 511 knots where that finished. The one-bridge braids are L -space knots by [20]; for the 40 smallest this was also confirmed directly by knot Floer homology, with the genus matching the braid-surface genus.

Generated families, counted by knot. Three rows of the table come from parameters or words rather than from a list: the one-bridge braids, the twisted torus knots $K(p, q; r, s)$, the closures of $(\sigma_1 \cdots \sigma_{p-1})^q (\sigma_1 \cdots \sigma_{r-1})^s$, and the closures of random braid words. Different parameters often close to the same knot, and an earlier version of this note counted diagrams, which overstated these families several times over. Counted by knot type, exactly for hyperbolic knots through the isometry signature of the complement and approximately otherwise, the 274 one-bridge parameter triples give 195 knots, 149 of them hyperbolic and at least 37 of those outside the SnapPy census; the 549 twisted torus records give 87 knots, 42 hyperbolic; and the 14452 random records give 46 knots, 17 hyperbolic. Every hyperbolic knot in the last two families is a census knot, so they confirm the census rows

rather than extend them; the hyperbolic evidence beyond the census is the one-bridge braids and the Baker–Kegel and Himeno families. In the twisted torus and random families the L -space property was decided by each knot’s own knot Floer homology, and the determinant was computed twice, from the knot Floer gradings and from the Seifert matrix, agreeing on all 15001 records.

Independent check. A language model with no access to our computations re-derived the lattice count and recomputed, independently: the six torus signatures used here, Lemma 1 and the bound on $N_<$ for all coprime $p \leq 24$, $q < 120$, the closed forms and their reductions, the list of leftover small pairs in the induction, the induction step under worst-case hypotheses, and the identity (1) on three examples. It found no errors. The gaps it flagged (an unsupported sign assertion, boundary points in the count, the $p < q$ convention, the nontriviality hypothesis in (F4), the derivation of the symmetry of S , missing references) are addressed in this version. A third model corrected the proof of Lemma 1, where an earlier version claimed an equivalence that is only a sufficient condition, and asked for the hypotheses and chirality conventions now stated; the identity $\sigma = 4N_< - 2g$ was verified on all coprime $p \leq 15$, $q < 50$, where $N_</g \leq 0.234$, so the branch proved here is the operative one. A second model, asked to attack the open case, contributed the congruence of Section 5, the one-bridge braid family of the table, and the rigidity observation of Section 7; its computations of the one-bridge braids (274 parameter triples, 0 violations, 70 equalities) and of the Baker–Kegel invariants were reproduced independently here.

The hypothesis is not redundant. Evaluating the same inequality on census knots whose braid word is positive or negative but which are *not* L -space knots produces 25 violations. So the L -space condition does the work; this is unlike several of the 42 candidates, whose hypotheses turned out to be irrelevant.

Sharpness. Equality $\det = 1 + |\sigma|$ holds for $T(2, n)$, for the $(2, q)$ -cables of the trefoil, and, among the families tested beyond the census, for 62 of 68 knots, including every Baker–Kegel K_n computed there, $n \leq 25$ (Section 7 extends this to $n \leq 60$). These are hyperbolic: K_1 is the census knot o9_30634, the one census L -space knot that is not braid positive, and K_n for $n \geq 2$ is not known to be braid positive. Of the 632 census L -space knots, exactly 1 attains equality, and it is K_1 .

4 A reformulation

For an L -space knot let $S \subset \mathbb{Z}_{\geq 0}$ be its staircase set (the *formal semigroup*; for an algebraic knot it is the semigroup of the singularity, but for a general L -space knot it need not be closed under addition), the complement of the g gaps, determined by $t^g \Delta_K(t) = (1 - t) \sum_{s \in S, s < 2g} t^s + t^{2g}$. Evaluating at $t = -1$ and using the symmetry $s \leftrightarrow 2g - 1 - s$ between $S \cap [0, 2g)$ and the gaps, which follows from $\Delta_K(t) = \Delta_K(t^{-1})$ and gives $|S \cap [0, 2g)| = g$ as well, one obtains

$$\det(K) = |2D + 1|, \quad D = \#\{\text{odd gaps}\} - \#\{\text{even gaps}\}. \quad (1)$$

Hence Conjecture 1 is equivalent to $|\sigma| \geq 2D$ when $D \geq 0$, and to $|\sigma| \geq 2|D| - 2$ when $D < 0$. Identity (1) was checked on 11728 iterated torus knots with 0 mismatches.

Remark 1. The cruder route fails. Since the coefficients of Δ_K are ± 1 , $\det(K)$ is at most the number of nonzero coefficients, so Conjecture 1 would follow from $|\sigma| \geq \#\{\text{terms}\} - 1$. That inequality is false: it fails on 401 of the 11728 iterated torus knots tested, and the failures are exactly the knots where Conjecture 1 is sharp. Any proof must therefore see the cancellation in $\Delta_K(-1)$.

5 An arithmetic refinement

For any knot, $\Delta_K(-1) \equiv 1 \pmod{4}$ and $\text{sign } \Delta_K(-1) = (-1)^{\sigma/2}$ (Murasugi), so $\det(K) \equiv (-1)^{\sigma/2} \pmod{4}$ and therefore

$$1 + |\sigma(K)| - \det(K) \equiv 0 \pmod{4}. \quad (2)$$

Checked on all 1523 L -space knots computed here with an exact signature and determinant: 0 failures. Two consequences. First, a counterexample to Conjecture 1 must satisfy $\det(K) \geq |\sigma(K)| + 5$; the inequality cannot fail narrowly. Second, combining (2) with (1), the *signature defect* $2g - |\sigma|$ of an L -space knot satisfies $2g - |\sigma| \equiv 0 \pmod{4}$ when $D \geq 0$, and Conjecture 1 becomes

$$2g - |\sigma| \leq 4 \# \{\text{even gaps}\} \quad (D \geq 0),$$

both sides being multiples of 4. (For $D < 0$ the corresponding statement is $2g - |\sigma| \leq 4 \# \{\text{odd gaps}\} + 2$.) In particular the conjecture holds automatically for every L -space knot with $\det \leq 3$ and for every definite one.

6 Proof for iterated torus knots

Throughout, $p \geq 2$ and q are coprime, $T(p, q)$ is the positive torus knot with $g(T(p, q)) = (p-1)(q-1)/2$, and $K_{p,q}$ denotes the (p, q) -cable of K (winding number p). We use:

- (F1) $\Delta_{K_{p,q}}(t) = \Delta_K(t^p) \Delta_{T(p,q)}(t)$; since $\Delta_K(1) = 1$ this gives $\det(K_{p,q}) = \det(K) \det(T(p, q))$ for p odd and $\det(K_{p,q}) = \det(T(p, q))$ for p even.
- (F2) Litherland's cabling formula $\sigma(K_{p,q}) = \sigma_{(-1)^p}(K) + \sigma(T(p, q))$, so $\sigma(K_{p,q}) = \sigma(K) + \sigma(T(p, q))$ for p odd and $= \sigma(T(p, q))$ for p even.
- (F3) $\det(T(p, q)) = 1$ if p, q are both odd, $= p$ if q is even, $= q$ if p is even.
- (F4) (Hedden; Hom) for K nontrivial, $K_{p,q}$ is an L -space knot if and only if K is one and $q \geq p(2g(K) - 1)$.

(F1)–(F3) were verified on 4088 cable instances, with 0, 0 and 0 mismatches respectively.

Lemma 1. *For integers $2 \leq p < q$ with $\gcd(p, q) = 1$, $\sigma(T(p, q)) \leq -g(T(p, q))$, where $g(T(p, q)) = \frac{(p-1)(q-1)}{2}$; in particular $|\sigma| \geq g$.*

Proof. We use the lattice count for torus knot signatures (Brieskorn; Gordon–Litherland–Murasugi [13]; Litherland [24]), in the normalisation $\sigma(T(2, 3)) = -2$ of Section 1: $\sigma(T(p, q)) = N_{\text{out}} - N_{\text{in}}$ over the grid $G = \{1, \dots, p-1\} \times \{1, \dots, q-1\}$, where (i, j) is counted in N_{in} when $\frac{1}{2} < \frac{i}{p} + \frac{j}{q} < \frac{3}{2}$. The involution $(i, j) \mapsto (p-i, q-j)$ of G exchanges the two outside regions, so $N_{\text{out}} = 2N_{<} with $N_{<} = \#\{(i, j) \in G : \frac{i}{p} + \frac{j}{q} < \frac{1}{2}\}$. No lattice point of G lies on a boundary: $\frac{i}{p} + \frac{j}{q} \in \{\frac{1}{2}, \frac{3}{2}\}$ gives $2(iq + jp) \in \{pq, 3pq\}$, so $p \mid 2i$ and $q \mid 2j$, forcing $2i = p$ and $2j = q$ with p and q both even, contradicting $\gcd(p, q) = 1$. As $N_{\text{in}} + N_{\text{out}} = 2g$, the count reads $\sigma(T(p, q)) = 4N_{<} - 2g$, so $|\sigma| \geq g$ holds if $N_{<} \leq g/4$, and also if $N_{<} \geq 3g/4$. We prove the first branch, $N_{<} \leq (p-1)(q-1)/8$, which gives $\sigma(T(p, q)) \leq -g < 0$.$

For fixed i the number of admissible j is $\lceil f(i) \rceil - 1 \leq f(i)$ with $f(i) = \frac{q}{2} - \frac{iq}{p}$, and $f(i) > 0$ exactly for $i < p/2$. Summing over those i : if p is even, with $m = p/2 - 1$, one gets $N_{<} \leq q(p-2)/8$, and $q(p-2) \leq (p-1)(q-1)$ reduces to $q \geq p-1$; if p is odd, with $m = (p-1)/2$, one gets $N_{<} \leq q(p-1)^2/(8p)$, and $q(p-1)/p \leq q-1$ reduces to $p \leq q$. Both hold because $q > p$. $\square$

The lattice inequality is machine-checked. The combinatorial core of this proof, the bound $N_{<} \leq (p-1)(q-1)/8$ for all integers $2 \leq p < q$, has been formalised in Lean 4 against Mathlib and carries no sorry: `#print axioms` reports only `propext`, `Classical.choice` and `Quot.sound`. Coprimality is not needed for it. The formalisation covers the column decomposition of the grid count, the per-column bound, the cut at $(p-1)/2$ in both parities, the Gauss sum, the closed forms, and the two final comparisons; it does not cover the topology the lemma is used with, namely the lattice count for $\sigma(T(p, q))$, which is cited. So the part of Lemma 1 proved here from scratch, rather than cited, is the part that is now checked by a machine. The file is `K33Lattice/Basic.lean`.

Every step of this proof was also checked numerically on 429 pairs with $p \leq 15$, $q < 60$ (0 failures), and the conclusion holds on all 2166 pairs with $p \leq 20$, $q < 200$. Lemma 1 says $-\sigma \geq b_1/2$ for torus knots, which is stronger than the linear bounds known for arbitrary positive braid links (Feller proved a linear bound; Greene–Liechti give $-\sigma \geq b_1/4$, see the abstract of [12]). We did not find Lemma 1 in the literature; it may be folklore.

Theorem 1. *Every L -space knot which is an iterated torus knot satisfies $\det(K) \leq 1 + |\sigma(K)|$. In particular this holds for all positive torus knots, hence for their mirrors since $\det$ and $|\sigma|$ are mirror invariant, and for all algebraic knots, which are iterated torus knots with positive parameters [8] and hence L -space knots.*

Proof. The induction is on the cabling depth, with torus knots at depth one; by (F4) applied to the companion (Hom’s “only if”), every companion of an L -space iterated torus knot is again an L -space knot, so the hypothesis is available at each step. We prove simultaneously that $\det(K) \leq 1 + |\sigma(K)|$ and $\sigma(K) < 0$. For torus knots we may assume $p < q$, since $T(p, q) = T(q, p)$; in a cable $K_{p,q}$ of a nontrivial L -space knot, (F4) gives $q \geq p(2g_K - 1) \geq p$, and $\gcd(p, q) = 1$ excludes $q = p$.

Base case. For $T(p, q)$, $\sigma < 0$ is part of Lemma 1, and: if p, q are both odd then $\det = 1$. If q is even then $\det = p$ and Lemma 1 gives $|\sigma| \geq 3(p-1)/2$ (using $q \geq 4$), whence $1 + |\sigma| \geq p$. If p is even then $\det = q$; for $p = 2$ we have $|\sigma(T(2, q))| = q - 1$, so $1 + |\sigma| = q$, with equality; for $p \geq 4$, Lemma 1 gives $|\sigma| \geq 3(q-1)/2 \geq q - 1$.

Induction step. Let K be an L -space knot with $\det(K) \leq 1 + A$, where $A = |\sigma(K)| \leq 2g_K$, and let $K_{p,q}$ be an L -space cable, so $q \geq p(2g_K - 1)$ by (F4). Put $B = |\sigma(T(p, q))|$. By the induction hypothesis $\sigma(K) < 0$, and $\sigma(T(p, q)) < 0$ by Lemma 1, so (F2) gives $\sigma(K_{p,q}) < 0$ and $|\sigma(K_{p,q})| = A + B$ for p odd, $= B$ for p even; this also propagates the sign claim.

If p is even, then $\det(K_{p,q}) = \det(T(p, q)) = q$ by (F1) and (F3), and $q \leq 1 + B$ was shown in the base case.

If p and q are both odd, then $\det(K_{p,q}) = \det(K) \leq 1 + A \leq 1 + A + B$.

If p is odd and q even, then $\det(K_{p,q}) = p \det(K) \leq p(1 + A)$, and the claim $p(1 + A) \leq 1 + A + B$ is $(p-1)(1 + A) \leq B$. By Lemma 1 it suffices that $1 + A \leq (q-1)/2$, i.e. $q \geq 2A + 3$, and since $A \leq 2g_K$ it suffices that $q \geq 4g_K + 3$. As $p \geq 3$, (F4) gives $q \geq 6g_K - 3$, which is $\geq 4g_K + 3$ exactly when $g_K \geq 3$. For $g_K \in \{1, 2\}$ the uncovered pairs are finite: for $g_K = 1$ they are $(p, q) = (3, 4)$ and $(5, 6)$, where $B = 6 \geq 6$ and $B = 16 \geq 12$; for $g_K = 2$ it is $(3, 10)$, where $B = 14 \geq 10$. $\square$

The induction step was also verified directly, with the worst admissible values $A = 2g_K$ and $\det(K) = 1 + A$, over $g_K \leq 11$, $p \leq 11$ and the full range of q allowed by (F4): 0 failures.

7 The open case

Theorem 1 does not reach the hyperbolic L -space knots, and that is where the inequality is sharp. Baker and Kegel (arXiv:2203.12013) construct hyperbolic L -space knots K_n , the closures of the braid words

$$[(2, 1, 3, 2)^{2n+1}, -1, 2, 1, 1, 2],$$

which are not braid positive for $n = 1$ and are not known to be braid positive for $n \geq 2$; Himeno (arXiv:2506.22934) constructs infinitely many hyperbolic L -space knots that are provably not braid positive. Equality $\det = 1 + |\sigma|$ holds for every Baker–Kegel K_n tested in Section 3 ($n \leq 25$), and more is true.

Proposition 1 (Baker–Kegel). *For every $k \geq 0$, $\Delta_{K_k} = \Delta_{C_k}$, where C_k is the $(2, 4k + 5)$ -cable of $T(2, 2k + 1)$, and $\det(K_k) = 4k + 5$.*

Baker and Kegel compute Δ_{K_k} [5, proof of Theorem 4.4] and note that a satellite with this polynomial would have to be C_k [5, §2, second proof of Lemma 2.3]; the determinant follows. We rechecked both independently.

Proof. Let ρ be the reduced Burau representation of B_4 , $P = \sigma_2\sigma_1\sigma_3\sigma_2$ and W the tail of the word, so that $\Delta_{K_k} \doteq \det(I - \rho(P)^{2k+1}\rho(W))(1 - t)/(1 - t^4)$. Expanding the 3×3 determinant, the right side is a sum of traces of powers of $A = \rho(P)^2$, of its second exterior power and of $\det A$, so by Cayley–Hamilton it satisfies a monic linear recurrence in k of order 8 over $\mathbb{Q}(t)$. So does $\Delta_{C_k}(t) = \Delta_{T(2, 2k+1)}(t^2) \Delta_{T(2, 4k+5)}(t)$, with characteristic roots $1, t^4, t^8$; the difference satisfies a recurrence of order 11 and vanishes for $k = 0, \dots, 10$, hence for all k . At $t = -1$ the order-8 recurrence becomes $(x-1)^8$, which $4k+5$ also satisfies, and the two agree for $k = 0, \dots, 7$. `bk_alex.sage` and `bk_det.sage` carry out these computations exactly. $\square$

Both K_k and C_k are L -space knots (C_k by (F4)), so they have the same knot Floer complex. C_k is an iterated torus knot and attains equality in Theorem 1: $\det(C_k) = 4k + 5$ and $\sigma(C_k) = \sigma(T(2, 4k + 5)) = -(4k + 4)$ by (F1)–(F3). For K_k we computed $\sigma(K_k) = -(4k + 4)$ for $k \leq 60$, from the Seifert matrix of the braid word (sizes up to 486, every eigenvalue certified away from zero), in agreement with [5, Remark 4.6], which states $|\sigma(K_k)| = g + 2$. On this family the hyperbolic knot and the cable share the Floer complex, the determinant and the signature.

Question 1. *Does $\det(K) \leq 1 + |\sigma(K)|$ hold for every L -space knot, in particular for the hyperbolic ones? Equivalently, by (1), is $|\sigma(K)| \geq 2(\#\{\text{odd gaps}\} - \#\{\text{even gaps}\})$?*

The sharp case carries a rigidity statement. If all g gaps are odd then $S \cap [0, 2g)$ consists of the even numbers, so $\Delta_K = \Delta_{T(2, 2g+1)}$, $D = g$ and $\det = 2g + 1$; Conjecture 1 then forces $|\sigma(K)| = 2g$, that is, the symmetrised Seifert form is definite. So the conjecture implies: every L -space knot with the Alexander polynomial of $T(2, 2g + 1)$ has $\sigma = -2g$. This is a weak form of Problem 1.21(c)(i) of the K3 problem list [21] (knot Floer homology detects $T(2g + 1, 2)$). Restricted to L -space knots, that problem is the conjecture that the only thin L -space knots are the $T(2, n)$ [7, §1]. It is proved for $g = 1$ by Ghiggini [11], for $g = 2$ by Farber–Reinoso–Wang [10] together with Baldwin–Hu–Sivek [2], and for no larger g [1, §1.1]. Any proof of Conjecture 1 therefore proves this definiteness statement.

What is known about the sharp case. Three further facts, contributed by a fourth reader and checked here against the exact data. (i) Such a K is not a nontrivial cable: the staircase set of a cable $J_{p,q}$ is $\{ps + qk : s \in S_J, 0 \leq k \leq p - 1\}$, whose smallest positive element is $\min(p s_{\min}, q) \geq 4$, whereas $S_{T(2, 2g+1)}$ has smallest positive element 2; here $s_{\min} \geq 2$ because 1 is a gap of every nontrivial L -space knot, which holds on all 11728 iterated torus knots computed. Among those knots, the only ones with $\Delta = \Delta_{T(2, 2g+1)}$ are the $T(2, 2g + 1)$ themselves. Baldwin and Sivek prove more: an L -space knot with $\Delta = \Delta_{T(2, 2g+1)}$ is not a satellite [1, Proposition 6.8]. (ii) The roots of $\Delta_{T(2, 2g+1)}$ are simple and on the unit circle, so the Levine–Tristram function has g jumps of ± 2 on the upper half circle and $\sigma(K) = -2g + 4m$, where m counts the positive jumps; the question is exactly $m = 0$, that is, that the signature function of K agrees with that of $T(2, 2g + 1)$. (iii) Definiteness does not by

itself force the torus knot: among the 11889 L -space iterated torus knots computed, exactly 21 are definite, namely the $T(2, n)$ together with $T(3, 4)$ and $T(3, 5)$ (the knots among the ADE singularity links). Neither of the last two has the Alexander polynomial of a $T(2, 2g + 1)$, so on this restricted class definiteness and detection still coincide.

The hypothesis of the sharp case is met only by $T(2, 2g + 1)$. For an L -space knot, $\det = 2g + 1$ exactly when every gap is odd, that is when $\Delta = \Delta_{T(2, 2g + 1)}$, which is also exactly when the knot is thin. Across the census, the families beyond it, the one-bridge braids, the twisted torus knots and the random search, reaching genus 145 in all, 2009 records meet this hypothesis. Each was put into braid form and found, individually, to be the closure of σ_1^{2g+1} in B_2 , which identifies it as $T(2, 2g + 1)$; together they cover $g \leq 11$, and none of the census knots is among them. So every thin L -space knot found here is a $T(2, 2g + 1)$, which is the L -space case of Problem 1.21(c)(i) appearing in the data, and on all of them the sharp case of Conjecture 1 holds.

Together these narrow a counterexample to Conjecture 1 in the sharp case: it would have genus $g \geq 3$, it would be hyperbolic (a torus knot with this Alexander polynomial is $T(2, 2g + 1)$, and satellites are excluded by [1, Proposition 6.8]), and it would need a positive Levine–Tristram jump at one of the g roots. The same hyperbolic case is the one left open in [1, Theorem 1.6], on detecting $T(2, 2g + 1)$ by its traces.

Why the induction stops. For a torus knot the gaps of $\langle p, q \rangle$ satisfy $2g - |\sigma| = 4 \#\{\text{gaps} > pq/2\}$, so Theorem 1 for torus knots says $\#\{\text{gaps} > pq/2\} \leq \#\{\text{even gaps}\}$. No analogue of the half-conductor $pq/2$ is available for a hyperbolic L -space knot, and that is exactly what the cabling induction replaces.

Where equality occurs. Among the hyperbolic L -space knots tested, equality occurs exactly on the Baker–Kegel family: on every K_n tested, and on no other census knot (Himeno’s K_2 is the same knot as K_1). Failing to be braid positive does not force it: Himeno’s K_n with n even provably are not braid positive, and none of $K_3, \dots, K_8$ attains equality, the last having $\det = 39$ against $1 + |\sigma| = 99$. Proposition 1 locates the equality: each K_k has the knot Floer complex and the signature of a cable that attains it. (None of the 70 one-bridge parameter triples with equality closes to a hyperbolic knot.)

The family $H(n, m)$. Both families are members of a two-parameter family of closed $2n$ -braids, $H(n, m)$, the closure of $X_n^m T_n$ with X_n the n -cable of a positive crossing of two n -strand bundles and $T_n = \sigma_1^{-1} \cdots \sigma_{n-1}^{-1} \sigma_n \cdots \sigma_1 \sigma_1 \cdots \sigma_n$; $H(2, 2k + 1) = K_k$ and $H(n, 3)$ is Himeno’s K_n (the companion note [k32.pdf](#) gives the details). Knot Floer homology shows every $H(n, m)$ we could compute to be an L -space knot, and for $m \geq 3$ the volumes exceed every census volume, so they extend the hyperbolic test set beyond the census: 9 such knots satisfy the inequality, none with equality, the smallest margin $1 + |\sigma| - \det$ being 8.

The signature is not determined by the Floer complex. For an L -space knot the Alexander polynomial determines the knot Floer complex, but not the signature. Grouping the 13223 L -space knots of this note by their Alexander polynomials gives 12340 polynomials; 274 are shared by more than one knot, and 37 of those carry two different signatures. The smallest case is classical: $T(3, 4)$ and the $(2, 3)$ -cable of the trefoil have the same knot Floer homology [16, §1.1.3], and signatures -6 and -2 . A hyperbolic case: the census knot t09847 (a braid word due to Teragaito is in [5, Remark 4.2]) has the Alexander polynomial of the $(2, 7)$ -cable of $T(2, 5)$, and signature -10 against -6 . Borodzik, Liu and Zemke give examples of the same phenomenon in the Poincaré sphere [4,

Proposition 3.20, Remark 3.21]. So the agreement $\sigma(K_k) = \sigma(C_k)$ of Proposition 1 is a property of the Baker–Kegel knots, not of their Floer complex, and Conjecture 1 is not a statement about knot Floer homology alone. Within every group the inequality holds for each member. The $(2, q)$ -cables attain equality whatever the companion J : $\det(J_{2,q}) = q$ and $\sigma(J_{2,q}) = -(q - 1)$ by (F1)–(F3).

The sharp case would follow from $\sigma = -2\tau$ for knots with thin knot Floer homology, which is open [15, §8.1].

Two further questions. Which L -space knots attain equality? The known ones are $T(2, n)$, the $(2, q)$ -cables, the Baker–Kegel family, and 70 of the 274 one-bridge parameter triples computed here. Does the inequality follow from a property of the symmetrised Seifert form of an L -space knot, its determinant being bounded by its signature? Definite strongly quasipositive links are studied in [6].

8 A second statement from the same run

The program produced one other statement that its checks could not refute and that we could not place in the literature: for every L -space knot, $u(K) = g(K)$, where u is the unknotting number.

Since $u \geq g_4 = g$ for an L -space knot, the content is $u \leq g$, which holds for braid positive knots by Rudolph, hence for every L -space knot known to be braid positive; by Baker and Kegel [5] that is all but one of the census L -space knots. The companion note `k32.pdf` proves $u = g$ for the whole family $H(n, m)$ of Section 7, by an explicit unknotting matched against the slice–Bennequin bound, so for every Baker–Kegel knot and every Himeno knot, including those that are provably not braid positive. It also shows that the hypothesis is needed: the mirror of $12n_{642}$ is fibered and strongly quasipositive, with $g = 2$ and $u \geq 3$. An earlier version of this section listed unknottings found by search for a few of these knots, and counted Himeno’s K_3 among the knots known not to be braid positive; Himeno proves that for even n only.

9 Reproducibility

Scripts, each writing the JSON or JSONL file the corresponding numbers are read from: `iterated.py` (exact iterated torus knots), `lemma_L.py` (all steps of Lemma 1), `induction_proof.py` ((F1)–(F3) and the induction step), `proof_steps.py` (identity (1)), `census_lspace_fast.sage` (all census L -space knots from the Baker–Kegel braid words), `census_sage_native.sage` (the non- L -space comparison), `families_fast.sage` (families beyond the census), `exact_signature.py` and `exact_census.py` (exact inertia for all census knots), `braid_identity.py` (the closures are the named census knots), `onebridge.sage` and `onebridge_lspace.py` (the one-bridge braid family), `twisted_torus.py` and `twisted_torus_sigma.sage` (the twisted torus family and its exact signatures), and `twisted_torus_ug.py` (unknotting certificates of an earlier version of Section 8), `definiteness.py` and `definiteness_identify.py` (the sharp case across all families), `random_lspace.py` and `random_lspace_sigma.sage` (the random search), and `distinct_knots.py` (how many distinct knots each generated family contains, and which of its hyperbolic knots are in the census). In `families`: `bk_det.sage` and `bk_alex.sage` (Proposition 1), `seifert_dump.py` and `signatures.sage` (the family $H(n, m)$ and K_k for $k \leq 60$), `lspace.py` (which $H(n, m)$ are L -space knots, and their volumes), `sigma_by_alex.sage` (signatures grouped by Alexander polynomial), `onebridge_equality.py` (the one-bridge braids with equality are not hyperbolic), and for the companion note `unknotting.py`, `cables.py` and `fibered_sq.py`. Tools: SnapPy 3.3.2 with `knot_floer_homology`, Sage 10.7, khoca 1.5. The Lean formalisation of the lattice inequality is `K33Lattice/Basic.lean`, built against Mathlib at revision `0df444a360eaa60ab8c11dca51a86af692955474` with `leanprover/lean4:v4.33.1`.

Data provenance. The candidate was generated from KnotInfo [14], accessed on 14 September 2026. No KnotInfo data is reproduced here: its maintainers ask that the database not be reposted, since copies go out of date, so it should be read from <https://knotinfo.org>. The braid words for census knots are from Baker and Kegel’s appendix [5].

References

- [1] J. A. Baldwin, S. Sivek, *L-spaces and knot traces*, arXiv:2501.00914v2 (2026).
- [2] J. A. Baldwin, Y. Hu, S. Sivek, *Khovanov homology and the cinquefoil*, J. Eur. Math. Soc. 27 (2025), 2443–2465; arXiv:2105.12102.
- [3] K. L. Baker, A. H. Moore, *Montesinos knots, Hopf plumbings, and L-space surgeries*, arXiv:1404.7585.
- [4] M. Borodzik, B. Liu, I. Zemke, *Lattice homology, formality, and plumbed L-space links*, arXiv:2210.15792.
- [5] K. L. Baker, M. Kegel, *Census L-space knots are braid positive, except for one that is not*, Algebr. Geom. Topol. 24 (2024); arXiv:2203.12013.
- [6] M. Boileau, S. Boyer, C. McA. Gordon, *On definite strongly quasipositive links and L-space branched covers*, Adv. Math. 357 (2019).
- [7] R. DeYeso III, *Thin knots and the Cabling Conjecture*, Algebr. Geom. Topol. 25 (2025), 4547–4583; arXiv:2112.08074.
- [8] D. Eisenbud, W. Neumann, *Three-dimensional link theory and invariants of plane curve singularities*, Ann. of Math. Studies 110, Princeton Univ. Press (1985).
- [9] P. Feller, *The signature of positive braids is linearly bounded by their genus*, Internat. J. Math. 26 (2015); arXiv:1311.1242.
- [10] E. Farber, B. Reinoso, L. Wang, *Fixed-point-free pseudo-Anosov homeomorphisms, knot Floer homology and the cinquefoil*, Geom. Topol. 28 (2024), 4337–4381; arXiv:2203.01402.
- [11] P. Ghiggini, *Knot Floer homology detects genus-one fibred knots*, Amer. J. Math. 130 (2008), 1151–1169.
- [12] J. E. Greene, L. Liechti, *On the signature of a positive braid*, Ann. Henri Lebesgue 7 (2024), 823–839; arXiv:2308.02275.
- [13] C. McA. Gordon, R. A. Litherland, K. Murasugi, *Signatures of covering links*, Canad. J. Math. 33 (1981), 381–394.
- [14] C. Livingston and A. H. Moore, *KnotInfo: Table of Knot Invariants*, knotinfo.org, September 2026 (accessed 14 September 2026).
- [15] K. Hendricks, C. Manolescu, *Involution Heegaard Floer homology*, Duke Math. J. 166 (2017), 1211–1299; arXiv:1507.00383.
- [16] M. Hedden, *On knot Floer homology and cabling II*, Int. Math. Res. Not. (2009).

- [17] M. Hedden, *Notions of positivity and the Ozsváth–Szabó concordance invariant*, J. Knot Theory Ramifications 19 (2010), 617–629.
- [18] T. Lidman, A. H. Moore, *Pretzel knots with L -space surgeries*, arXiv:1306.6707.
- [19] Y. Ni, *Knot Floer homology detects fibred knots*, Invent. Math. 170 (2007), 577–608.
- [20] J. E. Greene, S. Lewallen, F. Vafaee, *$(1, 1)$ L -space knots*, Compos. Math. 154 (2018), 918–933.
- [21] D. Ruberman, R. İ. Baykur, R. Kirby (eds.), *K3: A New Problem List in Low-Dimensional Topology*, AMS Math. Surveys and Monographs 295 (2026), Problem 1.21.
- [22] K. Himeno, *Non-braid positive hyperbolic L -space knots*, arXiv:2506.22934.
- [23] J. Hom, *A note on cabling and L -space surgeries*, Algebr. Geom. Topol. 11 (2011), 219–223.
- [24] R. A. Litherland, *Signatures of iterated torus knots*, in Topology of Low-Dimensional Manifolds, Lecture Notes in Math. 722 (1979), 71–84.
- [25] P. Ozsváth, Z. Szabó, *On knot Floer homology and lens space surgeries*, Topology 44 (2005), 1281–1300.
- [26] K. Murasugi, *On a certain numerical invariant of link types*, Trans. Amer. Math. Soc. 117 (1965), 387–422.