

Groups of Automata

Carlo Lanzi Luciani

Mentors:

izr. prof. Ganna Kudryavtseva

prof. Alessandro Logar


March 17, 2021

1. Introduction

The word **automaton**: from the greek "acting of one's own will".
Automata are important in:

- ▶ Information theory
- ▶ Theory of dynamical systems
- ▶ Algebra
- ▶ Others

My aim: study some of the groups constructed through a special class of them, the invertible deterministic Mealy automata, here called simply automata.



2. The automaton

Definition

An **automaton** is a 4-tuple $\mathcal{A} = \langle X, Q, \pi, \lambda \rangle$ where:

- ▶ $X = \{x_1, \dots, x_k\}$ is a finite set called the **alphabet**,
- ▶ Q is a set called the **set of internal states of the automaton**,
- ▶ $\pi : X \times Q \longrightarrow Q$ is a function called the **transition function**,
- ▶ $\lambda : X \times Q \longrightarrow X$ is a function such that $\lambda_q = \lambda(\cdot, q) : X \longrightarrow X$ is bijective, and is called the **output function**.

2. The automaton

Definition

An **automaton** is a 4-tuple $\mathcal{A} = \langle X, Q, \pi, \lambda \rangle$ where:

- ▶ $X = \{x_1, \dots, x_k\}$ is a finite set called the **alphabet**,
- ▶ Q is a set called the **set of internal states of the automaton**,
- ▶ $\pi : X \times Q \longrightarrow Q$ is a function called the **transition function**,
- ▶ $\lambda : X \times Q \longrightarrow X$ is a function such that $\lambda_q = \lambda(\cdot, q) : X \longrightarrow X$ is bijective, and is called the **output function**.




Figure: Moore diagram of a 2-state automaton over $X = \{0, 1\}$

3. The initial automaton

Definition

$X^* = \{x_1 x_2 \dots x_n : x_i \in X, n \in \mathbb{N} \cup \{0\}\}$ the **dictionary**.

Word composition: $x_1 \dots x_n \cdot z_1 \dots z_n := x_1 \dots x_n z_1 \dots z_n$

3. The initial automaton

Definition

$X^* = \{x_1x_2 \dots x_n : x_i \in X, n \in \mathbb{N} \cup \{0\}\}$ the **dictionary**.


Word composition: $x_1 \dots x_n \cdot z_1 \dots z_n := x_1 \dots x_n z_1 \dots z_n$

Definition

An **initial automaton** \mathcal{A}_{q_0} is an automaton \mathcal{A} with a fixed state q_0 .

The **action of** \mathcal{A}_{q_0} is the function $\bar{\lambda}_{q_0} : X^* \rightarrow X^*$ with

$\bar{\lambda}_{q_0}(x_1x_2 \dots x_n) = y_1y_2 \dots y_n$.



4. The word tree X^*

Definition

Given $w, v \in X^*$, w is a child of v if and only if $w = v.x = vx$ for some letter $x \in X$.





Figure: An example of the word tree X^* on $X = \{0, 1\}$.

5. Actions as tree-automorphisms

Proposition

A function $f : X^* \rightarrow X^*$ is the action of some initial automaton if and only if it is a **tree-automorphism** on the word tree X^* , i.e.:

- ▶ $f(\emptyset) = \emptyset$.
- ▶ if $w \in X^*$ is a child of v then $f(w)$ is a child of $f(v)$.
- ▶ f is bijective.



Proposition

If f, g are tree-automorphisms on X^* , then $g \circ f$ and f^{-1} are tree-automorphisms on X^* .

6. Groups defined by automata

Proposition

The functions $f : X^* \longrightarrow X^*$ defined by initial automata (i.e. tree-automorphisms), called **synchronous automatic transformations**, form a group denoted by $\mathcal{AUT}_{tree}(X^*)$.

6. Groups defined by automata

Proposition

The functions $f : X^* \longrightarrow X^*$ defined by initial automata (i.e. tree-automorphisms), called **synchronous automatic transformations**, form a group denoted by $\mathcal{AUT}_{tree}(X^*)$.

Definition

Let $\mathcal{A} = \langle X, \mathcal{Q}, \pi, \lambda \rangle$ be an automaton. The **group defined by \mathcal{A}** is the group generated by the set $\{\bar{\lambda}_q : q \in \mathcal{Q}\}$.

$$\begin{array}{ccc} \mathcal{A} \text{ with } & & \bar{\lambda}_{q_1} : X^* \longrightarrow X^* \\ \mathcal{Q} = \{q_1, \dots, q_n\} & \xrightarrow{\text{defines}} & \bar{\lambda}_{q_2} : X^* \longrightarrow X^* \\ & & \vdots \\ & & \bar{\lambda}_{q_n} : X^* \longrightarrow X^* \end{array} \quad \xrightarrow{\text{generate}} \quad \langle \{\bar{\lambda}_q : q \in \mathcal{Q}\} \rangle$$

7. Semidirect product and faithful actions

Definition

Let $(B, *_B), (N, *_N)$ be groups and $\varphi : B \longrightarrow \mathcal{AUT}(N)$ an homomorphism, where $\mathcal{AUT}(N)$ denotes the group of (group-)automorphisms on N . Then the semidirect product $B \ltimes_{\varphi} N$ is the group $(B \times N, *_\varphi)$, with composition rule:

$$(b_2, n_2) *_\varphi (b_1, n_1) = (b_2 *_B b_1, \varphi(b_1)(n_2) *_N n_1)$$

7. Semidirect product and faithful actions

Definition

Let $(B, *_B), (N, *_N)$ be groups and $\varphi : B \rightarrow \mathcal{AUT}(N)$ an homomorphism, where $\mathcal{AUT}(N)$ denotes the group of (group-)automorphisms on N . Then the semidirect product $B \ltimes_{\varphi} N$ is the group $(B \times N, *_\varphi)$, with composition rule:

$$(b_2, n_2) *_\varphi (b_1, n_1) = (b_2 *_B b_1, \varphi(b_1)(n_2) *_N n_1)$$

Definition

Let B be a group and $\mathcal{S}(Y)$ be the symmetric group on Y . Then a faithful action φ of B on Y is a monomorphism $\varphi : B \rightarrow \mathcal{S}(Y)$ and we sign $y^b := \varphi(b)(y)$.

8. Wreath product

Direct sum

Let A be a group and Y a set. The **direct sum of A on Y** is the group

$$A^{(Y)} := \{(a_\omega)_{\omega \in Y} : a_\omega \in A \text{ and } a_\omega \neq 1_A \text{ only for a finite number of } \omega\}.$$

equipped with the component-wise operation of A . If Y is finite we have $A^{(Y)} = A^Y = A \times A \cdots \times A$.

8. Wreath product

Direct sum

Let A be a group and Y a set. The **direct sum of A on Y** is the group

$$A^{(Y)} := \{(a_\omega)_{\omega \in Y} : a_\omega \in A \text{ and } a_\omega \neq 1_A \text{ only for a finite number of } \omega\}.$$

equipped with the component-wise operation of A . If Y is finite we have $A^{(Y)} = A^Y = A \times A \cdots \times A$.

Wreath Product

Let B be a group which acts faithfully on Y and let A be a group. Then the wreath product $B \wr A$ is the semidirect product $B \ltimes_\Phi A^{(Y)} = (B \times A^{(Y)}, *)$ where the composition rule is:

$$\begin{aligned} (g, (u_y)_{y \in Y}) * (h, (v_y)_{y \in Y}) &:= (gh, \Phi(h)((u_y)_{y \in Y})(v_y)_{y \in Y}) := \\ &= (gh, (u_{yh}v_y)_{y \in Y}). \end{aligned}$$

9. Application to automata

Proposition

Let $X = \{x_1, \dots, x_k\}$. Then the function $\psi : \mathcal{AUT}_{tree}(X^*) \longrightarrow \mathcal{S}(X) \wr \mathcal{AUT}_{tree}(X^*) = (\mathcal{S}(X) \times \mathcal{AUT}_{tree}(X^*)^X, *)$ defined by

$$\psi(\bar{\lambda}_{q_0}) = \lambda_{q_0}(\bar{\lambda}_{\pi(x_1, q_0)}, \dots, \bar{\lambda}_{\pi(x_n, q_0)})$$

is an isomorphism of groups.

9. Application to automata

Proposition

Let $X = \{x_1, \dots, x_k\}$. Then the function $\psi : \mathcal{AUT}_{tree}(X^*) \longrightarrow \mathcal{S}(X) \wr \mathcal{AUT}_{tree}(X^*) = (\mathcal{S}(X) \times \mathcal{AUT}_{tree}(X^*)^X, *)$ defined by

$$\psi(\bar{\lambda}_{q_0}) = \lambda_{q_0}(\bar{\lambda}_{\pi(x_1, q_0)}, \dots, \bar{\lambda}_{\pi(x_n, q_0)})$$

is an isomorphism of groups.

Proposition

The wreath product $\mathcal{S}(X) \wr \mathcal{AUT}_{tree}(X^*)$ acts faithfully on X^* as:

$$\lambda_{q_0}(\bar{\lambda}_{\pi(x_1, q_0)}, \dots, \bar{\lambda}_{\pi(x_n, q_0)})(w_1 w_2 w_3 \dots) = \lambda_{q_0}(w_1) \cdot (\bar{\lambda}_{\pi(w_1, q_0)})(w_2 w_3 \dots)$$

10. System of formulas

Proposition

Let \mathcal{A} be an automaton with $\mathcal{Q} = \{q_1, \dots, q_n\}$ over $X = \{x_1, \dots, x_k\}$. Then \mathcal{A} is described by n recurrent formulas

$$\begin{aligned} f_{q_1} &= \beta_{q_1}(h_{x_1, q_1}, \dots, h_{x_k, q_1}), \\ f_{q_2} &= \beta_{q_2}(h_{x_1, q_2}, \dots, h_{x_k, q_2}), \\ &\vdots \\ f_{q_n} &= \beta_{q_n}(h_{x_1, q_n}, \dots, h_{x_k, q_n}), \end{aligned}$$

where each h_{x_i, q_j} is equal to some f_{q_l} and each $\beta_{q_j} \in \mathcal{S}(X)$. Conversely, each such set of n recursive formulas defines an automaton \mathcal{A} such that $\bar{\lambda}_{q_j} = f_{q_j}$ for every $q_j \in \mathcal{Q}$.


$$\begin{array}{ccc} \bar{\lambda}_{q_1} = \lambda_{q_1}(\bar{\lambda}_{\pi(x_1, q_1)}, \dots, \bar{\lambda}_{\pi(x_n, q_1)}) & & f_{q_1} = \beta_{q_1}(h_{x_1, q_1}, \dots, h_{x_k, q_1}) \\ \bar{\lambda}_{q_2} = \lambda_{q_2}(\bar{\lambda}_{\pi(x_1, q_2)}, \dots, \bar{\lambda}_{\pi(x_n, q_2)}) & & f_{q_2} = \beta_{q_2}(h_{x_1, q_2}, \dots, h_{x_k, q_2}) \\ & \vdots & \vdots \\ \bar{\lambda}_{q_n} = \lambda_{q_n}(\bar{\lambda}_{\pi(x_1, q_n)}, \dots, \bar{\lambda}_{\pi(x_n, q_n)}) & & f_{q_n} = \beta_{q_n}(h_{x_1, q_n}, \dots, h_{x_k, q_n}) \end{array}$$

11. Preliminaries

Infinite dihedral group

The infinite dihedral group is the semidirect product $\mathbb{Z}_2 \ltimes_{\phi} \mathbb{Z}$ where $\phi(0)(z) = z$, $\phi(1)(z) = -z$. The group $\mathbb{Z}_2 \ltimes_{\phi} \mathbb{Z}$ acts on \mathbb{Z} from the left:

$$(0, m)z = m + z \quad (1, 0)z = -z \quad (1, m) = (1, 0) * (0, m)$$




11. Preliminaries

Infinite dihedral group

The infinite dihedral group is the semidirect product $\mathbb{Z}_2 \ltimes_{\phi} \mathbb{Z}$ where $\phi(0)(z) = z, \phi(1)(z) = -z$. The group $\mathbb{Z}_2 \ltimes_{\phi} \mathbb{Z}$ acts on \mathbb{Z} from the left:


$$(0, m)z = m + z \quad (1, 0)z = -z \quad (1, m) = (1, 0) * (0, m)$$



Lamplighter group

The lamplighter group is the wreath product $\mathbb{Z} \wr \mathbb{Z}_2 = \mathbb{Z} \ltimes \mathbb{Z}_2^{(\mathbb{Z})}$. The composition rule becomes:

$$(\mathbf{z}_2, (h_i)_{i \in \mathbb{Z}}) * (\mathbf{z}_1, (k_i)_{i \in \mathbb{Z}}) = (\mathbf{z}_2 + \mathbf{z}_1, (h_{i+\mathbf{z}_1} +_{\mathbb{Z}_2} k_i)_{i \in \mathbb{Z}})$$



12. The classification theorem

Theorem

Let \mathcal{A} be a 2-state automaton over the alphabet $X = \{0, 1\}$ and G the group defined by this automaton. Then G is isomorphic to one of the following groups:

- ▶ the trivial group $\{1_G\}$,
- ▶ \mathbb{Z}_2 ,
- ▶ $\mathbb{Z}_2 \times \mathbb{Z}_2$,
- ▶ \mathbb{Z} ,
- ▶ the infinite dihedral group $\mathbb{Z}_2 \rtimes_{\phi} \mathbb{Z}$,
- ▶ the lamplighter group $\mathbb{Z} \wr \mathbb{Z}_2 = \mathbb{Z} \ltimes \mathbb{Z}_2^{(\mathbb{Z})}$.

13. Sketch of proof

Define the cases

Let $Q = \{r, s\}$ and $a = \bar{\lambda}_r, b = \bar{\lambda}_s$, then

$$\mathcal{A} \leftarrow \text{-----} \rightarrow \begin{array}{l} a = \tau^{i_1}(x_{11}, x_{12}) \\ b = \tau^{i_2}(x_{21}, x_{22}) \end{array}$$

where $x_{ij} \in \{a, b\}$ and $\tau^{i_1}, \tau^{i_2} \in \mathcal{S}(X) = \mathcal{S}(\{0, 1\})$. There are 64 possibilities.

13. Sketch of proof

Define the cases

Let $\mathcal{Q} = \{r, s\}$ and $a = \bar{\lambda}_r, b = \bar{\lambda}_s$, then

$$\mathcal{A} \longleftrightarrow \begin{array}{l} a = \tau^{i_1}(x_{11}, x_{12}) \\ b = \tau^{i_2}(x_{21}, x_{22}) \end{array}$$

where $x_{ij} \in \{a, b\}$ and $\tau^{i_1}, \tau^{i_2} \in \mathcal{S}(X) = \mathcal{S}(\{0, 1\})$. There are 64 possibilities.

Case analysis

Let τ be the transposition of $\mathcal{S}(X)$. Every case is analogous to one of the 24 cases where $a \in \{\tau(a, a), \tau(a, b), \tau(b, b)\}$. We proceed by analysing part of them.

Thank you for your attention!