

Intensional paradoxes and the maxim of maximal recapture

Carlo Nicolai



Slides available at <https://carlonicolai.github.io>

*'There never were any set-theoretic paradoxes, but the
property theoretic paradoxes are still unresolved'
(Gödel via Myhill 1984)*

By intensional paradoxes one might refer to:

- ▶ semantic
- ▶ property-theoretic
- ▶ paradoxes involving modalities and propositional attitudes (in predicate form)

*'There never were any set-theoretic paradoxes, but the
property theoretic paradoxes are still unresolved'
(Gödel via Myhill 1984)*

By intensional paradoxes one might refer to:

- ▶ semantic
- ▶ property-theoretic
- ▶ paradoxes involving modalities and propositional attitudes (in predicate form)

In the talk I'll essentially deal with the first two. The gist of the talk will mainly be methodological: paradoxical sentences are so far away from empirical testing or conceptual practices that not much consensus can be gained by reflecting on our intuitions about them.

Part I: Some theories of self-applicable predication

I will use the notion of 'predication' as a bridge concept lying between the semantic notion of **satisfaction** and the more metaphysical notion of **instantiation**.

I will use the notion of 'predication' as a bridge concept lying between the semantic notion of **satisfaction** and the more metaphysical notion of **instantiation**.

The theories that I introduce will all be 'solutions' to the intensional paradoxes in Gödel's sense, in the sense of involving **non-trivial self-referential constructions**:

I will use the notion of 'predication' as a bridge concept lying between the semantic notion of **satisfaction** and the more metaphysical notion of **instantiation**.

The theories that I introduce will all be 'solutions' to the intensional paradoxes in Gödel's sense, in the sense of involving **non-trivial self-referential constructions**:

- ▶ Nonclassical theories:
 - ▶ theories based on the logic **K3** of 'gaps' (Field, Reinhardt);
 - ▶ theories based on the logic **LP** of 'gluts' (Priest, Beall);
- ▶ Classical theories of predication (Kripke, Feferman, Turner).

I will use the notion of 'predication' as a bridge concept lying between the semantic notion of **satisfaction** and the more metaphysical notion of **instantiation**.

The theories that I introduce will all be 'solutions' to the intensional paradoxes in Gödel's sense, in the sense of involving **non-trivial self-referential constructions**:

- ▶ Nonclassical theories:
 - ▶ theories based on the logic **K3** of 'gaps' (Field, Reinhardt);
 - ▶ theories based on the logic **LP** of 'gluts' (Priest, Beall);
- ▶ Classical theories of predication (Kripke, Feferman, Turner).

They will capture the same such solution in a sense that will be made precise.

Let's first make precise our support theory of **syntax/nominalization**:

Let's first make precise our support theory of **syntax/nominalization**:

Definition (Basic^K)

Basic^K contains Q – conveniently formulated with $<$ – and axioms for sequences, projections, lengths. Crucially:

$$\exists s(\text{lh}(s) = 0)$$

$$\forall s(\text{lh}(s) = n \rightarrow \exists s'(\text{lh}(s') = n + 1 \wedge (\forall m < n)(s_m = s'_m)))$$

Let's first make precise our support theory of **syntax/nominalization**:

Definition (Basic^K)

Basic^K contains Q – conveniently formulated with $<$ – and axioms for sequences, projections, lengths. Crucially:

$$\begin{aligned} &\exists s(\text{lh}(s) = 0) \\ &\forall s(\text{lh}(s) = n \rightarrow \exists s'(\text{lh}(s') = n + 1 \wedge (\forall m < n)(s_m = s'_m))) \end{aligned}$$

Remarks:

- ▶ The superscript $(\cdot)^K$ stands for 'classical logic'
- ▶ Basic^K is interpretable in Q , although not in a way that preserves identity since Q does not have a **general** notion of sequence;
- ▶ So far we have **no induction**, either on strings or on numbers.

Let's consider now a language \mathcal{L} containing $\mathcal{L}_{\text{Basic}}$ and a predicate $P(x, y)$ for **predication**. From a canonical Gödel numbering of \mathcal{L} -expression we define

$$\lambda x A(x, \vec{y}) := \langle \ulcorner A \urcorner, \vec{y} \rangle$$

where $\ulcorner A \urcorner$ is a (term representing the) code of A . We mostly deal with unary wwfs $A(v)$.

Let's consider now a language \mathcal{L} containing $\mathcal{L}_{\text{Basic}}$ and a predicate $P(x, y)$ for **predication**. From a canonical Gödel numbering of \mathcal{L} -expression we define

$$\lambda x A(x, \vec{y}) := \langle \ulcorner A \urcorner, \vec{y} \rangle$$

where $\ulcorner A \urcorner$ is a (term representing the) code of A . We mostly deal with unary wwfs $A(v)$.

As it is well-known, Basic^K formulated in \mathcal{L} is **inconsistent** with full predication rules:

$$\frac{\Gamma \Rightarrow A(x), \Delta}{\Gamma \Rightarrow P(\ulcorner A(v) \urcorner, x), \Delta} \qquad \frac{\Gamma, A(x) \Rightarrow \Delta}{\Gamma, P(\ulcorner A(v) \urcorner, x) \Rightarrow \Delta}$$

This holds equally well for intuitionistic versions of the theory of sequences and predication rules.

Restricting the logic

Definition (Basic De Morgan Logic)

BDM is the **fully structural** logic obtained by considering and positive and negative rules for monotone connectives and quantifiers, e.g.:

$$\frac{\Gamma \Rightarrow \Delta, \neg A, \neg B}{\Gamma \Rightarrow \Delta, \neg(A \wedge B)}$$

$$\frac{\Gamma, \neg A \Rightarrow \Delta \quad \Gamma, \neg B \Rightarrow \Delta}{\Gamma, \neg(A \wedge B) \Rightarrow \Delta}$$

and only **double negation** rules:

$$\frac{\Gamma \Rightarrow \Delta, A}{\Gamma \Rightarrow \Delta, \neg\neg A}$$

$$\frac{A, \Gamma \Rightarrow \Delta}{\neg\neg A, \Gamma \Rightarrow \Delta}$$

Restricting the logic

Definition (Basic De Morgan Logic)

BDM is the **fully structural** logic obtained by considering and positive and negative rules for monotone connectives and quantifiers, e.g.:

$$\frac{\Gamma \Rightarrow \Delta, \neg A, \neg B}{\Gamma \Rightarrow \Delta, \neg(A \wedge B)}$$

$$\frac{\Gamma, \neg A \Rightarrow \Delta \quad \Gamma, \neg B \Rightarrow \Delta}{\Gamma, \neg(A \wedge B) \Rightarrow \Delta}$$

and only **double negation** rules:

$$\frac{\Gamma \Rightarrow \Delta, A}{\Gamma \Rightarrow \Delta, \neg\neg A}$$

$$\frac{A, \Gamma \Rightarrow \Delta}{\neg\neg A, \Gamma \Rightarrow \Delta}$$

Definition (LP, K3)

LP and K3 are now obtained by adding to BDM, respectively:

$$\frac{\Gamma, A \Rightarrow \Delta}{\Gamma \Rightarrow \Delta, \neg A}$$

$$\frac{\Gamma \Rightarrow \Delta, A}{\neg A, \Gamma \Rightarrow \Delta}$$

Definition (Basic^{K3} , Basic^{LP})

Basic^{K3} and Basic^{LP} are obtained by extending $K3$ and LP in $\mathcal{L}_{\text{Basic}}$, respectively, with

- ▶ Identity sequents (we can treat them classically given our applications)

$$\Rightarrow t = t$$

$$s = t, A(s) \Rightarrow A(t)$$

- ▶ Sequents $\Rightarrow A$ for A an axiom of Q or of sequences.

Principles for Predication/Satisfaction for properties/unary formulas

$$(\text{At}+) \quad P(\ulcorner R(v) \urcorner, x) \Leftrightarrow R(x)$$

$$(\text{At}-) \quad P(\ulcorner \neg R(v) \urcorner, x) \Leftrightarrow \neg R(x)$$

for R in $\mathcal{L}_{\text{Basic}}$

Principles for Predication/Satisfaction for properties/unary formulas

$$(At+) \quad P(\ulcorner R(v) \urcorner, x) \Leftrightarrow R(x)$$

$$(At-) \quad P(\ulcorner \neg R(v) \urcorner, x) \Leftrightarrow \neg R(x)$$

for R in $\mathcal{L}_{\text{Basic}}$

$$(P1) \quad P(\ulcorner P(u, v) \urcorner, (x, y)) \Leftrightarrow P(x, y)$$

$$(P2) \quad P(\ulcorner \neg P(u, v) \urcorner, (x, y)) \Leftrightarrow \neg P(x, y)$$

Principles for Predication/Satisfaction for properties/unary formulas

$$(At+) \quad P(\ulcorner R(v) \urcorner, x) \Leftrightarrow R(x)$$

$$(At-) \quad P(\ulcorner \neg R(v) \urcorner, x) \Leftrightarrow \neg R(x)$$

for R in $\mathcal{L}_{\text{Basic}}$

$$(P1) \quad P(\ulcorner P(u, v) \urcorner, (x, y)) \Leftrightarrow P(x, y)$$

$$(P2) \quad P(\ulcorner \neg P(u, v) \urcorner, (x, y)) \Leftrightarrow \neg P(x, y)$$

$$(\neg 1) \quad P(\ulcorner \neg \varphi \urcorner, x) \Leftrightarrow \neg P(\ulcorner \varphi \urcorner, x)$$

$$(\neg 2) \quad P(\ulcorner \neg \neg \varphi \urcorner, x) \Leftrightarrow P(\ulcorner \varphi \urcorner, x)$$

Principles for Predication/Satisfaction for properties/unary formulas

$$(At+) \quad P(\ulcorner R(v) \urcorner, x) \Leftrightarrow R(x)$$

$$(At-) \quad P(\ulcorner \neg R(v) \urcorner, x) \Leftrightarrow \neg R(x)$$

for R in $\mathcal{L}_{\text{Basic}}$

$$(P1) \quad P(\ulcorner P(u, v) \urcorner, (x, y)) \Leftrightarrow P(x, y)$$

$$(P2) \quad P(\ulcorner \neg P(u, v) \urcorner, (x, y)) \Leftrightarrow \neg P(x, y)$$

$$(\neg 1) \quad P(\ulcorner \neg \varphi \urcorner, x) \Leftrightarrow \neg P(\ulcorner \varphi \urcorner, x)$$

$$(\neg 2) \quad P(\ulcorner \neg \neg \varphi \urcorner, x) \Leftrightarrow P(\ulcorner \varphi \urcorner, x)$$

$$(\wedge 1) \quad P(\ulcorner \varphi \wedge \psi \urcorner, x) \Leftrightarrow P(\ulcorner \varphi \urcorner, x) \wedge P(\ulcorner \psi \urcorner, x)$$

$$(\wedge 2) \quad P(\ulcorner \neg(\varphi \wedge \psi) \urcorner, x) \Leftrightarrow P(\ulcorner \neg \varphi \urcorner, x) \vee P(\ulcorner \neg \psi \urcorner, x)$$

\vdots

Paracomplete and paraconsistent satisfaction

$$(At+) \quad P(\ulcorner R(v) \urcorner, x) \Leftrightarrow R(x)$$

~~$$P(\ulcorner \neg R(v) \urcorner, x) \Leftrightarrow \neg R(x)$$~~

~~for R in \mathcal{L}_{Basic}~~

$$(P1) \quad P(\ulcorner P(u, v) \urcorner, (x, y)) \Leftrightarrow P(x, y)$$

~~$$P(\ulcorner \neg P(u, v) \urcorner, (x, y)) \Leftrightarrow \neg P(x, y)$$~~

~~$$(\neg 1) \quad P(\ulcorner \neg \varphi \urcorner, x) \Leftrightarrow \neg P(\ulcorner \varphi \urcorner, x)$$~~

$$(\neg 2) \quad P(\ulcorner \neg \neg \varphi \urcorner, x) \Leftrightarrow P(\ulcorner \varphi \urcorner, x)$$

$$(\wedge 1) \quad P(\ulcorner \varphi \wedge \psi \urcorner, x) \Leftrightarrow P(\ulcorner \varphi \urcorner, x) \wedge P(\ulcorner \psi \urcorner, x)$$

~~$$(\wedge 2) \quad P(\ulcorner \neg(\varphi \wedge \psi) \urcorner, x) \Leftrightarrow P(\ulcorner \neg \varphi \urcorner, x) \vee P(\ulcorner \neg \psi \urcorner, x)$$~~

Paracomplete and paraconsistent satisfaction

$$\begin{aligned}(\text{At}+) \quad & P(\ulcorner R(v) \urcorner, x) \Leftrightarrow R(x) \\ & \cancel{P(\ulcorner \neg R(v) \urcorner, x) \Leftrightarrow \neg R(x)} \quad \text{for } R \text{ in } \mathcal{L}_{\text{Basic}}\end{aligned}$$

$$\begin{aligned}(\text{P1}) \quad & P(\ulcorner P(u, v) \urcorner, (x, y)) \Leftrightarrow P(x, y) \\ & \cancel{P(\ulcorner \neg P(u, v) \urcorner, (x, y)) \Leftrightarrow \neg P(x, y)}\end{aligned}$$

$$(\neg 1) \quad \cancel{P(\ulcorner \neg \varphi \urcorner, x) \Leftrightarrow \neg P(\varphi, x)}$$

$$(\neg 2) \quad P(\ulcorner \neg \neg \varphi \urcorner, x) \Leftrightarrow P(\varphi, x)$$

$$(\wedge 1) \quad P(\varphi \wedge \psi, x) \Leftrightarrow P(\varphi, x) \wedge P(\psi, x)$$

$$(\wedge 2) \quad \cancel{P(\ulcorner \neg(\varphi \wedge \psi) \urcorner, x) \Leftrightarrow P(\ulcorner \neg \varphi \urcorner, x) \vee P(\ulcorner \neg \psi \urcorner, x)}$$

Definition (K3P_0 , LPP_0)

Extend respectively Basic^{K3} and Basic^{LP} with (At), (P1), ($\neg 1$), ($\wedge 1$) (and quantifier sequents).

Classical, Self-referential satisfaction

$$(At+) \quad P(\ulcorner R(v) \urcorner, x) \Leftrightarrow R(x)$$

$$(At-) \quad P(\ulcorner \neg R(v) \urcorner, x) \Leftrightarrow \neg R(x)$$

for R in $\mathcal{L}_{\text{Basic}}$

$$(P1) \quad P(\ulcorner P(u, v) \urcorner, (x, y)) \Leftrightarrow P(x, y)$$

$$(P2) \quad P(\ulcorner \neg P(u, v) \urcorner, (x, y)) \Leftrightarrow \neg P(x, y)$$

~~$$(\neg 1) \quad P(\ulcorner \neg \varphi \urcorner, x) \Leftrightarrow \neg P(\ulcorner \varphi \urcorner, x)$$~~

$$(\neg 2) \quad P(\ulcorner \neg \neg \varphi \urcorner, x) \Leftrightarrow P(\ulcorner \varphi \urcorner, x)$$

$$(\wedge 1) \quad P(\ulcorner \varphi \wedge \psi \urcorner, x) \Leftrightarrow P(\ulcorner \varphi \urcorner, x) \wedge P(\ulcorner \psi \urcorner, x)$$

$$(\wedge 2) \quad P(\ulcorner \neg(\varphi \wedge \psi) \urcorner, x) \Leftrightarrow P(\ulcorner \neg \varphi \urcorner, x) \vee P(\ulcorner \neg \psi \urcorner, x) \dots$$

Definition (KFP_0)

Extend Basic^K with $(At+)$ - $(\wedge 2)$ (and quantifier sequents).

Models

Fixed Point Model

A set $S \subseteq \mathbb{N}$ is a **fixed point** iff

$$(\varphi, n) \in S \quad \text{iff} \quad P(\varphi, n) \in S$$

Models

Fixed Point Model

A set $S \subseteq \mathbb{N}$ is a **fixed point** iff

$$(\varphi, n) \in S \text{ iff } P(\varphi, n) \in S$$

Consistent Fixed Points

For S a **consistent** fixed point, i.e. for no φ, n : $(\varphi, n) \in S$ and $(\neg\varphi, n) \in S$,

$$(\mathbb{N}, S) \models \text{KFP}_0 + P(\neg\varphi, x) \Rightarrow \neg P(\varphi, x) \text{ iff } (\mathbb{N}, S) \models_{\text{K3}} \text{K3P}_0$$

Models

Fixed Point Model

A set $S \subseteq \mathbb{N}$ is a **fixed point** iff

$$(\varphi, n) \in S \quad \text{iff} \quad P(\varphi, n) \in S$$

Consistent Fixed Points

For S a **consistent** fixed point, i.e. for no φ, n : $(\varphi, n) \in S$ and $(\neg\varphi, n) \in S$,

$$(\mathbb{N}, S) \models \text{KFP}_0 + P(\neg\varphi, x) \Rightarrow \neg P(\varphi, x) \quad \text{iff} \quad (\mathbb{N}, S) \models_{\text{K3}} \text{K3P}_0$$

Complete Fixed point

For S a **complete** fixed point, i.e. for every φ, n : $(\varphi, n) \in S$ or $(\neg\varphi, n) \in S$,

$$(\mathbb{N}, S) \models \text{KFP}_0 + \neg P(\varphi, x) \Rightarrow P(\neg\varphi, x) \quad \text{iff} \quad (\mathbb{N}, S) \models_{\text{LP}} \text{LPP}_0$$

Part II: Recapture(s)?

No restrictions

Fact

The **unrestricted predication** rules

$$\frac{\Gamma \Rightarrow A(x), \Delta}{\Gamma \Rightarrow P(\ulcorner A(v) \urcorner, x), \Delta}$$

$$\frac{\Gamma, A(x) \Rightarrow \Delta}{\Gamma, P(\ulcorner A(v) \urcorner, x) \Rightarrow \Delta}$$

are **admissible** in $K3P_0$ and LPP_0 (with $A(v) \in \mathcal{L}$).

No restrictions

Fact

The **unrestricted predication** rules

$$\frac{\Gamma \Rightarrow A(x), \Delta}{\Gamma \Rightarrow P(\ulcorner A(v) \urcorner, x), \Delta}$$

$$\frac{\Gamma, A(x) \Rightarrow \Delta}{\Gamma, P(\ulcorner A(v) \urcorner, x) \Rightarrow \Delta}$$

are **admissible** in $K3P_0$ and LPP_0 (with $A(v) \in \mathcal{L}$).

The classical theory KFP_0 and its extensions with consistency / completeness principles don't have this property. The Russell property $R(x) \leftrightarrow \neg P(x, x)$ is such that:

No restrictions

Fact

The **unrestricted predication** rules

$$\frac{\Gamma \Rightarrow A(x), \Delta}{\Gamma \Rightarrow P(\ulcorner A(v) \urcorner, x), \Delta}$$

$$\frac{\Gamma, A(x) \Rightarrow \Delta}{\Gamma, P(\ulcorner A(v) \urcorner, x) \Rightarrow \Delta}$$

are **admissible** in $K3P_0$ and LPP_0 (with $A(v) \in \mathcal{L}$).

The classical theory KFP_0 and its extensions with consistency / completeness principles don't have this property. The Russell property $R(x) \leftrightarrow \neg P(x, x)$ is such that:

- ▶ $KFP_0 + \text{CONS}$ proves that the Russell's property is **not** predicable of itself but the predicate $R(x)$ **applies** to $\ulcorner R \urcorner$;

No restrictions

Fact

The **unrestricted predication** rules

$$\frac{\Gamma \Rightarrow A(x), \Delta}{\Gamma \Rightarrow P(\ulcorner A(v) \urcorner, x), \Delta}$$

$$\frac{\Gamma, A(x) \Rightarrow \Delta}{\Gamma, P(\ulcorner A(v) \urcorner, x) \Rightarrow \Delta}$$

are **admissible** in $K3P_0$ and LPP_0 (with $A(v) \in \mathcal{L}$).

The classical theory KFP_0 and its extensions with consistency / completeness principles don't have this property. The Russell property $R(x) \leftrightarrow \neg P(x, x)$ is such that:

- ▶ $KFP_0 + \text{CONS}$ proves that the Russell's property is **not** predicable of itself but the predicate $R(x)$ **applies** to $\ulcorner R \urcorner$;
- ▶ $KFP_0 + \text{COMP}$ proves that the Russell's property **is** predicable of itself but the predicate $R(x)$ **does not** apply to $\ulcorner R \urcorner$.

Recapture

The problem with $K3P_0$ and LPP_0 approaches is well-known: principles such as the **the law of excluded middle** – in the case of $K3P_0$ – or **material modus ponens** – for LPP_0 :

‘nothing like sustained ordinary reasoning can be carried out in either logic’ (Feferman 1984)

Recapture

The problem with $K3P_0$ and LPP_0 approaches is well-known: principles such as the [the law of excluded middle](#) – in the case of $K3P_0$ – or [material modus ponens](#) – for LPP_0 :

‘nothing like sustained ordinary reasoning can be carried out in either logic’ (Feferman 1984)

In our framework, the paracomplete/paracosistent theorist’s reassuring remarks to the mathematician can be translated as follows:

Lemma (Recapture)

For $A \in \mathcal{L}_{\text{Basic}}$:

- ▶ $K3P_0$ proves $\Rightarrow A \vee \neg A$;
- ▶ LPP_0 proves $A \wedge \neg A \Rightarrow$.

In particular, this means that if one restricts the attention to $\mathcal{L}_{\text{Basic}}$, one obtains that [Basic^K](#), [Basic^{LP}](#), and [Basic^{K3}](#) are identical theories and classical reasoning can be used unrestrictedly.

Schemata

Our understanding of the language of arithmetic is such that we anticipate that the Induction Axiom Schema will persist through all changes. [...] We are always capable of generating new Induction Axioms by expanding the language. (McGee, How we learn mathematical language, p. 58)

Schemata

Our understanding of the language of arithmetic is such that we anticipate that the Induction Axiom Schema will persist through all changes. [...] We are always capable of generating new Induction Axioms by expanding the language. (McGee, How we learn mathematical language, p. 58)

What happens if we add induction schemata to our theories?

$$\frac{\Gamma, A(x) \Rightarrow A(x+1), \Delta}{\Gamma, A(0) \Rightarrow A(y), \Delta}$$

for x not free in $A(0), \Gamma, \Delta$ and y arbitrary. We call the resulting theories KFP, K3P, LPP.

Schemata

Our understanding of the language of arithmetic is such that we anticipate that the Induction Axiom Schema will persist through all changes. [...] We are always capable of generating new Induction Axioms by expanding the language. (McGee, How we learn mathematical language, p. 58)

What happens if we add induction schemata to our theories?

$$\frac{\Gamma, A(x) \Rightarrow A(x+1), \Delta}{\Gamma, A(0) \Rightarrow A(y), \Delta}$$

for x not free in $A(0), \Gamma, \Delta$ and y arbitrary. We call the resulting theories **KFP**, **K3P**, **LPP**.

Notice that this is not a logical *divertissement*: there are foundational programs (Aczel, Feferman, Schütte, Turner) that wish to start with natural numbers and justify higher mathematics via self-referential predication/satisfaction axioms. Similar remarks can be made, with a bit of effort, for **ZFC** and classes over it.

Transfinite Induction

We assume a notation (O, \prec) for the ordinals up to the Feferman - Schütte ordinal Γ_0 . Transfinite induction up to any $\beta < \Gamma_0$ is the schema, for $\delta < \beta$ and $A(v) \in \mathcal{L}$,

$$\frac{\Gamma, \forall \gamma \prec \alpha A(\gamma) \Rightarrow A(\alpha), \Delta}{\Gamma \Rightarrow \forall \gamma \prec \delta A(\gamma), \Delta} \text{TI}_{\mathcal{L}}(< \beta)$$

Transfinite Induction

We assume a notation (O, \prec) for the ordinals up to the Feferman - Schütte ordinal Γ_0 . Transfinite induction up to any $\beta < \Gamma_0$ is the schema, for $\delta < \beta$ and $A(v) \in \mathcal{L}$,

$$\frac{\Gamma, \forall \gamma \prec \alpha A(\gamma) \Rightarrow A(\alpha), \Delta}{\Gamma \Rightarrow \forall \gamma \prec \delta A(\gamma), \Delta} \text{TI}_{\mathcal{L}}(< \beta)$$

Proposition

By the classical Gentzen argument, KFP proves $\text{TI}_{\mathcal{L}}(< \varepsilon_0)$, whereas K3P and LPP **can only** prove $\text{TI}_{\mathcal{L}}(< \omega^\omega)$. The former is proof-theoretically equivalent to $\text{ACA}_{<\varepsilon_0}$, the second to $\text{ACA}_{<\omega^\omega}$.

Transfinite Induction

We assume a notation (O, \prec) for the ordinals up to the Feferman - Schütte ordinal Γ_0 . Transfinite induction up to any $\beta < \Gamma_0$ is the schema, for $\delta < \beta$ and $A(v) \in \mathcal{L}$,

$$\frac{\Gamma, \forall \gamma \prec \alpha A(\gamma) \Rightarrow A(\alpha), \Delta}{\Gamma \Rightarrow \forall \gamma \prec \delta A(\gamma), \Delta} \text{TI}_{\mathcal{L}}(< \beta)$$

Proposition

By the classical Gentzen argument, KFP proves $\text{TI}_{\mathcal{L}}(< \varepsilon_0)$, whereas K3P and LPP **can only** prove $\text{TI}_{\mathcal{L}}(< \omega^\omega)$. The former is proof-theoretically equivalent to $\text{ACA}_{<\varepsilon_0}$, the second to $\text{ACA}_{<\omega^\omega}$.

And crucially:

Corollary

$\text{Basic}^{\text{LP}} + \text{Ind}(\mathcal{L})$, and $\text{Basic}^{\text{K3}} + \text{Ind}(\mathcal{L})$ can only prove $\text{TI}_{\mathcal{L}}(< \omega^\omega)$, whereas $\text{Basic}^{\text{K}} + \text{Ind}(\mathcal{L})$ proves $\text{TI}_{\mathcal{L}}(< \varepsilon_0)$.

Non-semantic consequences /1

If one carefully looks at the proof-theory of these systems, it's clear that also in terms of mathematical consequences they are very different:

Proposition

- ▶ KFP proves $\text{TI}_{\mathcal{L}^{\text{Basic}}}(< \phi_{\varepsilon_0} 0)$ (same as $\text{ACA}_{<\varepsilon_0}$);
- ▶ K3P and LS3 prove $\text{TI}_{\mathcal{L}^{\text{Basic}}}(< \phi_\omega 0)$ (same as $\text{ACA}_{<\omega^\omega}$).

Non-semantic consequences /1

If one carefully looks at the proof-theory of these systems, it's clear that also in terms of mathematical consequences they are very different:

Proposition

- ▶ KFP proves $\text{TI}_{\mathcal{L}^{\text{Basic}}}(< \phi_{\varepsilon_0} 0)$ (same as $\text{ACA}_{<\varepsilon_0}$);
- ▶ K3P and LS3 prove $\text{TI}_{\mathcal{L}^{\text{Basic}}}(< \phi_\omega 0)$ (same as $\text{ACA}_{<\omega^\omega}$).

But the defender of gaps or gluts is within their rights to ask whether the the loss of the classically accepted pattern of reasoning by transfinite induction is in fact a **mathematically significant** one.

Non-semantic consequences /2

INDEC is the assertion that:

every countable scattered indecomposable linear ordering is either indecomposable to the left, or indecomposable to the right.

It was proved to be true by Pierre Jullien in 1969.

Non-semantic consequences /2

INDEC is the assertion that:

every countable scattered indecomposable linear ordering is either indecomposable to the left, or indecomposable to the right.

It was proved to be true by Pierre Jullien in 1969.

Lemma (Montalbán, Friedman)

INDEC is implied by Σ_1^1 -CA, which is Π_2^1 -conservative over $\text{ACA}_{<\varepsilon_0}$.

Non-semantic consequences /2

INDEC is the assertion that:

every countable scattered indecomposable linear ordering is either indecomposable to the left, or indecomposable to the right.

It was proved to be true by Pierre Jullien in 1969.

Lemma (Montalbán, Friedman)

INDEC is implied by Σ_1^1 -CA, which is Π_2^1 -conservative over $\text{ACA}_{<\varepsilon_0}$.

Proposition (Eastaugh, N.)

$\text{RCA} + \text{INDEC}$ is proof-theoretically equivalent to KFP . It follows that KFP can ‘nicely’ interpret $\text{RCA} + \text{INDEC}$, but neither K3P nor LPP can.

- ▶ The advocate of **K3P** and **LPP** will rely on recapture theorems to claim that a purported solution to paradox does not mess around with **standard mathematical reasoning**.

- ▶ The advocate of **K3P** and **LPP** will rely on recapture theorems to claim that a purported solution to paradox does not mess around with **standard mathematical reasoning**.
- ▶ Under the presupposition that the schema of mathematical induction is to be understood as (weakly) open-ended, mathematical patterns of reasoning are **severely crippled** when one moves to from classical to non-classical theories of predication.

- ▶ The advocate of **K3P** and **LPP** will rely on recapture theorems to claim that a purported solution to paradox does not mess around with **standard mathematical reasoning**.
- ▶ Under the presupposition that the schema of mathematical induction is to be understood as (weakly) open-ended, mathematical patterns of reasoning are **severely crippled** when one moves to from classical to non-classical theories of predication.
- ▶ Even if one is suspicious about the status of 'axioms' or logical principles such as iterations of comprehension, there are **theorems**, such as **INDEC**, that separate the two clusters of theories.

Part III: Bridges

Reflection

One way to fill this mathematical gap is to resort to Feferman's theory of **implicit commitment**. We can enlarge **K3P** or **LPP** with a reflection principle of the form, for $S = \text{K3P}, \text{LPP}$,

$$(R) \frac{\text{Pr}_S^2(\ulcorner \Gamma \dot{x} \Rightarrow \Delta \dot{x} \urcorner, \ulcorner \Theta \dot{x} \Rightarrow \Lambda \dot{x} \urcorner) \quad \Gamma(x) \Rightarrow \Delta(x)}{\Theta(x) \Rightarrow \Lambda(x)}$$

Reflection

One way to fill this mathematical gap is to resort to Feferman's theory of **implicit commitment**. We can enlarge **K3P** or **LPP** with a reflection principle of the form, for $S = \text{K3P}, \text{LPP}$,

$$(R) \frac{\text{Pr}_S^2(\ulcorner \Gamma \dot{x} \Rightarrow \Delta \dot{x} \urcorner, \ulcorner \Theta \dot{x} \Rightarrow \Lambda \dot{x} \urcorner) \quad \Gamma(x) \Rightarrow \Delta(x)}{\Theta(x) \Rightarrow \Lambda(x)}$$

Proposition (Fischer, N., Horsten)

- ▶ Finitely many iterations of reflection over S give us $\text{TI}_{\mathcal{L}}(< \omega^{\omega^2})$;
- ▶ $< \varepsilon_0$ -iterations of reflection give us the same transfinite induction as **KFP** and variants thereof.

Reflection

One way to fill this mathematical gap is to resort to Feferman's theory of **implicit commitment**. We can enlarge **K3P** or **LPP** with a reflection principle of the form, for $S = \text{K3P}, \text{LPP}$,

$$(R) \frac{\text{Pr}_S^2(\ulcorner \Gamma \dot{x} \Rightarrow \Delta \dot{x} \urcorner, \ulcorner \Theta \dot{x} \Rightarrow \Lambda \dot{x} \urcorner) \quad \Gamma(x) \Rightarrow \Delta(x)}{\Theta(x) \Rightarrow \Lambda(x)}$$

Proposition (Fischer, N., Horsten)

- ▶ Finitely many iterations of reflection over S give us $\text{TI}_{\mathcal{L}}(< \omega^{\omega^2})$;
- ▶ $< \varepsilon_0$ -iterations of reflection give us the same transfinite induction as **KFP** and variants thereof.

An obvious problem with this strategy is that it is available also to the classical logician, especially the one willing to accept gluts internally.

A conditional

We can extend **Basic De Morgan Logic** with a new conditional satisfying

$$\frac{\Gamma, A \Rightarrow B}{\Gamma \Rightarrow A \rightarrow B} \qquad \frac{\Gamma \Rightarrow A \quad B, \Gamma \Rightarrow \Delta}{\Gamma, A \rightarrow B \Rightarrow \Delta}$$

Definition (HBP)

We extend **Basic**^{BDM \rightarrow} with the axiom

$$\Rightarrow P(\ulcorner A \urcorner, x) \leftrightarrow A(x)$$

for A containing P but not \rightarrow itself.

A conditional

We can extend **Basic De Morgan Logic** with a new conditional satisfying

$$\frac{\Gamma, A \Rightarrow B}{\Gamma \Rightarrow A \rightarrow B} \qquad \frac{\Gamma \Rightarrow A \quad B, \Gamma \Rightarrow \Delta}{\Gamma, A \rightarrow B \Rightarrow \Delta}$$

Definition (HBP)

We extend **Basic**^{BDM[→]} with the axiom

$$\Rightarrow P(\ulcorner A \urcorner, x) \leftrightarrow A(x)$$

for A containing P but not \rightarrow itself.

These rules are sound with respect to the class of fixed point models for satisfaction / predication extracted from Leitgeb's system **HYPE** (roughly a 'hyperintensional conditional').

By carefully reproducing Gentzen's argument one should be able to obtain:

Conjecture

HBP proves $\text{TI}_{\mathcal{L}}(< \varepsilon_0)$ (indeed already in $\text{Basic}^{\text{BDM}^+}$).

By carefully reproducing Gentzen's argument one should be able to obtain:

Conjecture

HBP proves $\text{TI}_{\mathcal{L}}(< \varepsilon_0)$ (indeed already in $\text{Basic}^{\text{BDM}^{\rightarrow}}$).

If true, it is still not clear whether this claim may help the gap / glut theorist, unless it is established that there is no classical, natural counterpart to HBP and its extensions with compositional clauses.

In addition, we cannot have unrestricted principles anymore.

THANK YOU.