

# *The least of all evils*

---

Carlo Nicolai



Bristol, May 3, 2019  
(slides at [carlonicolai.github.io](https://carlonicolai.github.io))

---



- I am presenting a cluster of nonclassical systems for (transparent) truth and consequence.

- I am presenting a cluster of nonclassical systems for (transparent) truth and consequence.
- Such (substructural) theories stand out among their fellow villains:

- I am presenting a cluster of nonclassical systems for (transparent) truth and consequence.
- Such (substructural) theories stand out among their fellow villains:
  - Unlike other substructural options, they have a nice semantics

- I am presenting a cluster of nonclassical systems for (transparent) truth and consequence.
- Such (substructural) theories stand out among their fellow villains:
  - Unlike other substructural options, they have a nice semantics
  - Unlike fully structural options, they have a nice proof theory.

*theories*

---





- $\mathcal{L}_C$  results from expanding a first-order language  $\mathcal{L}$  with the **binary** predicate  $C$ .

- $\mathcal{L}_C$  results from expanding a first-order language  $\mathcal{L}$  with the **binary** predicate  $C$ .
- $\Gamma, \Delta, \Theta, \Lambda, \dots$  are multisets of formulae of  $\mathcal{L}_C$ .

- $\mathcal{L}_C$  results from expanding a first-order language  $\mathcal{L}$  with the **binary** predicate  $C$ .
- $\Gamma, \Delta, \Theta, \Lambda, \dots$  are multisets of formulae of  $\mathcal{L}_C$ .
- $\ulcorner \cdot \urcorner$  *somehow* yields a name for each sentence of the language.

- $\mathcal{L}_C$  results from expanding a first-order language  $\mathcal{L}$  with the **binary** predicate  $C$ .
- $\Gamma, \Delta, \Theta, \Lambda, \dots$  are multisets of formulae of  $\mathcal{L}_C$ .
- $\ulcorner \cdot \urcorner$  *somehow* yields a name for each sentence of the language.
- on occasion I will work with  $\mathcal{L}_{\mathbb{N}}$ , which yields a proper way of dealing with formal syntax

$$(\text{ref}^-) \quad \begin{array}{c} \Gamma, A \Rightarrow A, \Delta \\ \text{with } A \in \text{AtFml}_{\mathcal{L}} \end{array}$$

$$(\text{cut}) \quad \frac{\Gamma \Rightarrow \Delta, A \quad A, \Gamma \Rightarrow \Delta}{\Gamma \Rightarrow \Delta}$$

$$(\neg l) \quad \frac{\Gamma \Rightarrow A, \Delta}{\Gamma, \neg A \Rightarrow \Delta}$$

$$(\neg r) \quad \frac{\Gamma, A \Rightarrow \Delta}{\Gamma \Rightarrow \neg A, \Delta}$$

$$(\wedge l) \quad \frac{\Gamma, A, B \Rightarrow \Delta}{\Gamma, A \wedge B \Rightarrow \Delta}$$

$$(\wedge r) \quad \frac{\Gamma \Rightarrow A, \Delta \quad \Gamma \Rightarrow B, \Delta}{\Gamma \Rightarrow \Delta, A \wedge B}$$

$$(\forall l) \quad \frac{\Gamma, \forall x A, A(s) \Rightarrow \Delta}{\Gamma, \forall x A \Rightarrow \Delta}$$

$$(\forall r) \quad \frac{\Gamma \Rightarrow A(y), \Delta}{\Gamma \Rightarrow \Delta, \forall x A} \quad y \notin \text{FV}(\Gamma, \Delta, \forall x A)$$

$$(\text{ref}^-) \quad \frac{\Gamma, A \Rightarrow A, \Delta}{\text{with } A \in \text{AtFml}_{\mathcal{L}}}$$

$$(\text{cut}) \quad \frac{\Gamma \Rightarrow \Delta, A \quad A, \Gamma \Rightarrow \Delta}{\Gamma \Rightarrow \Delta}$$

$$(\neg l) \quad \frac{\Gamma \Rightarrow A, \Delta}{\Gamma, \neg A \Rightarrow \Delta}$$

$$(\neg r) \quad \frac{\Gamma, A \Rightarrow \Delta}{\Gamma \Rightarrow \neg A, \Delta}$$

$$(\wedge l) \quad \frac{\Gamma, A, B \Rightarrow \Delta}{\Gamma, A \wedge B \Rightarrow \Delta}$$

$$(\wedge r) \quad \frac{\Gamma \Rightarrow A, \Delta \quad \Gamma \Rightarrow B, \Delta}{\Gamma \Rightarrow \Delta, A \wedge B}$$

$$(\forall l) \quad \frac{\Gamma, \forall x A, A(s) \Rightarrow \Delta}{\Gamma, \forall x A \Rightarrow \Delta}$$

$$(\forall r) \quad \frac{\Gamma \Rightarrow A(y), \Delta}{\Gamma \Rightarrow \Delta, \forall x A} \quad y \notin \text{FV}(\Gamma, \Delta, \forall x A)$$

## Observation

This logic is the substructural ‘dual’ of **FDE**:  $\Gamma \Rightarrow \Delta / \Gamma' \Rightarrow \Delta'$  is admissible iff  $\bigwedge \Gamma \rightarrow \bigvee \Delta \Rightarrow \bigwedge \Gamma' \rightarrow \bigvee \Delta'$  is FDE-valid.

$$(Cl) \frac{\Gamma \Rightarrow \Delta, A \quad B, \Gamma \Rightarrow \Delta}{\Gamma, C(\ulcorner A \urcorner, \ulcorner B \urcorner) \Rightarrow \Delta}$$

$$(Cr) \frac{\Gamma, A \Rightarrow B, \Delta}{\Gamma \Rightarrow C(\ulcorner A \urcorner, \ulcorner B \urcorner), \Delta}$$

$$\text{(Cl)} \frac{\Gamma \Rightarrow \Delta, A \quad B, \Gamma \Rightarrow \Delta}{\Gamma, C(\ulcorner A \urcorner, \ulcorner B \urcorner) \Rightarrow \Delta} \quad \text{(Cr)} \frac{\Gamma, A \Rightarrow B, \Delta}{\Gamma \Rightarrow C(\ulcorner A \urcorner, \ulcorner B \urcorner), \Delta}$$

... from which one can easily obtain (assuming  $\Gamma \Rightarrow \Delta, \top$ ):

$$\text{(Tr l)} \frac{A, \Gamma \Rightarrow \Delta}{\text{Tr } \ulcorner A \urcorner, \Gamma \Rightarrow \Delta} \quad \text{(Tr r)} \frac{\Gamma \Rightarrow \Delta, A}{\Gamma \Rightarrow \Delta, \text{Tr } \ulcorner A \urcorner}$$



$$\Gamma \Rightarrow \Delta, \varphi$$

for  $\varphi$  an axiom of PA in  $\mathcal{L}_{\mathbb{N}}$

$$(=\text{r}) \frac{\Gamma \Rightarrow s^{\circ} = t^{\circ}, \Delta}{\Gamma \Rightarrow \text{Tr}(s = t), \Delta}$$

$$(=\text{l}) \frac{\Gamma, s^{\circ} = t^{\circ} \Rightarrow \Delta}{\Gamma, \text{Tr}(s = t) \Rightarrow \Delta}$$

$$(\text{Tr } 1) \frac{\Gamma \Rightarrow \text{Tr}(x), \Delta}{\Gamma \Rightarrow \text{Tr}^{\ulcorner} \text{Tr}(\dot{x})^{\urcorner}, \Delta}$$

$$(\text{Tr } 2) \frac{\Gamma, \text{Tr}(x) \Rightarrow \Delta}{\Gamma, \text{Tr}^{\ulcorner} \text{Tr}(\dot{x})^{\urcorner} \Rightarrow \Delta}$$

$$(\text{Tr } \neg 1) \frac{\Gamma \Rightarrow \neg \text{Tr}(\sigma), \Delta}{\Gamma \Rightarrow \text{Tr}(\neg \sigma), \Delta}$$

$$(\text{Tr } \neg 2) \frac{\Gamma, \neg \text{Tr}(\sigma) \Rightarrow \Delta}{\Gamma, \text{Tr}(\neg \sigma), \Delta}$$

$$(\text{Tr } \wedge 1) \frac{\Gamma \Rightarrow \text{Tr}(\sigma), \text{Tr}(\tau), \Delta}{\Gamma \Rightarrow \text{Tr}(\sigma \wedge \tau) \Rightarrow \Delta}$$

$$(\text{Tr } \wedge 2) \frac{\Gamma, \text{Tr}(\sigma), \text{Tr}(\tau) \Rightarrow \Delta}{\Gamma, \text{Tr}(\sigma \wedge \tau) \Rightarrow \Delta}$$

$$(\text{Tr } \forall 1) \frac{\Gamma \Rightarrow \forall t \text{Tr}(\sigma(t/v)), \Delta}{\Gamma \Rightarrow \text{Tr}(\forall v \sigma), \Delta}$$

$$(\text{Tr } \forall 2) \frac{\Gamma, \forall t \text{Tr}(\sigma(t/v)) \Rightarrow \Delta}{\Gamma, \text{Tr}(\forall v \sigma) \Rightarrow \Delta}$$

It is possible to formulate compositional extensions of **LPC**, e.g.:

$$\frac{\Gamma, C(\sigma, \delta) \Rightarrow \Delta \quad \Gamma, C(\tau, \delta) \Rightarrow \Delta}{\Gamma, C(\sigma \wedge \tau, \delta) \Rightarrow \Delta}$$

$$\frac{\Gamma \Rightarrow C(\sigma, \delta), C(\tau, \delta), \Delta}{\Gamma \Rightarrow C(\sigma \wedge \tau, \delta), \Delta}$$

$$\frac{\Gamma, C(\gamma, \sigma), C(\gamma, \tau) \Rightarrow \Delta}{\Gamma, C(\gamma, \sigma \wedge \tau) \Rightarrow \Delta}$$

$$\frac{\Gamma \Rightarrow C(\gamma, \sigma), C(\gamma, \tau), \Delta}{\Gamma \Rightarrow C(\gamma, \sigma \wedge \tau), \Delta}$$

*semantics*

---

## Nicolai&Rossi 2018

LPC – and its compositional extension – has a natural model  $(\mathbb{N}, X_\Phi)$  where  $X_\Phi$  is a fixed point of a monotone operator  $\Phi$  such that:

## Nicolai&Rossi 2018

LPC – and its compositional extension – has a natural model  $(\mathbb{N}, X_\Phi)$  where  $X_\Phi$  is a fixed point of a monotone operator  $\Phi$  such that:

- $\Phi$  operates on subsets of the set of (codes) of all sequents

## Nicolai&Rossi 2018

LPC – and its compositional extension – has a natural model  $(\mathbb{N}, X_\Phi)$  where  $X_\Phi$  is a fixed point of a monotone operator  $\Phi$  such that:

- $\Phi$  operates on subsets of the set of (codes) of all sequents
- for any such  $X$ :

## Nicolai&amp;Rossi 2018

LPC – and its compositional extension – has a natural model  $(\mathbb{N}, X_\Phi)$  where  $X_\Phi$  is a fixed point of a monotone operator  $\Phi$  such that:

- $\Phi$  operates on subsets of the set of (codes) of all sequents
- for any such  $X$ :
  - $\#(\Gamma \Rightarrow s = t, \Delta) \in \Phi(X)$ , for  $\mathbb{N} \models s = t$

## Nicolai&amp;Rossi 2018

LPC – and its compositional extension – has a natural model  $(\mathbb{N}, X_\Phi)$  where  $X_\Phi$  is a fixed point of a monotone operator  $\Phi$  such that:

- $\Phi$  operates on subsets of the set of (codes) of all sequents
- for any such  $X$ :
  - $\#(\Gamma \Rightarrow s = t, \Delta) \in \Phi(X)$ , for  $\mathbb{N} \models s = t$
  - $\#(\Gamma \Rightarrow C(\ulcorner A \urcorner, \ulcorner B \urcorner), \Delta) \in \Phi(X)$  if  $\#(\Gamma, A \Rightarrow B, \Delta) \in X$



## Nicolai&amp;Rossi 2018

LPC – and its compositional extension – has a natural model  $(\mathbb{N}, X_\Phi)$  where  $X_\Phi$  is a fixed point of a monotone operator  $\Phi$  such that:

- $\Phi$  operates on subsets of the set of (codes) of all sequents
- for any such  $X$ :
  - $\#(\Gamma \Rightarrow s = t, \Delta) \in \Phi(X)$ , for  $\mathbb{N} \models s = t$
  - $\#(\Gamma \Rightarrow C(\ulcorner A \urcorner, \ulcorner B \urcorner), \Delta) \in \Phi(X)$  if  $\#(\Gamma, A \Rightarrow B, \Delta) \in X$
  - $\#(\Gamma, C(\ulcorner A \urcorner, \ulcorner B \urcorner) \Rightarrow \Delta) \in \Phi(X)$  if  $\#(\Gamma \Rightarrow A, \Delta) \in X$  and  $\#(\Gamma, B \Rightarrow \Delta) \in X \dots$

## Nicolai&amp;Rossi 2018

LPC – and its compositional extension – has a natural model  $(\mathbb{N}, X_\Phi)$  where  $X_\Phi$  is a fixed point of a monotone operator  $\Phi$  such that:

- $\Phi$  operates on subsets of the set of (codes) of all sequents
- for any such  $X$ :
  - $\#(\Gamma \Rightarrow s = t, \Delta) \in \Phi(X)$ , for  $\mathbb{N} \models s = t$
  - $\#(\Gamma \Rightarrow C(\ulcorner A \urcorner, \ulcorner B \urcorner), \Delta) \in \Phi(X)$  if  $\#(\Gamma, A \Rightarrow B, \Delta) \in X$
  - $\#(\Gamma, C(\ulcorner A \urcorner, \ulcorner B \urcorner) \Rightarrow \Delta) \in \Phi(X)$  if  $\#(\Gamma \Rightarrow A, \Delta) \in X$  and  $\#(\Gamma, B \Rightarrow \Delta) \in X \dots$
- with  $(\mathbb{N}, X) \models^* \Gamma \Rightarrow \Delta$  iff all  $\gamma \in \Gamma$  are **true/both/neither** only if there is a  $\delta \in \Delta$  that is **true/both**, we have:  
 $\#(\Gamma, A \Rightarrow B, \Delta) \in X$  **iff**  $\#(\Gamma \Rightarrow C(\ulcorner A \urcorner, \ulcorner B \urcorner), \Delta) \in X$ .

## Nicolai&amp;Rossi 2018

LPC – and its compositional extension – has a natural model  $(\mathbb{N}, X_\Phi)$  where  $X_\Phi$  is a fixed point of a monotone operator  $\Phi$  such that:

- $\Phi$  operates on subsets of the set of (codes) of all sequents
- for any such  $X$ :
  - $\#(\Gamma \Rightarrow s = t, \Delta) \in \Phi(X)$ , for  $\mathbb{N} \models s = t$
  - $\#(\Gamma \Rightarrow C(\ulcorner A \urcorner, \ulcorner B \urcorner), \Delta) \in \Phi(X)$  if  $\#(\Gamma, A \Rightarrow B, \Delta) \in X$
  - $\#(\Gamma, C(\ulcorner A \urcorner, \ulcorner B \urcorner) \Rightarrow \Delta) \in \Phi(X)$  if  $\#(\Gamma \Rightarrow A, \Delta) \in X$  and  $\#(\Gamma, B \Rightarrow \Delta) \in X \dots$
- with  $(\mathbb{N}, X) \models^* \Gamma \Rightarrow \Delta$  iff all  $\gamma \in \Gamma$  are **true/both/neither** only if there is a  $\delta \in \Delta$  that is **true/both**, we have:
 
$$\#(\Gamma, A \Rightarrow B, \Delta) \in X \text{ iff } \#(\Gamma \Rightarrow C(\ulcorner A \urcorner, \ulcorner B \urcorner), \Delta) \in X.$$

## Observation

$(\mathbb{N}, X_{\Phi \upharpoonright \text{Tr}}) \models \text{KF} \upharpoonright$  iff  $(\mathbb{N}, X_{\Phi \upharpoonright \text{Tr}}) \models \text{PKF} \upharpoonright$  iff  $(\mathbb{N}, X_{\Phi \upharpoonright \text{Tr}}) \models^* \text{RKF} \upharpoonright$

**LPC<sup>∞</sup>** is obtained from LPC in  $\mathcal{L}_{\mathbb{N}}$  by omitting free variables and:

- adding axioms  $\Gamma \Rightarrow r = s, \Delta$  and  $\Gamma, r = s \Rightarrow \Delta$ ; where  $r, s$  are closed terms of  $\mathcal{L}_{\mathbb{C}}^{\mathbb{N}}$  and, respectively,  $r^{\mathbb{N}} = s^{\mathbb{N}}$  and  $r^{\mathbb{N}} \neq s^{\mathbb{N}}$ .
- Replacing  $(\forall r)$  with:

$$\frac{\Gamma \Rightarrow \varphi(s), \Delta \quad \text{for any closed term } s}{\Gamma \Rightarrow \forall x \varphi(x), \Delta} (\omega)$$

$\text{LPC}^\infty$  is obtained from LPC in  $\mathcal{L}_\mathbb{N}$  by omitting free variables and:

- adding axioms  $\Gamma \Rightarrow r = s, \Delta$  and  $\Gamma, r = s \Rightarrow \Delta$ ; where  $r, s$  are closed terms of  $\mathcal{L}_\mathbb{C}^\mathbb{N}$  and, respectively,  $r^\mathbb{N} = s^\mathbb{N}$  and  $r^\mathbb{N} \neq s^\mathbb{N}$ .
- Replacing  $(\forall r)$  with:

$$\frac{\Gamma \Rightarrow \varphi(s), \Delta \quad \text{for any closed term } s}{\Gamma \Rightarrow \forall x \varphi(x), \Delta} (\omega)$$

### Proposition

For  $\mathcal{I}_\Phi$  the  $(\Pi_1^1\text{-complete})$  minimal fixed point of  $\Phi$ , one has (by employing cut-elimination):

$$A \Rightarrow B \in \mathcal{I}_\Phi \quad \text{iff} \quad \vdash_{\text{LPC}^\infty}^\alpha C(\ulcorner A \urcorner, \ulcorner B \urcorner) \text{ for some } \alpha < \omega_1^{\text{ck}}$$

*proof theory*

---



- Cut-elimination for theories with non-logical rules such as (CI) and (Cr) is a piece of cake for non-contractive systems ([Cantini 2003](#), [Grišin 1982](#)): derivation trees don't 'grow' in the reduction, so one can simply count the applications of Tr- or C-rules.



- Cut-elimination for theories with non-logical rules such as (CI) and (Cr) is a piece of cake for non-contractive systems (Cantini 2003, Grišin 1982): derivation trees don't 'grow' in the reduction, so one can simply count the applications of Tr- or C-rules.
- When contraction is admissible, one has to find finer grained measures, and the task becomes notoriously hard. To my knowledge, there is no **simple** reduction strategy for fully structural, transparent theories of truth (and consequence), either for full or partial cut elimination. But...

- Cut-elimination for theories with non-logical rules such as (CI) and (Cr) is a piece of cake for non-contractive systems (Cantini 2003, Grišin 1982): derivation trees don't 'grow' in the reduction, so one can simply count the applications of Tr- or C-rules.
- When contraction is admissible, one has to find finer grained measures, and the task becomes notoriously hard. To my knowledge, there is no **simple** reduction strategy for fully structural, transparent theories of truth (and consequence), either for full or partial cut elimination. But...
- ... there is one for LPC (and extensions thereof), and RKF<sub>↑</sub>.



One defines the C-complexity of a formula in a derivation  $\mathcal{D}$ :

One defines the C-complexity of a formula in a derivation  $\mathcal{D}$ :

- formulae of  $\mathcal{L}$  have C-complexity 0 in any  $\mathcal{D}$ ;

One defines the C-complexity of a formula in a derivation  $\mathcal{D}$ :

- formulae of  $\mathcal{L}$  have C-complexity 0 in any  $\mathcal{D}$ ;
- In initial sequents  $\Gamma, \varphi \Rightarrow \varphi, \Delta$ , each formula  $\psi$  is such that  $\kappa(\psi) = 0$ .

One defines the C-complexity of a formula in a derivation  $\mathcal{D}$ :

- formulae of  $\mathcal{L}$  have C-complexity 0 in any  $\mathcal{D}$ ;
- In initial sequents  $\Gamma, \varphi \Rightarrow \varphi, \Delta$ , each formula  $\psi$  is such that  $\kappa(\psi) = 0$ .
- If  $\mathcal{D}$  ends with

$$\frac{\Gamma, A \Rightarrow B, \Delta}{\Gamma \Rightarrow C(\ulcorner A \urcorner, \ulcorner B \urcorner), \Delta}$$

then  $\kappa(C(\ulcorner A \urcorner, \ulcorner B \urcorner)) = \max(\kappa(A), \kappa(B)) + 1$  and the C-complexity of the formulas in  $\Gamma, \Delta$  is unchanged.

One defines the C-complexity of a formula in a derivation  $\mathcal{D}$ :

- formulae of  $\mathcal{L}$  have C-complexity 0 in any  $\mathcal{D}$ ;
- In initial sequents  $\Gamma, \varphi \Rightarrow \varphi, \Delta$ , each formula  $\psi$  is such that  $\kappa(\psi) = 0$ .
- If  $\mathcal{D}$  ends with

$$\frac{\Gamma, A \Rightarrow B, \Delta}{\Gamma \Rightarrow C(\ulcorner A \urcorner, \ulcorner B \urcorner), \Delta}$$

then  $\kappa(C(\ulcorner A \urcorner, \ulcorner B \urcorner)) = \max(\kappa(A), \kappa(B)) + 1$  and the C-complexity of the formulas in  $\Gamma, \Delta$  is unchanged.

- If  $\mathcal{D}$  ends with

$$\frac{\gamma_1^1, \dots, \gamma_n^1 \Rightarrow \delta_1^1, \dots, \delta_m^1, A \quad B, \gamma_1^2, \dots, \gamma_n^2 \Rightarrow \delta_1^2, \dots, \delta_m^2}{\gamma_1^3, \dots, \gamma_n^3, C(\ulcorner A \urcorner, \ulcorner B \urcorner) \Rightarrow \delta_1^3, \dots, \delta_m^3}$$

then

$$\kappa(C(\ulcorner A \urcorner, \ulcorner B \urcorner)) = \max(\kappa(A), \kappa(B)) + 1$$

$$\kappa(\gamma_i^3) = \max(\kappa(\gamma_i^1), \kappa(\gamma_i^2)) \quad 1 \leq i \leq n$$

$$\kappa(\delta_j^3) = \max(\kappa(\delta_j^1), \kappa(\delta_j^2)) \quad 1 \leq j \leq m$$



### Lemma (Inversion)

*All rules of LPC are  $\kappa$ -invertible: the C-complexity of the complex formulae is **no greater** than the one of their constituents, and **strictly smaller** for  $\kappa > 0$ .*

## Lemma (Inversion)

*All rules of LPC are  $\kappa$ -invertible: the C-complexity of the complex formulae is **no greater** than the one of their constituents, and **strictly smaller** for  $\kappa > 0$ .*

**For instance:** If  $\mathcal{D} \vdash_{\text{LPC}} \Gamma^1, C(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \Rightarrow \Delta^1$ , then there are  $\mathcal{D}' \vdash_{\text{LPC}} \Gamma^{10} \Rightarrow \Delta^{10}, \varphi$  and  $\mathcal{D}'' \vdash_{\text{LPC}} \psi, \Gamma^{11} \Rightarrow \Delta^{11}$  with  $\kappa(\Gamma^{10}), \kappa(\Gamma^{11}) \leq \kappa(\Gamma^1)$ , with  $\kappa(\Delta^{10}), \kappa(\Delta^{11}) \leq \kappa(\Delta^1)$ , and

$$\begin{aligned} \kappa(\varphi), \kappa(\psi) &\leq \kappa(C(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)), \text{ if } \kappa(C(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)) = 0, \text{ or} \\ \kappa(\varphi), \kappa(\psi) &< \kappa(C(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)), \text{ if } \kappa(C(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)) > 0. \end{aligned}$$

### Lemma (Inversion)

*All rules of LPC are  $\kappa$ -invertible: the C-complexity of the complex formulae is **no greater** than the one of their constituents, and **strictly smaller** for  $\kappa > 0$ .*

**For instance:** If  $\mathcal{D} \vdash_{\text{LPC}} \Gamma^1, C(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \Rightarrow \Delta^1$ , then there are  $\mathcal{D}' \vdash_{\text{LPC}} \Gamma^{10} \Rightarrow \Delta^{10}, \varphi$  and  $\mathcal{D}'' \vdash_{\text{LPC}} \psi, \Gamma^{11} \Rightarrow \Delta^{11}$  with  $\kappa(\Gamma^{10}), \kappa(\Gamma^{11}) \leq \kappa(\Gamma^1)$ , with  $\kappa(\Delta^{10}), \kappa(\Delta^{11}) \leq \kappa(\Delta^1)$ , and

$$\begin{aligned} \kappa(\varphi), \kappa(\psi) &\leq \kappa(C(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)), \text{ if } \kappa(C(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)) = 0, \text{ or} \\ \kappa(\varphi), \kappa(\psi) &< \kappa(C(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)), \text{ if } \kappa(C(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)) > 0. \end{aligned}$$

### Lemma

*All **truth rules** of  $\text{RKF} \upharpoonright$  are  $\kappa$ -invertible – again, with a strict inequality if the active formula has Tr-complexity  $> 0$ .*

Proposition ( $I\Delta_0 + \text{superexp}$ )

LPC admits full cut elimination.

Proposition ( $I\Delta_0 + \text{superexp}$ )

LPC admits full cut elimination.

*Proof strategy.* For  $\varphi$  the cut formula, the proof is by main induction on  $\kappa(\varphi)$ , and side inductions on the complexity of  $\varphi$  and on the level of the cut (i.e.  $d_0 + d_1$ ). This yields the usual superexponential bound.

### Proposition ( $I\Delta_0 + \text{superexp}$ )

LPC admits full cut elimination.

*Proof strategy.* For  $\varphi$  the cut formula, the proof is by main induction on  $\kappa(\varphi)$ , and side inductions on the complexity of  $\varphi$  and on the level of the cut (i.e.  $d_0 + d_1$ ). This yields the usual superexponential bound.

### Corollary

$\text{LPC}^\infty$  admits full cut elimination.

A **Tr-cut** is a cut on formulae of the form  $\text{Tr } t$ .

A **Tr-cut** is a cut on formulae of the form  $\text{Tr } t$ .

### Lemma (Tr-cut elimination)

*Let  $\mathcal{D}_0, \mathcal{D}_1$  be cut-free derivations, in  $\text{RKF}_\downarrow$ , of  $\Gamma^1 \Rightarrow \Delta^1, \text{Tr}(t)$  and  $\text{Tr}(t), \Gamma^2 \Rightarrow \Delta^2$  respectively. Then we can find a derivation  $\mathcal{D}$  of  $\Gamma \Rightarrow \Delta$  with  $\kappa(\Gamma^k) \leq \kappa(\Gamma)$  and  $\kappa(\Delta^k) \leq \tau(\Delta)$  and  $k \in \{1, 2\}$ .*



A **Tr-cut** is a cut on formulae of the form  $\text{Tr } t$ .

### Lemma (Tr-cut elimination)

Let  $\mathcal{D}_0, \mathcal{D}_1$  be cut-free derivations, in  $\text{RKF}\downarrow$ , of  $\Gamma^1 \Rightarrow \Delta^1, \text{Tr}(t)$  and  $\text{Tr}(t), \Gamma^2 \Rightarrow \Delta^2$  respectively. Then we can find a derivation  $\mathcal{D}$  of  $\Gamma \Rightarrow \Delta$  with  $\kappa(\Gamma^k) \leq \kappa(\Gamma)$  and  $\kappa(\Delta^k) \leq \tau(\Delta)$  and  $k \in \{1, 2\}$ .

### Corollary ( $\text{I}\Delta_0(\text{superexp})$ )

$\text{RKF}\downarrow$  is **conservative** over PA: if  $\Rightarrow \varphi$  is derivable in  $\text{RKF}\downarrow$  and  $\varphi \in \mathcal{L}_{\mathbb{N}}$ , then there is a derivation of  $\Rightarrow \varphi$  **without Tr-cuts**, i.e. a PA-derivation of  $\Rightarrow \varphi$ . A fortiori,  $\text{RKF}\downarrow$  is consistent if PA is.

*a story within the story*

---

- The strategy of removing **Tr-cuts** to obtain conservativity of truth theories originated in **Halbach 1999**.

- The strategy of removing **Tr-cuts** to obtain conservativity of truth theories originated in [Halbach 1999](#).
- A proof of the conservativeness of Tarskian (non self-referential) compositional clauses – a.k.a. **CT<sub>↑</sub>** – is presented in [Leigh 2015](#).

- The strategy of removing **Tr-cuts** to obtain conservativity of truth theories originated in [Halbach 1999](#).
- A proof of the conservativeness of Tarskian (non self-referential) compositional clauses – a.k.a. **CT<sub>↑</sub>** – is presented in [Leigh 2015](#).
- The version of **CT<sub>↑</sub>** with restricted initial sequents – a subtheory of **RKF<sub>↑</sub>** – admits, by contrast, a smooth cut-elimination procedure.

- The strategy of removing **Tr-cuts** to obtain conservativity of truth theories originated in **Halbach 1999**.
- A proof of the conservativeness of Tarskian (non self-referential) compositional clauses – a.k.a. **CT<sub>↑</sub>** – is presented in **Leigh 2015**.
- The version of **CT<sub>↑</sub>** with restricted initial sequents – a subtheory of **RKF<sub>↑</sub>** – admits, by contrast, a smooth cut-elimination procedure.
- What I presented for truth can be extended to a *compositional theory of consequence* in the style of **RKF<sub>↑</sub>**.

- I've presented a cluster of nonclassical systems for (transparent) truth and consequence.
- Such (substructural) theories stand out among their fellow villains:
  - Unlike other substructural options, they have a nice semantics
  - Unlike fully structural options, they have a nice proof theory.