

Copyright © 2017 Grupo Estudiantil de Matemática PUBLICADO POR GEM EDITORES WWW.FACEBOOK/GEMFCUNI Licensed under the Creative Commons Attribution-NonCommercial 3.0 Unported License (the "License"). You may not use this file except in compliance with the License. You may obtain a copy of the License at http://creativecommons.org/licenses/by-nc/3.0. Unless required by applicable law or agreed to in writing, software distributed under the License is distributed on an "AS IS" BASIS, WITHOUT WARRANTIES OR CONDITIONS OF ANY KIND, either express or implied. See the License for the specific language governing permissions and limitations under the License.

Primera impresión, abril 2017



-1	Primera parte	
1	Primera clase	. 7
1.1	La antiderivada o integral indefinida	7
1.2	Citas	7
1.3	Listas	7
1.3.1	Observación	
1.3.2 1.3.3	Bullet Points	
2	Segunda clase	. 9
2.1	Teoremas	9
2.1.1	Several equations	
2.1.2	Single Line	
2.2	Definiciones	9
2.3	Notaciones	10
2.4	Remarks	10
2.5	Corolarios	10
2.6	Proposiciones	10
2.6.1	Varias ecuaciones	
2.6.2	Single Line	10
2.7	Ejemplos	10
2.7.1	Ecuación y texto	11
2.7.2	Párrafo de texto	11

2.8	Ejercicios	11
2.9	Problemas	11
2.10	Vocabulario	11
Ш	Segunda parte	
3	Presentación de la información	15
3.1	Tablas	15
3.2	Figuras	15
	Bibliografía	17
	Libros	17
	Artículos	17

Primera parte

1.1 1.2 1.3	Primera clase La antiderivada o integral indefinida Citas Listas	7
2	Segunda clase	9
2.1	Teoremas	
2.2	Definiciones	
2.3	Notaciones	
2.4	Remarks	
2.5	Corolarios	
2.6	Proposiciones	
2.7	Ejemplos	
2.8	Ejercicios	
2.9	Problemas	
2.10	Vocabulario	



1.1 La antiderivada o integral indefinida

Definition 1.1.1 — Antiderivada. Sean $\mathscr I$ un intervalo, $f:\mathscr I\to\mathbb R$ una función y $F:\mathscr I\to\mathbb R$ otra función. Se dirá que F es una antiderivada de f si $F'(x)=f(x)\ \forall x\in\mathscr I$.

1.2 Citas

This statement requires citation [book_key]; this one is more specific [article_key].

1.3 Listas

Lists are useful to present information in a concise and/or ordered way¹.

1.3.1 Observación

- 1. F es derivable en \mathscr{I} .
- 2. F es continua en \mathscr{I} .

1.3.2 Bullet Points

• The first item

1.3.3 Descripciones y definiciones

Nombre Descripción Palabra Definición Comentario Elaboración

¹Footnote example...



2.1 Teoremas

This is an example of theorems.

2.1.1 Several equations

This is a theorem consisting of several equations.

Theorem 2.1.1 — Nombre del teorema. In $E = \mathbb{R}^n$ all norms are equivalent. It has the properties:

$$|||\mathbf{x}|| - ||\mathbf{y}||| \le ||\mathbf{x} - \mathbf{y}||$$
 (2.1)

$$\left|\left|\sum_{i=1}^{n} \mathbf{x}_{i}\right|\right| \leq \sum_{i=1}^{n} \left|\left|\mathbf{x}_{i}\right|\right| \quad \text{where } n \text{ is a finite integer}$$

$$(2.2)$$

2.1.2 Single Line

This is a theorem consisting of just one line.

Theorem 2.1.2 A set $\mathcal{D}(G)$ in dense in $L^2(G)$, $|\cdot|_0$.

2.2 Definiciones

This is an example of a definition. A definition could be mathematical or it could define a concept.

Definition 2.2.1 — Nombre de la definición. Given a vector space E, a norm on E is an

application, denoted $||\cdot||$, E in $\mathbb{R}^+ = [0, +\infty[$ such that:

$$||\mathbf{x}|| = 0 \Rightarrow \mathbf{x} = \mathbf{0} \tag{2.3}$$

$$||\mathbf{x}|| = 0 \Rightarrow \mathbf{x} = \mathbf{0}$$

$$||\lambda \mathbf{x}|| = |\lambda| \cdot ||\mathbf{x}||$$
(2.3)

$$||\mathbf{x} + \mathbf{y}|| \le ||\mathbf{x}|| + ||\mathbf{y}|| \tag{2.5}$$

2.3 Notaciones

Notation 2.1. Given an open subset G of \mathbb{R}^n , the set of functions φ are:

- 1. Bounded support G;
- 2. Infinitely differentiable;

a vector space is denoted by $\mathcal{D}(G)$.

2.4 Remarks

This is an example of a remark.



The concepts presented here are now in conventional employment in mathematics. Vector spaces are taken over the field $\mathbb{K} = \mathbb{R}$, however, established properties are easily extended to $\mathbb{K} = \mathbb{C}$.

2.5 Corolarios

This is an example of a corollary.

Corollary 2.5.1 — Nombre del corolario. The concepts presented here are now in conventional employment in mathematics. Vector spaces are taken over the field $\mathbb{K} = \mathbb{R}$, however, established properties are easily extended to $\mathbb{K} = \mathbb{C}$.

Proposiciones 2.6

This is an example of propositions.

2.6.1 Varias ecuaciones

Proposition 2.6.1 — PNombre de la proposición. It has the properties:

$$|||\mathbf{x}|| - ||\mathbf{y}||| \le ||\mathbf{x} - \mathbf{y}|| \tag{2.6}$$

$$\left|\left|\sum_{i=1}^{n} \mathbf{x}_{i}\right|\right| \leq \sum_{i=1}^{n} \left|\left|\mathbf{x}_{i}\right|\right| \quad \text{where } n \text{ is a finite integer}$$
(2.7)

2.6.2 Single Line

Proposition 2.6.2 Let $f, g \in L^2(G)$; if $\forall \varphi \in \mathcal{D}(G)$, $(f, \varphi)_0 = (g, \varphi)_0$ then f = g.

2.7 **Ejemplos**

This is an example of examples.

2.8 Ejercicios

2.7.1 Ecuación y texto

Example 2.1 Let $G = \{x \in \mathbb{R}^2 : |x| < 3\}$ and denoted by: $x^0 = (1,1)$; consider the function:

$$f(x) = \begin{cases} e^{|x|} & \text{si } |x - x^0| \le 1/2\\ 0 & \text{si } |x - x^0| > 1/2 \end{cases}$$
 (2.8)

The function f has bounded support, we can take $A = \{x \in \mathbb{R}^2 : |x - x^0| \le 1/2 + \varepsilon\}$ for all $\varepsilon \in]0; 5/2 - \sqrt{2}[$.

2.7.2 Párrafo de texto

■ Example 2.2 — Nombre ejemplo. Aeiou.

2.8 Ejercicios

This is an example of an exercise.

Exercise 2.1 This is a good place to ask a question to test learning progress or further cement ideas into students' minds.

2.9 Problemas

Problem 2.1 What is the average airspeed velocity of an unladen swallow?

2.10 Vocabulario

Define a word to improve a students' vocabulary.

Vocabulary 2.1 — Palabra. Definition of word.

Segunda parte

3	Presentación de la información	15
3.1	Tablas	
3.2	Figuras	
	Bibliografía	17
	Libros	

Artículos



3.1 Tablas

Treatments	Response 1	Response 2
Treatment 1	0.0003262	0.562
Treatment 2	0.0015681	0.910
Treatment 3	0.0009271	0.296

Table 3.1: Table caption

3.2 Figuras

Placeholder Image

Figure 3.1: Figure caption



Libros Artículos