

Chapter 1

Laplace Equation

If does not have boundary conditions, the problem is ill. The equation needs two conditions.

It is very easy to partition the interval $[a, b]$

Definition 1. We call $\Omega \subset \mathbb{R}^d$ for $d = 1, 2, 3$ a **domain** if

1. Ω is open, is connected. It has no holes. It must be smooth.

Definition 2. We call $\Gamma = \partial\Omega$ the **boundary** of the domain Ω .

Definition 3. By \hat{n} we denote the **unit normal vector** (facing outwards) on the boundary.

Definition 4. We define **function space of differentiable functions**

$$C^m * (\Omega) = \{f: \Omega \rightarrow \mathbb{R} | f(x_1, x_2, \dots, x_d)\}$$

Definition 5. We define **laplace operator**

- Let $f \in C^0(\Omega)$ be the **right hand side function**.
- Let $g \in C^0(\Gamma)$ be the **boundary value function**.
- **Dirichlet Problem** we are looking for $u \in C^2(\Omega)$ such that

$$-\Delta u = f \text{ in } \Omega$$

- Let Ω be the unit sphere

$$\Omega = \{\mathbf{x} = (x, y) \in \mathbb{R}^2 | x^2 + y^2 \leq 1\}$$

- Let $f = 1$ and $g = 0$
- There is **no solution** to the Dirichlet Problem

$$-\Delta u = 1 \text{ in } \Omega, u = 0 \text{ on } \Gamma.$$

which is 2 times differentiable.

1.1 The variational formulation

- Assume that $u \in C^2(\Omega)$ is a solution to the Laplace problem

$$-\Delta u(x, y) = f(x, y) \text{ in } \Omega \text{ with } u = 0 \text{ on } \Gamma.$$

- The, we can multiply this equation with a **test function** ϕ

$$-\Delta u(x, y) \cdot \phi(x, y) = f(x, y) \cdot \phi(x, y) \text{ in } \Omega$$

- The, we can **integrate by parts over the domain**

$$-\int_D \Delta u(x, y) \cdot \phi(x, y) dx dy = \int_{\Omega} f(x, y) \cdot \phi(x, y) dx dy$$

- We assume that the test function is differentiable $\phi \in C^1(\Omega)$. Then, we can **integrate by parts**

$$\int_{\Omega} \nabla u(x, y) \cdot \nabla \phi(x, y) dx dy - \int_{\Gamma} (\hat{n} \cdot \nabla) u \cdot \phi dS = \int_{\Omega} f(x, y) \cdot \phi(x, y) dx dy$$

- We assume that the test function is zero on the boundary. Then

$$\int_{\Omega} \nabla u(x, y) \cdot \nabla \phi(x, y) dx dy = \int_{\Omega} f(x, y) \cdot \phi(x, y) dx dy.$$

If the boundary is given by the graph of a function in C^2 , then there exists a classical solution $u \in C^2(\Omega)$.

- We introduce L^2 scalar product

$$(u, \phi) = \int_{\Omega}$$

Theorem 1. Let $\Omega \subset \mathbb{R}^d$ for $d = 1, 2, 3$ be a domain and $f \in L^2(\Omega)$. Then, there exists a solution

$$u \in \nu = H_0^1(\Omega)$$

to the Laplace problem in variational formulation

Chapter 2

Finite Element Method

Steps for a finite element discretization

1. We discretize the domain Ω by a mesh Ω_h .
2. On Ω_h we discretize the function space $\nu = H_0^1(\Omega)$ by a finite element space V_h .
3. We restrict the variational formulation to V_h

$$u_h \in V_h(\nabla u_h, \nabla \phi_h) = (f, \phi_h) \quad \forall \phi_h$$

4. We solve a linear system of equations.

2.1 Construction

- We discretize the domain Ω by splitting it into simple **open elements**, e.g, triangles, quadrilaterals (in $2D$) or tetrahedras, prisms, hexaedras, pyramids (in $3D$)
- The **finite element mesh** Ω_h

2.2 Some examples

2.3 Shape assumption

Local Finite Element space

- On every element $T \in \Omega_h$ define the basis functions of a simple polynomial space
- **bi-linear finite elements**
- Let T be a quadrilateral with the points $x^{(1)} = (0,0)$, $x^{(2)} = (h,0)$, $x^{(3)} = (0,h)$, $x^{(4)} = (h,h)$.

- $\phi^{(1)}(x, y) = (1 - \frac{x}{h})(1 - \frac{y}{h})$, $\phi^{(2)}(x, y) = \frac{x}{h}(1 - \frac{y}{h})$, $\phi^{(3)}(x, y) = (1 - \frac{x}{h})\frac{y}{h}$,
 $\phi^{(4)}(x, y) = \frac{xy}{h^2}$

- The Lagrange basis of the finite element space is given as

$$V_h = \{\phi_h \in C(\Omega) | \phi|_T \in Q^1 = \text{span}(\phi_h^{(1)}, \phi_h^{(2)}, \phi_h^{(3)}, \phi_h^{(4)})\}$$

- The **Lagrange basis** of **nodal basis** is given by

$$V$$

- Starting point: weak formulation of Laplace equation

$$u \in \nu$$

•

$$u_h \in V_h(\nabla u_h, \nabla \phi_h) = (f, \phi_h) \quad \forall \phi_h \in V_h$$

- The finite element is given by a local basis

$$V_h = \text{span}\{\phi_h^{(1)}, \dots, \phi_h^{(N)} \quad \forall i = 1, \dots, N\}$$

- We write the unknown solution $u_h \in V_h$

2.4 Assembling the matrix

- We must compute the matrix entries

$$A_{ij}(\nabla \phi_h^{(j)}, \nabla \phi_h^{(i)}) = \int_{\Omega} \nabla \phi_h^{(j)} \cdot \nabla \phi_h^{(i)} dx = \sum_{T \subset \Omega_h} \int_T \nabla \phi_h^{(j)} \cdot \nabla \phi_h^{(i)} dx$$

- For every **nodal**...

- We combine the result in a **stencil**

$$S = \begin{bmatrix} s_{31} & s_{32} & s_{33} \\ s_{21} & s_{22} & s_{23} \\ s_{11} & s_{12} & s_{13} \end{bmatrix}$$

- The finite element matrix on a small mesh with $16 = 4 \cdot 4$ nodes like

$$A = \frac{1}{3} [1]$$

- The main difference between 1D and 2D (or 3D)