

Math 214

VISHAL RAMAN

February 12, 2021

Contents

1	Topology	2
2	Topological Manifolds	3
3	Smooth Structures	3
4	Manifolds with Boundary	4
5	Smooth Maps	4

§1 Topology

Definition 1.1 (Topological Space). $(X, O_X \subset \mathcal{P}(X))$, where $A \in O_x$ are the open sets which satisfy the following:

1. $\emptyset, X \in O_X$.
2. $A, B \in O_X$ implies $A \cap B \in O_X$
3. $A_i \in O_X, i \in I$, then $\bigcup_{i \in I} A_i \in O_X$.

We say that $A \subset X$ is closed if $X \setminus A$ is open. $U \subset X$ is a neighborhood of $p \in X$ if $\exists A$ such that $p \in A \subset U$.

Definition 1.2. $\mathcal{B} \subset \mathcal{P}(X)$ is called a **basis** for the topology on X if for every subset $A \subset X$, A is open if and only if A is a union of elements of \mathcal{B} .

Let $(X, O_X), (Y, O_Y)$ be topological spaces.

Definition 1.3. A function $\varphi : X \rightarrow Y$ is continuous if for any open subset $B \subset Y$, $\varphi^{-1}(B) \subset X$ is open.

Definition 1.4. $\varphi : X \rightarrow Y$ is a homeomorphism if it is a continuous bijection whose inverse is continuous.

Definition 1.5. Let $Y \subset X$ a topological space. We set $O_Y = \{A \cap Y : A \in O_X\}$.

Definition 1.6. Given a topological space X , X is called locally Euclidean (of dimension n) at $p \in X$ if there is an open neighborhood about $p \in U \subset X$ that is homeomorphic to an open subset of \mathbb{R}^n .

Definition 1.7. A space X is **Hausdorff** if for any $p, q \in X, p \neq q$ there exists open subsets U, V with $p \in U, q \in V$ so that $U \cap V = \emptyset$.

Definition 1.8. $K \subset X$ is compact if every open cover of K has a finite subcover.

Given a topological space X , we have the following definitions:

Definition 1.9. X is connected if the only subsets that are open and closed are \emptyset, X .

Definition 1.10. A space is path-connected if for any $p, q \in X$ there is a continuous path between them.

Definition 1.11. An exhaustion by compact subsets is an increasing sequence of subsets $K_1 \subset K_2 \subset \dots \subset X$ such that K_i is compact and $K_i \subset \text{Int}(K_{i+1})$ and $\bigcup_i K_i = X$.

Definition 1.12. Take $\mathcal{U} \subset \mathcal{P}(X)$. This is a cover of X if $X = \bigcup_{U \in \mathcal{U}} U$. A collection is called locally finite if every $p \in X$ has a neighborhood $p \in W \subset X$ such that W only intersects finitely many $U \in \mathcal{U}$.


Definition 1.13. A collection of subsets \mathcal{V} is called a refinement of some other collection \mathcal{U} if for every $V \in \mathcal{V}$ there is $U \in \mathcal{U}$ such that $V \subset U$.

§2 Topological Manifolds

Definition 2.1. A topological space M is called an n -dimensional **topological manifold** if M satisfies the following:

- M is locally Euclidean at any point,
- M is Hausdorff,
- M is second countable.


Definition 2.2. A **coordinate chart** on M is a pair (U, φ) where $U \subset M$ is open and $\varphi : U \rightarrow \hat{U}$ is a homeomorphism to an open subset $\hat{U} \subset \mathbb{R}^n$.



Definition 2.3. X is called paracompact if every open cover has a locally finite refinement.

§3 Smooth Structures

Definition 3.1. Let M^n be a topological manifold. Two charts $(U, \varphi), (V, \psi)$ of M have a transition map: $\psi \circ \varphi^{-1}$. This map is a homeomorphism.



Definition 3.2. Two charts are smoothly compatible if the transition maps in both directions are smooth.

Definition 3.3. An atlas \mathcal{A} of M is a collection of charts such that the domains of the charts cover M . An atlas \mathcal{A} is smooth if any two charts in \mathcal{A} are smoothly compatible. An atlas \mathcal{A} is called a maximal smooth atlas on M if there is no smooth atlas containing \mathcal{A}' that contains \mathcal{A} .

Definition 3.4. A maximal smooth atlas \mathcal{A} on a topological manifold M is called a smooth structure on M .

Definition 3.5. A smooth manifold is a pair (M^n, \mathcal{A}) , where M^n is a topological manifold and \mathcal{A} is a smooth structure.

§4 Manifolds with Boundary

Definition 4.1. We denote $H^n = \{x^n \geq 0\} \subset \mathbb{R}^n$, the upper half space, the most basic example. Note that $\partial H^n = \{x^n = 0\} \cong \mathbb{R}^{n-1}$. The interior $\text{Int } H^n = \{x^n > 0\}$.

Definition 4.2. A topological manifold with boundary M^n is a topological space such that is Hausdorff, second countable, and every point $p \in H^n$ has an open neighborhood $p \in U \subset M$ that is homeomorphic to some (relatively) open subset $\widehat{U} \subset H$.

§5 Smooth Maps

Definition 5.1. $f : M \rightarrow \mathbb{R}^m$ is smooth if for every $p \in M$, there is a smooth chart (U, φ) , $\widehat{U} = \varphi(U)$ such that $p \in U$ and $\widehat{f} = f \circ \varphi^{-1} : \widehat{U} \rightarrow \mathbb{R}^m$ is smooth. We denote $C^\infty(M) : \{f : M \rightarrow \mathbb{R}^m \text{ smooth}\}$.

Definition 5.2. Suppose we have M^m, N^n smooth manifolds (with boundary) and take $F : M \rightarrow N$. F is called smooth if for any $p \in M$ there are smooth charts (U, φ) of M and (V, ψ) of N such that $p \in U$, $F(U) \subset V$ and $\psi \circ F \circ \varphi^{-1}$ is smooth.

Other equivalent definitions:

- For every $p \in M$, there exist smooth charts (U, φ) containing p and (V, ψ) containing $F(p)$ such that $U \cap F^{-1}(V)$ is open in M and the composite map $\psi \circ F \circ \varphi^{-1}$ is smooth from $\varphi(U \cap F^{-1}(V))$ to $\psi(V)$.
- F is continuous and there exist smooth atlases $\{(U_\alpha, \varphi_\alpha)\}$ and $\{(V_\beta, \psi_\beta)\}$ for M and N respectively so that for each α, β $\psi_\beta \circ F \circ \varphi_\alpha^{-1}$ is a smooth map from $\varphi_\alpha(U_\alpha \cap F^{-1}(V_\beta))$ to $\psi_\beta(V_\beta)$.

Definition 5.3. Let $\mathcal{X} = (X_\alpha)_{\alpha \in A}$ be an open cover of some topological space X . A partition of unity subordinate to \mathcal{X} is a family $(\psi_\alpha)_{\alpha \in A}$ of continuous maps on $\psi_\alpha : X \rightarrow \mathbb{R}$ such that $0 \leq \psi_\alpha \leq 1$, $\text{supp } \psi_\alpha \subset X_\alpha$, $(\text{supp } \psi_\alpha)_{\alpha \in A}$ is locally finite, and $\sum_{\alpha \in A} \psi_\alpha(x) = 1$ for all $x \in X$.

Definition 5.4. An open subset $B \subset m$ is called a regular coordinate ball if there is a smooth chart (U, φ) such that $\varphi(U) = B_{r'}(0)$, $\varphi(B) = B_r(0)$ where $0 < r < r'$.

Definition 5.5. If M is a topological space, $A \subset M$ is a closed subset, and $U \subset M$ is an open subset containing A , a continuous function $\psi : M \rightarrow \mathbb{R}$ is called a bump function for A supported in U if $0 \leq \psi \leq 1$ on M and $\psi \equiv 1$ on A , $\text{supp } \psi \subset U$.