

USAJMO 2011 - Problems and Solutions

VISHAL RAMAN

May 20, 2018

§1 Problems

Problem 1.1 (USAJMO 2011/1). Find, with proof, all positive integers n for which $2^n + 12^n + 2011^n$ is a perfect square.

Problem 1.2 (USAJMO 2011/2). Let a, b, c be positive real numbers such that $a^2 + b^2 + c^2 + (a + b + c)^2 \leq 4$. Prove that

$$\frac{ab+1}{(a+b)^2} + \frac{bc+1}{(b+c)^2} + \frac{ca+1}{(c+a)^2} \geq 3.$$

Problem 1.3 (USAJMO 2011/3). For a point $P = (a, a^2)$ in the coordinate plane, let $l(P)$ denote the line passing through P with slope $2a$. Consider the set of triangles with vertices of the form $P_1 = (a_1, a_1^2), P_2 = (a_2, a_2^2), P_3 = (a_3, a_3^2)$, such that the intersection of the lines $l(P_1), l(P_2), l(P_3)$ form an equilateral triangle \triangle . Find the locus of the center of \triangle as $P_1P_2P_3$ ranges over all such triangles.

Problem 1.4 (USAJMO 2011/4). A word is defined as any finite string of letters. A word is a palindrome if it reads the same backwards and forwards. Let a sequence of words W_0, W_1, W_2, \dots be defined as follows: $W_0 = a, W_1 = b$, and for $n \geq 2$, W_n is the word formed by writing W_{n-2} followed by W_{n-1} . Prove that for any $n \geq 1$, the word formed by writing $W_1, W_2, W_3, \dots, W_n$ in succession is a palindrome.

Problem 1.5 (USAJMO 2011/5). Points A, B, C, D, E lie on a circle ω and point P lies outside the circle. The given points are such that (i) lines PB and PD are tangent to ω , (ii) P, A, C are collinear, and (iii) $DE \parallel AC$. Prove that BE bisects AC .

Problem 1.6 (USAJMO 2011/6). Consider the assertion that for each positive integer $n \geq 2$, the remainder upon dividing 2^{2^n} by $2^n - 1$ is a power of 4. Either prove the assertion or find (with proof) a counterexample.

§2 Solutions

Problem 1 (USAJMO 2011/1)

Find, with proof, all positive integers n for which $2^n + 12^n + 2011^n$ is a perfect square.

Proof. The only possible value of n is $n = 1$. Note that $2^1 + 12^1 + 2011^1 = 2025 = 45^2$ is a perfect square.

Suppose there exists $m > 1$ so that $2^m + 12^m + 2011^m$ is a perfect square. Then

$$2^m + 12^m + 2011^m \equiv 2^m + (-1)^m \equiv (-1)^m \pmod{4}.$$

Since the quadratic residues of 4 are 0 and 1, m must be even so that $2^m + 12^m + 2011^m \equiv 1 \pmod{4}$. Furthermore,

$$2^m + 12^m + 2011^m \equiv (-1)^m + (1)^m \equiv 1 + (-1)^m \pmod{3}.$$

Since the quadratic residues of 3 are 0 and 1, m must be odd so that $2^m + 12^m + 2011^m \equiv 1 + (-1) \equiv 0 \pmod{3}$. However, m cannot be simultaneously odd and even. Therefore, there $n = 1$ is the only value of n so that $2^n + 12^n + 2011^n$ is a perfect square. \square

Problem 2 (USAJMO 2011/2)

Let a, b, c be positive real numbers such that $a^2 + b^2 + c^2 + (a + b + c)^2 \leq 4$. Prove that

$$\frac{ab+1}{(a+b)^2} + \frac{bc+1}{(b+c)^2} + \frac{ca+1}{(c+a)^2} \geq 3.$$

Proof. Note that

$$\begin{aligned} a^2 + b^2 + c^2 + (a + b + c)^2 &= 2a^2 + 2b^2 + 2c^2 + 2ab + 2bc + 2ca \\ &= (a^2 + 2ab + b^2) + (b^2 + 2bc + c^2) + (c^2 + 2ca + a^2) \\ &= (a + b)^2 + (b + c)^2 + (c + a)^2. \end{aligned}$$

Let $x = a + b, y = b + c, z = c + a$. Then the given condition is equivalent to $x^2 + y^2 + z^2 \leq 4$ which implies $4 - y^2 - z^2 \geq x^2$.

Also note that $a = \frac{x-y+z}{2}, b = \frac{x+y-z}{2}, c = \frac{-x+y+z}{2}$.

Thus, we have

$$\begin{aligned} \sum_{\text{cyc}} \frac{ab+1}{(a+b)^2} &= \sum_{\text{cyc}} \frac{4 + (x-y+z)(x+y-z)}{4x^2} \\ &= \sum_{\text{cyc}} \frac{4 + (x - (y-z))(x + (y-z))}{4x^2} \\ &= \sum_{\text{cyc}} \frac{4 + x^2 - (y-z)^2}{4x^2} \\ &= \sum_{\text{cyc}} \frac{4 - y^2 - z^2 + x^2 + 2yz}{4x^2} \\ &\geq \sum_{\text{cyc}} \frac{2x^2 + 2yz}{4x^2} \\ &= \sum_{\text{cyc}} \frac{1}{2} + \frac{1}{2} \frac{yz}{x^2} \\ &= \frac{3}{2} + \frac{1}{2} \sum_{\text{cyc}} \frac{yz}{x^2}, \end{aligned}$$

and

$$\sum_{\text{cyc}} \frac{yz}{x^2} = \sum_{\text{cyc}} \frac{(b+c)(c+a)}{(a+b)^2} \geq 3 \sqrt[3]{\prod_{\text{cyc}} \frac{(b+c)(c+a)}{(a+b)^2}} = 3,$$

from the AM-GM Inequality.

Therefore,

$$\sum_{\text{cyc}} \frac{ab+1}{(a+b)^2} \geq \frac{3}{2} + \frac{1}{2}(3) = 3,$$

as desired. □

Problem 3 (USAJMO 2011/3)

For a point $P = (a, a^2)$ in the coordinate plane, let $l(P)$ denote the line passing through P with slope $2a$. Consider the set of triangles with vertices of the form $P_1 = (a_1, a_1^2), P_2 = (a_2, a_2^2), P_3 = (a_3, a_3^2)$, such that the intersection of the lines $l(P_1), l(P_2), l(P_3)$ form an equilateral triangle \triangle . Find the locus of the center of \triangle as $P_1P_2P_3$ ranges over all such triangles.

Proof. Note that $l(P) = a^2 + 2a(x - a) = 2ax - a^2$.

Therefore, we have

$$l(P_1) = 2a_1x - a_1^2,$$

$$l(P_2) = 2a_2x - a_2^2,$$

$$l(P_3) = 2a_3x - a_3^2.$$

Let $Q_1 = l(P_1) \cap l(P_2), Q_2 = l(P_2) \cap l(P_3), Q_3 = l(P_3) \cap l(P_1)$. Solving for the intersection points, we have

$$Q_1 = \left(\frac{a+b}{2}, ab \right), Q_2 = \left(\frac{b+c}{2}, bc \right), Q_3 = \left(\frac{c+a}{2}, ca \right).$$

Denote H as the orthocenter and G as the centroid of $\triangle Q_1Q_2Q_3$. If $\triangle Q_1Q_2Q_3$ is equilateral, then $H = G$.

Firstly,

$$G = \left(\frac{a+b+c}{3}, \frac{ab+bc+ca}{3} \right),$$

the average of the coordinates.

Let $l(H_3)$ and $l(H_1)$ denote the altitudes from Q_3 and Q_1 respectively. The slope of $\overline{Q_1Q_2} = \frac{a_1a_2 - a_2a_3}{\frac{a_1+a_2}{2} - \frac{a_2+a_3}{2}} = 2a_2$, which means the slope of $l(H_3)$ is $-\frac{1}{2a_2}$, which gives

$$l(H_3) = a_1a_3 - \frac{1}{2a_2} \left(x - \frac{a_1+a_3}{2} \right),$$

and similarly,

$$l(H_1) = a_1a_2 - \frac{1}{2a_3} \left(x - \frac{a_1+a_2}{2} \right).$$

Then,

$$H = l(H_3) \cap l(H_1) = \left(\frac{a_1+a_2+a_3+4a_1a_2a_3}{2}, -\frac{1}{4} \right).$$

Call the x and y coordinates of G , x and y respectively. Since $G = H$, $x = \frac{a_1+a_2+a_3}{3} = \frac{a_1+a_2+a_3+4a_1a_2a_3}{2}$, which gives $x = -4a_1a_2a_3$.

Also, $y = \frac{a_1a_2+a_2a_3+a_3a_1}{3} = -\frac{1}{4}$, so $-12a_1y = -4a_1^2a_2 - 4a_1a_2a_3 - 4a_1^2a_3 = 3a_1$, and since $-4a_1a_2a_3 = x$, we find $x(a_1) = \frac{3a_1-4a_1^3}{1-12a_1^2}$, where $x(a_1)$ denotes x as a function of $a_1 \in \mathbb{R} - \{-\frac{1}{\sqrt{12}}, \frac{1}{\sqrt{12}}\}$.

Note that $\lim_{a_1 \rightarrow -\frac{1}{\sqrt{12}}} x = -\infty$, and $\lim_{a_1 \rightarrow \frac{1}{\sqrt{12}}} x = \infty$. Therefore x takes on all reals in the interval $\left(-\frac{1}{\sqrt{12}}, \frac{1}{\sqrt{12}} \right)$. Therefore, the locus of centers of the equilateral triangles is the line $y = -\frac{1}{4}$.

□

Problem 4 (USAJMO 2011/4)

A word is defined as any finite string of letters. A word is a palindrome if it reads the same backwards and forwards. Let a sequence of words W_0, W_1, W_2, \dots be defined as follows: $W_0 = a, W_1 = b$, and for $n \geq 2$, W_n is the word formed by writing W_{n-2} followed by W_{n-1} . Prove that for any $n \geq 1$, the word formed by writing $W_1, W_2, W_3, \dots, W_n$ in succession is a palindrome.

Proof. We will use strong induction on n . We will let the AB represent the word formed by writing the words A and B in succession. From the problem statement, $W_{k+1} = W_k W_{k-1}$. Let $W(n) = W_1 W_2 \dots W_n$, and let $r(l_1 l_2 l_3 \dots l_n) = l_n \dots l_3 l_2 l_1$, where l_1, l_2, \dots, l_n are letters. Note that $r(XY) = r(Y)r(X)$ and $W(n) = W_1 W_2 \dots W_{n-2} W_{n-1} W_n = W(n-2)W_{n-1}W_n = W_{n-2}W_{n+1}$. It suffices to show $r(W(n)) = W(n)$ for all $n \in \mathbb{N}$.

First, note that $W(1) = W_1 = b$ is a palindrome. Also, $W(2) = W_1 W_2 = bab$ is a palindrome. Now, suppose $W(m) = r(W(m))$ for all $3 \leq m \leq k$. Then

$$\begin{aligned} r(W(k+1)) &= r(W(k)W_{k+1}) \\ &= r(W_{k+1})r(W(k)) \\ &= r(W_{k+1})W(k) \\ &= r(W_{k+1})W(k-2)W_{k+1}. \end{aligned}$$

So

$$\begin{aligned} r(r(W(k+1))) &= r(r(W_{k+1})W(k-2)W_{k+1}) \\ &= r(W_{k+1})r(W(k-2))r(r(W_{k+1})) \\ &= r(W_{k+1})W(k-2)W_{k+1}. \end{aligned}$$

But $r(r(W(k+1))) = W(k+1)$. Therefore, $W(k+1) = r(W(k+1))$, which means $W(n)$ is a palindrome for all $n \in \mathbb{N}$. \square

Problem 5 (USAJMO 2011/5)

Points A, B, C, D, E lie on a circle ω and point P lies outside the circle. The given points are such that (i) lines PB and PD are tangent to ω , (ii) P, A, C are collinear, and (iii) $DE \parallel AC$. Prove that BE bisects AC .

Proof. Let \angle denote directed angles (mod 180°). Let O be the center of the circle, and let $F = BE \cap AC$.

Since PB and PD are tangents, $OB \perp BP$ and $OD \perp DP$, so $OBPD$ is cyclic.

Now,

$$\angle BOP = \angle BDP = \angle BAD = \angle BED = \angle BFP,$$

where the last equality follows from $DE \parallel AC$, and the others follow from the Inscribed Angles Theorem. Since $\angle BOP = \angle BFP$, $BOFP$ is cyclic, which means $OF \perp AC$. Since $OA = OC$, we must have $AF = FC$. Therefore, BE bisects AC . □

Problem 6 (USAJMO 2011/6)

Consider the assertion that for each positive integer $n \geq 2$, the remainder upon dividing 2^{2^n} by $2^n - 1$ is a power of 4. Either prove the assertion or find (with proof) a counterexample.

Proof. Let $r(a, b)$ denote the remainder upon dividing a by b .

From long division,

$$2^{2^n} = (2^n - 1)(2^{2^n-n} + 2^{2^n-2n} + \dots + 2^{2^n-(k-1)n}) + 2^{2^n-kn},$$

where k is the largest integer so that $2^n - kn \geq 0$. In other words, $2^n - kn = r(2^n, n)$. Thus, $r(2^{2^n}, n) = 2^{r(2^n, n)}$.

If n is even, $r(2^n, n)$ is clearly even, so $r(2^{2^n}, n)$ will be a power of 4.

Thus, it suffices to check odd composite numbers. One quickly finds a counterexample $n = 25$, since $r(2^{25}, 25) = 7$, so $r(2^{2^n}) = 2^7$, which is not a power of 4.

□