

# **Math 222a Lecture Notes, Fall 2020**

## **Partial Differential Equations**

Professor: Daniel Tataru

Vishal Raman

# Contents

<b>1 September 1st, 2020</b>	<b>3</b>
1.1 Introduction . . . . .	3
1.2 Classification of PDE's . . . . .	3
1.3 Initial Value Problems . . . . .	4
1.4 Boundary Value Problems . . . . .	5
1.5 Fluid Classification . . . . .	5
<b>2 September 3rd, 2020</b>	<b>7</b>
2.1 Picard-Lindeloff Theorem . . . . .	7
2.2 Contraction Principle . . . . .	7
2.3 Bootstrap Argument . . . . .	9

## §1 September 1st, 2020

### §1.1 Introduction

Partial differential equations apply to functions  $u : \mathbb{R}^n \rightarrow \mathbb{R}(\mathbb{C})$ , where  $u$  refers to the space dimension. Usually,  $n \geq 2$  ( $n = 1$  corresponds to ODEs).

We present the following notation:

- $\frac{\partial}{\partial x_i} u = \partial_i u$
- There is also multi-index notation, where  $\alpha = (\alpha_1, \dots, \alpha_n)$  and  $\partial^\alpha u = \partial_1^{\alpha_1} \partial_2^{\alpha_2} \dots \partial_n^{\alpha_n} u$ . The size of  $\alpha$  is given by  $|\alpha| = \sum_{i=1}^n \alpha_i$ .
- $C(\mathbb{R}^n)$ , continuous functions in  $\mathbb{R}^n$ .
- $C(\Omega)$ ,  $\Omega \subset \mathbb{R}^n$ , continuous functions in  $\Omega$ .
- $C^1(\mathbb{R}^n)$ ,  $C^1(\Omega)$ , continuously differentiable functions.
- $C^k(\mathbb{R}^n)$ ,  $C^k(\Omega)$ ,  $k$ -times differentiable.
- $C^\infty(\mathbb{R}^n) = \bigcap_{k=0}^\infty C^k(\mathbb{R}^n)$ .

We consider an example PDE,

$$F(u, \partial u, \partial^2 u, \dots, \partial^k u) = 0.$$

In the above,  $k \geq 1$  and  $k$  is the **order** of the equation. We also have the shorthand  $F(\partial^{\leq k} u) = 0$ .

### §1.2 Classification of PDE's

**Definition 1.1** (Linear PDE). The PDE is a linear function of its arguments. We can apply multi-index notation, as follows:

$$\sum_{|\alpha| < k} c_\alpha \partial^\alpha u = f(x).$$

If  $f(x) = 0$ , the PDE is **homogeneous**, otherwise it is **inhomogeneous**.

This can be separated into linear PDEs with constant coefficients,  $c_\alpha \in \mathbb{R}, \mathbb{C}$  and variable coefficients,  $c_\alpha = c_\alpha(x)$ . [In this class, we focus on constant coefficient PDEs, but many of the techniques can be extended to variable coefficient PDEs.]

**Definition 1.2** (Nonlinear PDE). We look at a function  $F = F(u, \partial u, \dots, \partial^k u)$ . The highest order terms are take the *leading role*.

- Semilinear PDE's:  $F$  is linear, with constant or variable coefficients in  $\partial^k u$ :

$$\sum_{|\alpha|=k} c_\alpha(x) \partial^\alpha u = N(\partial^{\leq k-1} u).$$

The LHS is called the principal part, and the RHS is the perturbative role.

- Quasilinear PDE's:

$$\sum_{|\alpha|=k} c_\alpha(\partial^{\leq k-1} u) \partial^\alpha u = N(\partial^{\leq k-1} u).$$

- Fully Nonlinear PDE's:  $F(\partial^{\leq k} u) = 0$ , with a nonlinear dependence on  $\partial^k u$ .

Some examples:

- Linear, homogeneous, variable coefficients, order 1:

$$\sum_{k=1}^u c_k(x) \partial_k(u) = 0.$$

- Define  $\Delta = \partial_1^2 + \cdots + \partial_n^2$ , the Laplacian operator. We have a linear, constant coefficients, inhomogeneous, order 2:

$$\Delta u = f.$$

- Semilinear, order 2:

$$\Delta u = u^3.$$

[Note that translation invariance makes homogeneous vs inhomogeneous not useful for classification in the case of nonlinear PDE's.]

- Harmonic Map Equation:

$$\Delta u = u |\nabla u|^2.$$

It is still semilinear, but with a stronger nonlinearity.

- Monge Ampere Equation:

$$\mathbb{R}^2, \partial_1^2 u \partial_2^2 u - (\partial_1 \partial_2 u)^2 = 0.$$

It is a fully nonlinear equation.

### §1.3 Initial Value Problems

We have various types of problems:

- (Stationary Problems) With  $u : \mathbb{R}^n \rightarrow \mathbb{R}$ ,

$$F(\partial^{\leq k} u) = 0,$$

might describe an equilibrium configuration of a physical system.

- (Evolution Equations) With  $u : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $u(t, x)$  describes the state at time  $t$ . We can think about the order in  $x$  or in  $t$ .

**Definition 1.3** (Initial Value Problem/Cauchy Problem). A PDE with initial conditions.

#### Example 1.4

Consider the heat equation:

$$\begin{aligned} \partial_t u &= \Delta_x u, \\ u(t=0, x) &= u_o(x). \end{aligned}$$

The equation is first order in  $t$ , but second order in  $x$ .

**Example 1.5**

In  $[\mathbb{R} \times \mathbb{R}]$ , the vibrating string:

$$\partial_t^2 u = \partial_x^2 u,$$

$$u(t = 0, x) = u_0(x),$$

$$\partial_t u(t = 0, x) = u_1(x).$$

Note that this equation is second order in time, and requires 2 pieces of initial data.

An easier problem: Compute the Taylor series of  $u$  at some point  $(0, x_0)$ . It requires  $\partial_t^\alpha \partial_x^\beta u(0, x_0)$ .

- This is obvious if we have no time derivative or exactly 1.
- Second order time derivatives come from the equation.
- Third order or higher time derivatives come from differentiating the equation:

$$\partial_t^3 u = \partial_x^2 \partial_t u.$$

**§1.4 Boundary Value Problems**

We begin with an example.

**Example 1.6**

Take  $\Delta u = f$  in  $\Omega \subset \mathbb{R}^3$ , which represents equilibrium for temperature in a solid. To solve, we need information about the boundary of  $\Omega$ . For example,

$$\Delta u = f \in \Omega,$$

$$u = g \in \partial\Omega.$$

**§1.5 Fluid Classification**

We take  $u : \mathbb{R}^n \rightarrow \mathbb{R}(\mathbb{C})$ , and

$$F(\partial^{\leq k} u) = 0.$$

This is considered to be a **scalar equation**.

We could also take a **system** of equations, where  $u : \mathbb{R}^n \rightarrow \mathbb{R}^m(\mathbb{C}^m)$ , where  $u = [u_i]$  a column of equations. These are often more difficult than scalar equation. We should have

$$F(\partial^{\leq k} u) = 0,$$

but  $F : \mathbb{R}^{(\cdot)} \rightarrow \mathbb{R}^m(\mathbb{C}^m)$ .

**Example 1.7**

A 2-system:

$$\Delta u = v,$$

$$\Delta v = -u.$$

We can often reduce the order of a scalar equation by turning it into a system:

**Example 1.8**

Consider the vibrating string,

$$\partial_t^2 u = \partial_x^2 u.$$

If we take  $v = \partial_t u$ , then it suffices to solve the system,

$$\partial_t u = v,$$

$$\partial_t v = \partial_x^2 u.$$

We can reduce it further by saying  $u_1 = \partial_x u, u_2 = \partial_t u$  for the system,

$$\partial_t u_1 = \partial_x u_2,$$

$$\partial_t u_2 = \partial_x u_1.$$

## §2 September 3rd, 2020

### §2.1 Picard-Lindeloff Theorem

Consider the example,  $x' = f(x)$ ,  $x(0) = x_0$ ,  $x : \mathbb{R} \rightarrow \mathbb{R}^n$ . We ask for existence, uniqueness, continuous dependence on initial data.

**Definition 2.1** (Locally Lipschitz). A **Lipschitz** continuous function  $f$  is one that satisfies,

$$|f(x) - f(y)| \leq c|x - y|.$$


A function is **Locally Lipschitz** if for each  $R$ , there exists  $c(R)$  such that

$$|f(x) - f(y)| \leq c(r)|x - y|, x, y \in \text{Ball}(0, R).$$

As examples,  $f(x) = x$  is Lipschitz,  $f(x) = x^2$  is not Lipschitz, but is locally Lipschitz.

**Definition 2.2** (Locally well-posed). For each  $x_0 \in \mathbb{R}^n$ , there exists  $T > 0$  (lifespan) and a unique solution  $u \in C^1[0, T; \mathbb{R}^n]$  with the property that  $u_0 = x_0$  and the solution has a Lipschitz dependence on the data:  $x_0, y_0$  initial data,  $T = T(x_0)$ . For  $T_1 < T$ , there exists  $\epsilon > 0$  such that if  $|y_0 - x_0| \leq \epsilon$  then  $T(y) > T_1$  and

$$\sup_{t \in [0, T_1]} |x(t) - y(t)| \leq \tilde{C}|x_0 - y_0|.$$



#### Theorem 1 (Picard-Lindelof)

Assume that  $f$  is locally Lipschitz continuous. Then the ODE is locally well-posed.

### §2.2 Contraction Principle

We will use the "Contraction principle" - recall the following definitions:

**Definition 2.3** (Fixed-point Problem). Let  $X$  be a Banach space, let  $D \subset X$  be a closed subset of  $X$ , and let  $F : D \rightarrow D$ . Question: Can we solve the equation  $F(u) = u$  where  $u \in D$ .

**Definition 2.4** (Contraction).

$$\|F(u) - F(v)\|_X \leq L\|u - v\|,$$


where  $L < 1$ .

If  $F$  is a contraction, then it has a unique fixed point. The existence proof follows an iterative construction: start with an arbitrary element  $u_0 \in D$  and define  $u_{n+1} = F(u_n)$ . We would show  $\{u_n\}$  is a Cauchy sequence, so it converges.

We now prove the theorem. We have  $x' = f(x)$ ,  $x(0) = x_0$ , so

$$x(t) = x_0 + \int_0^t f(x(s))ds, t \in [0, T].$$

We choose  $X = C[0, T; \mathbb{R}^n]$ ,  $F(x)(t) = x_0 + \int_0^t f(x(s))ds$ . Then  $x$  solves the ODE in  $(0, T)$  if  $F(x) = x$ .



We have to choose  $R, T$ . Then

$$D = \{x \in X : \|x - x_0\|_X \leq R\}.$$

Let  $R = |x_0|$ . Next, we choose  $T$  so that  $F : D \rightarrow D$  is Lipschitz. For  $F : D \rightarrow D$ , we estimate the size of  $F(x) - x_0$ .

$$\begin{aligned} |F(x)(t) - x_0| &= \left| \int_0^t f(x(s))ds \right| \\ &\leq \left| \int_0^t f(x_0(s))ds \right| + \left| \int_0^t f(x) - f(x_0)ds \right| \\ &\leq T|f(x_0)| + CT\|x - x_0\|_X \end{aligned}$$

Hence,

$$\|F(x) - x_0\| \leq T(|f(x_0)| + CR).$$

Thus, we choose  $T$  such that  $T(|f(x_0)| + CR) \leq R$ .

Now look at differences: For  $x, y \in D$ ,

$$\begin{aligned} |F(x)(t) - F(y)(t)| &\leq \int_0^t |f(x(s)) - f(y(s))|ds \\ &\leq TC \sup_{s \in [0, T]} |x(s) - y(s)| \end{aligned}$$

thus,


$$\|F(x) - F(y)\|_X \leq CT\|x - y\|_X,$$

so we can choose  $T$  so that  $CT\|x - y\|_X < 1$ .

By the contraction principle, there exists a unique solution  $x \in D$ .

To prove uniqueness of a solution, we have to show that any solution has to stay in  $D$ , up to time  $T$ .






Suppose a solution  $\tilde{x}$  leave the ball before time  $T$ . We repeat the above computation up to the exit time  $T_0$ . Then,  $T_0(|f(x_0) + CR|) < T$ , since  $T_0 < T$ . This is a contradiction since  $T_0$  is the exit time.

### §2.3 Bootstrap Argument

Consider a bootstrap argument: try to solve an equation and show that the solution  $x$  satisfies some bound  $\|x\|_T \leq R$ . The difficulty is that a priori, we do not know any bound on  $\|x\|_T$ . The solution: make a bootstrap assumption,  $\|x\|_T \leq 2R$  and show that  $\|x\|_T \leq R$  under this assumption.



So far, we know uniqueness in  $[0, T]$ , where  $T = T(x_0)$  given by the contraction argument. We now show global uniqueness: Suppose we have a solution  $x_0$  with maximal lifespan  $T_{max}(x_0)$ . Suppose  $y$  is another solution. We look at the maximal  $T$  so that  $x = y$  in  $[0, T)$ . We now think of  $T$  as the initial time. We  $x(T) = y(T)$  from continuity. Then, the solution is unique up to some time  $T + T_0$ , so  $x = y$  in  $[T, T + T_0]$ , contradicting the maximality of  $T$ . This is called a "continuity argument".

Next, we compare two solutions: We have  $x(0) = x_0, x : [0, T) \rightarrow \mathbb{R}^n$ . We choose  $T_1 < T$ . Then  $x : [0, T_1] \rightarrow \mathbb{R}^n$ . We compare  $x$  with a "nearby" solution  $y(0) = y_0$  close to  $x_0$ . We have  $\|x\|_{X_{T_1}} \leq R$  since we have continuity on a compact set. We claim the following: if  $|y_0 - x_0| < \epsilon$ , then  $x, y$  stay close. We make a bootstrap assumption  $\|y\|_{X_{T_1}} \leq 2R$ .

$$\frac{d}{dt}|x - y|^2 = 2(x - y)(f(x) - f(y)) \leq 2C|x - y|^2.$$

This is the *Gronwall Inequality*. It follows that

$$|x - y|^2(t) \leq e^{2ct}|x - y|^2(0) = e^{2ct}|x_0 - y_0|^2.$$

To close the bootstrap:

$$\|y\|_{X_{T_1}} \leq \|x\|_{X_{T_1}} + \|x - y\|_{X_{T_1}} \leq R + e^{cT_1}\|x_0 - y_0\| \leq \frac{3R}{2},$$

which is better than the bootstrap assumption.