

Mathematics for Informatics

Carlos Areces and Patrick Blackburn

`areces@loria.fr`

`blackbur@loria.fr`

<http://www.loria.fr/~areces>

<http://www.loria.fr/~blackbur>

INRIA Lorraine
Nancy, France

2007/2008

Summatory and Productory (from 0)

Theorem

Let \mathcal{C} be a PRC class. If $f : \mathbb{N}^{n+1} \rightarrow \mathbb{N}$ is in \mathcal{C} then the functions

$$g(y, x_1, \dots, x_n) = \sum_{t=0}^y f(t, x_1, \dots, x_n)$$

$$h(y, x_1, \dots, x_n) = \prod_{t=0}^y f(t, x_1, \dots, x_n)$$

are also in \mathcal{C} .

Summatory and Productory (from 0)

Theorem

Let \mathcal{C} be a PRC class. If $f : \mathbb{N}^{n+1} \rightarrow \mathbb{N}$ is in \mathcal{C} then the functions

$$g(y, x_1, \dots, x_n) = \sum_{t=0}^y f(t, x_1, \dots, x_n)$$

$$h(y, x_1, \dots, x_n) = \prod_{t=0}^y f(t, x_1, \dots, x_n)$$

are also in \mathcal{C} .

Proof.

$$\begin{aligned} g(0, x_1, \dots, x_n) &= f(0, x_1, \dots, x_n) \\ g(t+1, x_1, \dots, x_n) &= g(t, x_1, \dots, x_n) + f(t+1, x_1, \dots, x_n) \end{aligned}$$

Idem for h with \cdot instead of $+$. □

Observe that it is not important in which variable we do the recursion (we can define $g'(x, t)$ as in the last class and then $g(t, x) = g'(x, t)$)

Sumatorias y productorias (desde 1)

Theorem

Let \mathcal{C} be a PRC class. If $f : \mathbb{N}^{n+1} \rightarrow \mathbb{N}$ is in \mathcal{C} then also the following functions are in \mathcal{C}

$$g(y, x_1, \dots, x_n) = \sum_{t=1}^y f(t, x_1, \dots, x_n)$$

$$h(y, x_1, \dots, x_n) = \prod_{t=1}^y f(t, x_1, \dots, x_n)$$

(as usual, empty summatory = 0, empty productory = 1)

Sumatorias y productorias (desde 1)

Theorem

Let \mathcal{C} be a PRC class. If $f : \mathbb{N}^{n+1} \rightarrow \mathbb{N}$ is in \mathcal{C} then also the following functions are in \mathcal{C}

$$g(y, x_1, \dots, x_n) = \sum_{t=1}^y f(t, x_1, \dots, x_n)$$

$$h(y, x_1, \dots, x_n) = \prod_{t=1}^y f(t, x_1, \dots, x_n)$$

(as usual, empty summatory = 0, empty productory = 1)

Proof.

$$g(0, x_1, \dots, x_n) = 0$$

$$g(t+1, x_1, \dots, x_n) = g(t, x_1, \dots, x_n) + f(t+1, x_1, \dots, x_n)$$

Idem for h with \cdot instead of $+$ and 1 instead of 0 in the base case. □

Bounded quantification

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate

$(\forall t)_{\leq y} p(t, x_1, \dots, x_n)$ is true iff

▶ $p(0, x_1, \dots, x_n)$ is true **and**

⋮

▶ $p(y, x_1, \dots, x_n)$ is true

$(\exists t)_{\leq y} p(t, x_1, \dots, x_n)$ is true iff

▶ $p(0, x_1, \dots, x_n)$ is true **or**

⋮

▶ $p(y, x_1, \dots, x_n)$ is true

We can also define versions with $< y$ instead of $\leq y$.

$$(\exists t)_{< y} p(t, x_1, \dots, x_n) \quad \text{and} \quad (\forall t)_{< y} p(t, x_1, \dots, x_n)$$

Bounded quantification (with \leq)

Theorem

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} . The following predicates are also in \mathcal{C} :

$$(\forall t)_{\leq y} p(t, x_1, \dots, x_n)$$

$$(\exists t)_{\leq y} p(t, x_1, \dots, x_n)$$

Bounded quantification (with \leq)

Theorem

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} . The following predicates are also in \mathcal{C} :

$$(\forall t)_{\leq y} p(t, x_1, \dots, x_n)$$

$$(\exists t)_{\leq y} p(t, x_1, \dots, x_n)$$

Proof.

$$(\forall t)_{\leq y} p(t, x_1, \dots, x_n) \text{ iff } \prod_{t=0}^y p(t, x_1, \dots, x_n) = 1$$

Bounded quantification (with \leq)

Theorem

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} . The following predicates are also in \mathcal{C} :

$$(\forall t)_{\leq y} p(t, x_1, \dots, x_n)$$

$$(\exists t)_{\leq y} p(t, x_1, \dots, x_n)$$

Proof.

$$(\forall t)_{\leq y} p(t, x_1, \dots, x_n) \text{ iff } \prod_{t=0}^y p(t, x_1, \dots, x_n) = 1$$

$$(\exists t)_{\leq y} p(t, x_1, \dots, x_n) \text{ iff } \sum_{t=0}^y p(t, x_1, \dots, x_n) \neq 0$$

Bounded quantification (with \leq)

Theorem

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} . The following predicates are also in \mathcal{C} :

$$(\forall t)_{\leq y} p(t, x_1, \dots, x_n)$$

$$(\exists t)_{\leq y} p(t, x_1, \dots, x_n)$$

Proof.

$$(\forall t)_{\leq y} p(t, x_1, \dots, x_n) \text{ iff } \prod_{t=0}^y p(t, x_1, \dots, x_n) = 1$$

$$(\exists t)_{\leq y} p(t, x_1, \dots, x_n) \text{ iff } \sum_{t=0}^y p(t, x_1, \dots, x_n) \neq 0$$

- ▶ summatory and productory are in \mathcal{C}
- ▶ comparison by $=$ is in \mathcal{C}



Bounded quantification (with $<$)

Theorem

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} . Then the following predicates are also in \mathcal{C} :

$$(\forall t)_{<y} p(t, x_1, \dots, x_n)$$

$$(\exists t)_{<y} p(t, x_1, \dots, x_n)$$

Proof.

$$(\forall t)_{<y} p(t, x_1, \dots, x_n) \text{ iff } (\forall t)_{\leq y} (t = y \vee p(t, x_1, \dots, x_n))$$

$$(\exists t)_{<y} p(t, x_1, \dots, x_n) \text{ iff } (\exists t)_{\leq y} (t \neq y \wedge p(t, x_1, \dots, x_n))$$



More examples of primitive recursive functions

- ▶ $y|x$ iff y divides x . It can be defined as

$$(\exists t)_{\leq x} y \cdot t = x$$

Note that with this definition $0|0$.

- ▶ $prime(x)$ iff x is prime.

Minimization

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} .

$$g(y, x_1, \dots, x_n) = \sum_{u=0}^y \prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n))$$

What does g do?

Minimization

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} .

$$g(y, x_1, \dots, x_n) = \sum_{u=0}^y \prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n))$$

What does g do?

- ▶ suppose that there is $t \leq y$ such that $p(t, x_1, \dots, x_n)$ is true

Minimization

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} .

$$g(y, x_1, \dots, x_n) = \sum_{u=0}^y \prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n))$$

What does g do?

- ▶ suppose that there is $t \leq y$ such that $p(t, x_1, \dots, x_n)$ is true
 - ▶ let t_0 be the maximal such t

Minimization

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} .

$$g(y, x_1, \dots, x_n) = \sum_{u=0}^y \prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n))$$

What does g do?

- ▶ suppose that there is $t \leq y$ such that $p(t, x_1, \dots, x_n)$ is true
 - ▶ let t_0 be the maximal such t
 - ▶ $p(t, x_1, \dots, x_n) = 0$ for each $t < t_0$

Minimization

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} .

$$g(y, x_1, \dots, x_n) = \sum_{u=0}^y \prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n))$$

What does g do?

- ▶ suppose that there is $t \leq y$ such that $p(t, x_1, \dots, x_n)$ is true
 - ▶ let t_0 be the maximal such t
 - ▶ $p(t, x_1, \dots, x_n) = 0$ for each $t < t_0$
 - ▶ $p(t_0, x_1, \dots, x_n) = 1$

Minimization

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} .

$$g(y, x_1, \dots, x_n) = \sum_{u=0}^y \prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n))$$

What does g do?

- ▶ suppose that there is $t \leq y$ such that $p(t, x_1, \dots, x_n)$ is true
 - ▶ let t_0 be the maximal such t
 - ▶ $p(t, x_1, \dots, x_n) = 0$ for each $t < t_0$
 - ▶ $p(t_0, x_1, \dots, x_n) = 1$
 - ▶ $\prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n)) = \begin{cases} 1 & \text{if } u < t_0 \\ 0 & \text{otherwise} \end{cases}$

Minimization

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} .

$$g(y, x_1, \dots, x_n) = \sum_{u=0}^y \prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n))$$

What does g do?

- ▶ suppose that there is $t \leq y$ such that $p(t, x_1, \dots, x_n)$ is true
 - ▶ let t_0 be the maximal such t
 - ▶ $p(t, x_1, \dots, x_n) = 0$ for each $t < t_0$
 - ▶ $p(t_0, x_1, \dots, x_n) = 1$
 - ▶ $\prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n)) = \begin{cases} 1 & \text{if } u < t_0 \\ 0 & \text{otherwise} \end{cases}$
 - ▶ $g(y, x_1, \dots, x_n) = \underbrace{1 + 1 + \dots + 1}_{t_0 \text{ veces}} = t_0$
- ▶ then $g(x_1, \dots, x_n)$ is the maximal t such that $p(t, x_1, \dots, x_n)$ is true

Minimization

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} .

$$g(y, x_1, \dots, x_n) = \sum_{u=0}^y \prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n))$$

What does g do?

- ▶ suppose that there is $t \leq y$ such that $p(t, x_1, \dots, x_n)$ is true
 - ▶ let t_0 be the maximal such t
 - ▶ $p(t, x_1, \dots, x_n) = 0$ for each $t < t_0$
 - ▶ $p(t_0, x_1, \dots, x_n) = 1$
 - ▶ $\prod_{t=0}^u \alpha(p(t, x_1, \dots, x_n)) = \begin{cases} 1 & \text{if } u < t_0 \\ 0 & \text{otherwise} \end{cases}$
 - ▶ $g(y, x_1, \dots, x_n) = \underbrace{1 + 1 + \dots + 1}_{t_0 \text{ veces}} = t_0$
 - ▶ then $g(x_1, \dots, x_n)$ is the maximal t such that $p(t, x_1, \dots, x_n)$ is true
- ▶ if there is no such t , then $g(y, x_1, \dots, x_n) = y + 1$

Minimization

We note

$$\min_{t \leq y} p(t, x_1, \dots, x_n) = \begin{cases} \text{smallest } t \leq y \text{ such that} \\ p(t, x_1, \dots, x_n) \text{ is true} & \text{if there is such a } t \\ 0 & \text{otherwise} \end{cases}$$

Theorem

Let $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ be a predicate in a PRC class \mathcal{C} . Then the function

$$\min_{t \leq y} p(t, x_1, \dots, x_n)$$

is also in \mathcal{C} .

More examples of primitive recursive functions

- ▶ $x \text{ div } y$ is the integer division of x by y

$$\min_{t \leq x} ((t + 1) \cdot y > x)$$

Note that with this definition $0 \text{ div } 0$ is false.

More examples of primitive recursive functions

- ▶ $x \text{ div } y$ is the integer division of x by y

$$\min_{t \leq x} ((t + 1) \cdot y > x)$$

Note that with this definition $0 \text{ div } 0$ is false.

- ▶ $x \text{ mod } y$ is the remainder of dividing x by y

More examples of primitive recursive functions

- ▶ $x \text{ div } y$ is the integer division of x by y

$$\min_{t \leq x} ((t + 1) \cdot y > x)$$

Note that with this definition $0 \text{ div } 0$ is false.

- ▶ $x \text{ mod } y$ is the remainder of dividing x by y
- ▶ p_n is the n -th prime ($n > 0$). It is defined as $p_0 = 0, p_1 = 2, p_2 = 3, p_3 = 5, \dots$

More examples of primitive recursive functions

- ▶ $x \text{ div } y$ is the integer division of x by y

$$\min_{t \leq x} ((t + 1) \cdot y > x)$$

Note that with this definition $0 \text{ div } 0$ is false.

- ▶ $x \text{ mod } y$ is the remainder of dividing x by y
- ▶ p_n is the n -th prime ($n > 0$). It is defined as $p_0 = 0, p_1 = 2, p_2 = 3, p_3 = 5, \dots$

$$\begin{aligned} p_0 &= 0 \\ p_{n+1} &= \min_t (\text{primo}(t) \wedge t > p_n) \end{aligned}$$

More examples of primitive recursive functions

- ▶ $x \text{ div } y$ is the integer division of x by y

$$\min_{t \leq x} ((t + 1) \cdot y > x)$$

Note that with this definition $0 \text{ div } 0$ is false.

- ▶ $x \text{ mod } y$ is the remainder of dividing x by y
- ▶ p_n is the n -th prime ($n > 0$). It is defined as $p_0 = 0, p_1 = 2, p_2 = 3, p_3 = 5, \dots$

$$\begin{aligned} p_0 &= 0 \\ p_{n+1} &= \min_{t \leq K(n)} (\text{primo}(t) \wedge t > p_n) \end{aligned}$$

We need a good bound $K(n)$, i.e.

- ▶ sufficiently big and
- ▶ primitive recursive

More examples of primitive recursive functions

- ▶ $x \text{ div } y$ is the integer division of x by y

$$\min_{t \leq x} ((t + 1) \cdot y > x)$$

Note that with this definition $0 \text{ div } 0$ is false.

- ▶ $x \text{ mod } y$ is the remainder of dividing x by y
- ▶ p_n is the n -th prime ($n > 0$). It is defined as $p_0 = 0, p_1 = 2, p_2 = 3, p_3 = 5, \dots$

$$\begin{aligned} p_0 &= 0 \\ p_{n+1} &= \min_{t \leq K(n)} (\text{primo}(t) \wedge t > p_n) \end{aligned}$$

We need a good bound $K(n)$, i.e.

- ▶ sufficiently big and
- ▶ primitive recursive

$K(n) = p_n! + 1$ is enough (note that $p_{n+1} \leq p_n! + 1$).

Bounded minimization

Recall the definition of **bounded minimization**:

$$\min_{t \leq y} p(t, x_1, \dots, x_n) = \begin{cases} \text{smallest } t \leq y \text{ such that} \\ p(t, x_1, \dots, x_n) \text{ is true} & \text{if there is such a } t \\ 0 & \text{otherwise} \end{cases}$$

Bounded minimization

Recall the definition of **bounded minimization**:

$$\min_{t \leq y} p(t, x_1, \dots, x_n) = \begin{cases} \text{smallest } t \leq y \text{ such that} \\ p(t, x_1, \dots, x_n) \text{ is true} & \text{if there is such a } t \\ 0 & \text{otherwise} \end{cases}$$

What happens if there is no bound? We define **unbounded minimization** as

$$\min_t p(t, x_1, \dots, x_n) = \begin{cases} \text{smallest } t \text{ such that} \\ p(t, x_1, \dots, x_n) \text{ is true} & \text{if there is such a } t \\ \uparrow & \text{otherwise} \end{cases}$$

Unbounded minimisation

Theorem

If $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ is a computable predicate then

$$\min_t p(t, x_1, \dots, x_n)$$

is a partially computable predicate.

Unbounded minimisation

Theorem

If $p : \mathbb{N}^{n+1} \rightarrow \{0, 1\}$ is a computable predicate then

$$\min_t p(t, x_1, \dots, x_n)$$

is a partially computable predicate.

Proof.

The following program computes $\min_t p(t, x_1, \dots, x_n)$:

```
[A]   IF  $p(X_1, \dots, X_n, Y) = 1$  GOTO  $E$   
       $Y \leftarrow Y + 1$   
      GOTO  $A$ 
```



Bounded and unbounded minimization

What we know:

- ▶ a `while` can simulate an unbounded minimization
- ▶ a `for` can simulate a bounded minimization

Bounded and unbounded minimization

What we know:

- ▶ a `while` can simulate an unbounded minimization
- ▶ a `for` can simulate a bounded minimization

Questions:

- ▶ Can we write every program of \mathcal{S} as a `for`-program?
- ▶ can we find, for any unbounded minimization, a proper bound to turn it into a bounded minimization?
- ▶ computable = primitive recursive?