# Logics for Computation

Lecture #5: About Trees, and How to Cut Them

Carlos Areces and Patrick Blackburn {carlos.areces,patrick.blackburn}@loria.fr

INRIA Nancy Grand Est Nancy, France

ESSLLI 2008 - Hamburg - Germany

▶ We have introduced the  $\langle R \rangle$  operator to talk about complex relational structures.

Nothing fancy (yet), just a simple extension of PL.

- ▶ We have introduced the  $\langle R \rangle$  operator to talk about complex relational structures.
  - Nothing fancy (yet), just a simple extension of PL.
- ▶ We have used it to describe some properties over models.

E.g., the following formulas are valid: 
$$\langle R \rangle (p \wedge q) \rightarrow \langle R \rangle p \langle R \rangle \top \wedge \neg \langle R \rangle \neg p \rightarrow \langle R \rangle \neg p$$

- ▶ We have introduced the  $\langle R \rangle$  operator to talk about complex relational structures.
  - Nothing fancy (yet), just a simple extension of PL.
- ▶ We have used it to describe some properties over models.

▶ We have discussed when two models are the same.

- ▶ We have introduced the  $\langle R \rangle$  operator to talk about complex relational structures.
  - Nothing fancy (yet), just a simple extension of PL.
- ▶ We have used it to describe some properties over models.

E.g., the following formulas are valid: 
$$\langle R \rangle (p \wedge q) \rightarrow \langle R \rangle p \ \langle R \rangle \top \wedge \neg \langle R \rangle \neg p \rightarrow \langle R \rangle \neg p$$

- ▶ We have discussed when two models are the same.
- We have seen an algorithm to check whether a formula is true in a given model.

▶ We will define a tableax algorithm for satisfiability of formulas containing  $\langle R \rangle$ .

- ▶ We will define a tableax algorithm for satisfiability of formulas containing  $\langle R \rangle$ .
  - ▶ We already know how to check, given a model, if the formula holds in the model (model checking).
  - Today, we will see how do we check whether a formula has a model.

- ▶ We will define a tableax algorithm for satisfiability of formulas containing  $\langle R \rangle$ .
  - We already know how to check, given a model, if the formula holds in the model (model checking).
  - Today, we will see how do we check whether a formula has a model.
- We will also go back to the question When are two models the same?

- ▶ We will define a tableax algorithm for satisfiability of formulas containing  $\langle R \rangle$ .
  - ▶ We already know how to check, given a model, if the formula holds in the model (model checking).
  - Today, we will see how do we check whether a formula has a model.
- We will also go back to the question When are two models the same?
  - and talk about trees . . .

- ▶ We will define a tableax algorithm for satisfiability of formulas containing  $\langle R \rangle$ .
  - We already know how to check, given a model, if the formula holds in the model (model checking).
  - Today, we will see how do we check whether a formula has a model.
- We will also go back to the question When are two models the same?
  - ▶ and talk about trees . . .
  - ...and how to cut them.

► The proof that the satisfiability problem for PL is decidable is very simple:

- ► The proof that the satisfiability problem for PL is decidable is very simple:
  - ▶ Suppose that you are given a formula  $\varphi$  and you are looking for a model of  $\varphi$ .

- ► The proof that the satisfiability problem for PL is decidable is very simple:
  - ▶ Suppose that you are given a formula  $\varphi$  and you are looking for a model of  $\varphi$ .
  - ightharpoonup First note that propositional symbols that do not appear in  $\varphi$  are irrelevant.

- ► The proof that the satisfiability problem for PL is decidable is very simple:
  - ▶ Suppose that you are given a formula  $\varphi$  and you are looking for a model of  $\varphi$ .
  - First note that propositional symbols that do not appear in  $\varphi$  are irrelevant.
  - We know that our models has only one point.

- ► The proof that the satisfiability problem for PL is decidable is very simple:
  - ▶ Suppose that you are given a formula  $\varphi$  and you are looking for a model of  $\varphi$ .
  - First note that propositional symbols that do not appear in  $\varphi$  are irrelevant.
  - We know that our models has only one point.
  - ▶ Hence, we only need to list all possible ways of labelling that single node with propositional symbols in  $\varphi$ .

- ► The proof that the satisfiability problem for PL is decidable is very simple:
  - ▶ Suppose that you are given a formula  $\varphi$  and you are looking for a model of  $\varphi$ .
  - First note that propositional symbols that do not appear in  $\varphi$  are irrelevant.
  - We know that our models has only one point.
  - ▶ Hence, we only need to list all possible ways of labelling that single node with propositional symbols in  $\varphi$ .
- ▶ What about the  $\langle R \rangle$  language?

We want to devise a tableau method for the language we introduced to talk about complex relational structures.

- ▶ We want to devise a tableau method for the language we introduced to talk about complex relational structures.
- Let's review the tableaux method that we introduced for propositional logic:

$$\frac{(\varphi \wedge \psi)}{\varphi} (\wedge)$$

$$\frac{\neg(\varphi \wedge \psi)}{\neg \varphi} (\neg \wedge)$$

$$\frac{\neg \neg \varphi}{\varphi} (\neg \neg)$$

- ► We want to devise a tableau method for the language we introduced to talk about complex relational structures.
- Let's review the tableaux method that we introduced for propositional logic:

$$\frac{(\varphi \wedge \psi)}{\varphi} (\wedge)$$

$$\frac{\neg (\varphi \wedge \psi)}{\neg \varphi} (\neg \wedge)$$

$$\frac{\neg \neg \varphi}{\varphi} (\neg \neg)$$

Pretty neat: 3 rules for an NP-complete problem!

- ► We want to devise a tableau method for the language we introduced to talk about complex relational structures.
- Let's review the tableaux method that we introduced for propositional logic:

$$\frac{(\varphi \wedge \psi)}{\varphi} \ (\wedge)$$

$$\frac{\neg(\varphi \wedge \psi)}{\neg \varphi} \ (\neg \wedge)$$

$$\frac{\neg\neg\varphi}{\varphi}$$
 (¬¬)

- Pretty neat: 3 rules for an NP-complete problem!
- But now we want to deal with more than a single point.

- ▶ We want to devise a tableau method for the language we introduced to talk about complex relational structures.
- Let's review the tableaux method that we introduced for propositional logic:

$$\frac{s:(\varphi \wedge \psi)}{\underset{s:\psi}{s:\varphi}} \ (\wedge)$$

 $\frac{s:\neg\neg\varphi}{s:\omega}$   $(\neg\neg)$ 

$$\frac{s:\neg(\varphi \wedge \psi)}{s:\neg\varphi} (\neg \wedge)$$

- Pretty neat: 3 rules for an NP-complete problem!
- But now we want to deal with more than a single point.
- ► The solution is: labels! ¬

- ► We want to devise a tableau method for the language we introduced to talk about complex relational structures.
- Let's review the tableaux method that we introduced for propositional logic:

$$\frac{s:(\varphi \wedge \psi)}{\underset{s:\psi}{s:\varphi}} \ (\wedge)$$

$$\frac{s:\neg(\varphi \wedge \psi)}{s:\neg\varphi} \ (\neg \wedge)$$

$$\frac{s:\neg\neg\varphi}{s:\varphi}$$
 (¬¬)

- Pretty neat: 3 rules for an NP-complete problem!
- But now we want to deal with more than a single point.
- ► The solution is: labels! ¬
- ► They will help us keep track of what is going on in each point in our model.

▶ We have dealt in the previous slide with multiple points. What about lines?

- ▶ We have dealt in the previous slide with multiple points. What about lines?
- ▶ Remember that the operator we introduced to talk about lines in our language was  $\langle R \rangle \varphi$  and we said that

$$\mathcal{M}, w \models \langle R \rangle \varphi$$
 iff there is  $w'$  s.t.  $wRw'$  and  $\mathcal{M}, w' \models \varphi$ .

- ▶ We have dealt in the previous slide with multiple points. What about lines?
- ▶ Remember that the operator we introduced to talk about lines in our language was  $\langle R \rangle \varphi$  and we said that

$$\mathcal{M}, w \models \langle R \rangle \varphi$$
 iff there is  $w'$  s.t.  $wRw'$  and  $\mathcal{M}, w' \models \varphi$ .

Start with the labelled formula 
$$s:\langle R \rangle \varphi$$
.  $\longrightarrow \frac{s:\langle R \rangle \varphi}{sRt \atop t:\varphi}$  ( $\langle R \rangle$ )

for t a new label

- ► We have dealt in the previous slide with multiple points. What about lines?
- ▶ Remember that the operator we introduced to talk about lines in our language was  $\langle R \rangle \varphi$  and we said that

$$\mathcal{M}, w \models \langle R \rangle \varphi$$
 iff there is  $w'$  s.t.  $wRw'$  and  $\mathcal{M}, w' \models \varphi$ .

Start with the labelled formula  $s:\langle R\rangle\varphi$ .

If this formula is satisfiable, it is because there is an R-successor t where  $\varphi$  holds.  $s:\langle R\rangle\varphi$  sRt  $t:\varphi$ for t a new label

- ▶ We have dealt in the previous slide with multiple points. What about lines?
- ▶ Remember that the operator we introduced to talk about lines in our language was  $\langle R \rangle \varphi$  and we said that

$$\mathcal{M}, w \models \langle R \rangle \varphi$$
 iff there is  $w'$  s.t.  $wRw'$  and  $\mathcal{M}, w' \models \varphi$ .

Start with the labelled formula  $s:\langle R \rangle \varphi$ . If this formula is satisfiable, it is because there is an R-sucessor t where  $\varphi$  holds.

$$\frac{s:\langle R\rangle\varphi}{\underset{t:\varphi}{sRt}}\ (\langle R\rangle)$$

for t a new label

Start with the labelled formula 
$$s: \neg \langle R \rangle \varphi. \longrightarrow s: \neg \langle R \rangle \varphi$$

$$\frac{sRt}{t: \neg \varphi} (\neg \langle R \rangle)$$

- ▶ We have dealt in the previous slide with multiple points. What about lines?
- ▶ Remember that the operator we introduced to talk about lines in our language was  $\langle R \rangle \varphi$  and we said that

$$\mathcal{M}, w \models \langle R \rangle \varphi$$
 iff there is  $w'$  s.t.  $wRw'$  and  $\mathcal{M}, w' \models \varphi$ .

▶ Start with the labelled formula  $s:\langle R \rangle \varphi$ . If this formula is satisfiable, it is because there is an R-sucessor t where  $\varphi$  holds.

$$\frac{s:\langle R \rangle \varphi}{\underset{t:\varphi}{sRt}} (\langle R \rangle)$$
 for t a new label

Start with the labelled formula  $s: \neg \langle R \rangle \varphi$ . If there is an R-successor t, then  $\varphi$  should not hold at t.

$$\stackrel{S:\neg\langle R\rangle\varphi}{\longrightarrow} sRt \\
\xrightarrow{t:\neg\varphi} (\neg\langle R\rangle)$$

The Complete Cast, plus an Example

# The Complete Cast, plus an Example

$$\frac{s:(\varphi \wedge \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt}$$

$$\frac{s:\neg(\varphi \wedge \psi)}{s:\neg(\varphi \wedge \psi)} \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \wedge \psi)}{s:\neg(\varphi \wedge \psi)} \qquad \frac{s:\neg(\varphi \wedge \psi)}{s:\neg(\varphi \wedge \psi)}$$

$$\frac{s:\neg(\varphi \wedge \psi)}{s:\neg(\varphi \wedge \psi)} \qquad \frac{s:\neg(\varphi \wedge \psi)}{t:\neg(\varphi \wedge \psi)}$$

# The Complete Cast, plus an Example

$$\frac{s:(\varphi \wedge \psi)}{s:\varphi} \frac{s:\langle R \rangle \varphi}{sRt}$$

$$\frac{s:\neg(\varphi \wedge \psi)}{s:\neg(\varphi \wedge \psi)}$$

$$s:(\neg\langle R\rangle p \wedge (\langle R\rangle q \wedge \langle R\rangle p))$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg(\varphi \land \psi)}{s:\neg(\varphi \land \psi)} \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg(\varphi \land \psi)} \qquad \frac{s:\neg(\varphi \land \psi)}{s:\neg(\varphi \land \psi)} \qquad \frac{s:\neg(\varphi \land \psi)}{s:\neg(\varphi \land \psi)}$$

$$s: (\neg \langle R \rangle p \bigcirc (\langle R \rangle q \land \langle R \rangle p))$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg(\varphi \land \psi)}{s:\neg(\varphi \land \psi)} \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg\varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg\varphi}$$

$$s: (\neg \langle R \rangle p \wedge (\langle R \rangle q \wedge \langle R \rangle p)) \\ s: \neg \langle R \rangle p \\ s: (\langle R \rangle q \wedge \langle R \rangle p)$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg(\varphi \land \psi)}{s:\neg(\varphi \land \psi)} \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg\varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg\varphi}$$

$$s: (\neg \langle R \rangle p \wedge (\langle R \rangle q \wedge \langle R \rangle p)) \\ s: \neg \langle R \rangle p \\ s: (\langle R \rangle q \bigcirc \langle R \rangle p)$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg(\varphi \land \psi)}{s:\neg(\varphi \land \psi)} \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg\varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg\varphi}$$

$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg(\varphi \land \psi)}{s:\neg(\varphi \land \psi)} \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg\varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg\varphi}$$

$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: (R)q$$

$$s: \langle R \rangle p$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ t:\varphi \qquad \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg\psi}{s:\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg\neg\varphi}{s:\varphi} \qquad \frac{s:\neg\varphi}{t:\neg\varphi}$$

$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

$$sRt$$

$$t: q$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ \frac{s:(\varphi \land \psi)}{s:\neg \varphi} \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg \varphi}{s:\neg \varphi} \qquad \frac{s:\neg \varphi}{sRt} \\ \frac{s:\neg \varphi}{t:\neg \varphi}$$

$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: (R) p$$

$$sRt$$

$$t: q$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ t:\varphi \qquad \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg\psi}{s:\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg\neg\varphi}{s:\varphi} \qquad \frac{s:\neg\varphi}{t:\neg\varphi}$$

$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

$$sRt$$

$$t: q$$

$$sRu$$

$$u: p$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ t:\varphi \qquad \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg \langle R \rangle \varphi}{sRt} \\ \frac{s:\neg \neg \varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg \varphi}$$

$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

$$sRt$$

$$t: q$$

$$sRu$$

$$u: p$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ t:\varphi \qquad \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg \langle R \rangle \varphi}{sRt} \\ \frac{s:\neg \neg \varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg \varphi}$$

$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

$$sRt$$

$$t: q$$

$$sRu$$

$$u: p$$

$$t: \neg p$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ t:\varphi \qquad \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg \langle R \rangle \varphi}{sRt} \\ \frac{s:\neg \neg \varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg \varphi}$$

$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

$$sRt$$

$$t: q$$

$$sRu$$

$$u: p$$

$$t: \neg p$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ t:\varphi \qquad \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg\neg\varphi}{s:\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{t:\neg\varphi}$$

$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

$$sRt$$

$$t: q$$

$$sRu$$

$$u: p$$

$$t: \neg p$$

$$u: \neg p$$

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ t:\varphi \qquad \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg \langle R \rangle \varphi}{sRt} \\ \frac{s:\neg \neg \varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg \varphi}$$

```
s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))
                    s: \neg \langle R \rangle p
            s:(\langle R \rangle q \wedge \langle R \rangle p)
                      s:\langle R\rangle q
                       s:\langle R\rangle p
                          sRt
                           t:q
                          sRu
                           u:p
                          t:\neg p
                          u:¬p
               contradiction!!!
```

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg(\varphi \land \psi)}{t:\varphi} \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg\varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg\varphi}$$

```
s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))
s: \neg \langle R \rangle p
s: (\langle R \rangle q \land \langle R \rangle p)
s: \langle R \rangle q
s: \langle R \rangle p
sRt
t: q
sRu
u: p
t: \neg p
u: \neg p
```

contradiction!!!

Which are the similarities/differences with tableaux for PL?

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ t:\varphi \qquad \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg\neg\varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg\varphi}$$

```
s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))
s: \neg \langle R \rangle p
s: (\langle R \rangle q \land \langle R \rangle p)
s: \langle R \rangle q
s: \langle R \rangle p
sRt
t: q
sRu
u: p
t: \neg p
u: \neg p
```

contradiction!!!

- Which are the similarities/differences with tableaux for PL?
- ▶ How do we know that we got it right?

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ t:\varphi \qquad \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad s:\neg\psi \qquad s:\neg\langle R \rangle \varphi \\ \frac{s:\neg\varphi}{s:\varphi} \qquad \frac{sRt}{t:\neg\varphi}$$

```
s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))
s: \neg \langle R \rangle p
s: (\langle R \rangle q \land \langle R \rangle p)
s: \langle R \rangle q
s: \langle R \rangle p
sRt
t: q
sRu
u: p
t: \neg p
u: \neg p
```


contradiction!!!

- ▶ Which are the similarities/differences with tableaux for PL?
- ▶ How do we know that we got it right?
- ▶ What can we learn from the calculus?


► Which similarities / differences with tableaux for PL?

$\frac{s:(\varphi \wedge \psi)}{s:\varphi}$ $s:\psi$	$\frac{s:\langle R\rangle\varphi}{sRt}$ $t:\varphi$
$s: \neg (\varphi \wedge \psi)$	for $t$ a new label
$\overline{s}:\neg\varphi$ $\overline{s}:\neg\psi$	$s:\neg\langle R\rangle\varphi$
$\frac{s:\neg\neg\varphi}{s:\varphi}$	$\frac{s\kappa t}{t:\neg\varphi}$


- ► Which similarities / differences with tableaux for PL?
  - Does the calculus terminate?




- ► Which similarities / differences with tableaux for PL?
  - ► Does the calculus terminate?
  - What are labels? What are they doing? Can we use them?




- ► Which similarities / differences with tableaux for PL?
  - Does the calculus terminate?
  - What are labels? What are they doing? Can we use them?
  - Is this an algorithm?




- ► Which similarities / differences with tableaux for PL?
  - ► Does the calculus terminate?
  - What are labels? What are they doing? Can we use them?
  - ► Is this an algorithm?
  - Is it a good algorithm?



- ► Which similarities / differences with tableaux for PL?
  - ► Does the calculus terminate?
  - What are labels? What are they doing? Can we use them?
  - ► Is this an algorithm?
  - ► Is it a good algorithm?
- ▶ Did we get it right?



- ► Which similarities / differences with tableaux for PL?
  - ► Does the calculus terminate?
  - What are labels? What are they doing? Can we use them?
  - ▶ Is this an algorithm?
  - Is it a good algorithm?
- ▶ Did we get it right?
  - Did we get it right in the PL case, to start with?!
    Consider the rule:



$$\frac{s:\neg(\varphi \land \psi)}{s:\varphi}$$
$$s:\neg\varphi$$
$$s:\neg\psi$$

- ► Which similarities / differences with tableaux for PL?
  - ▶ Does the calculus terminate?
  - What are labels? What are they doing? Can we use them?
  - ▶ Is this an algorithm?
  - Is it a good algorithm?
- ▶ Did we get it right?
  - Did we get it right in the PL case, to start with?!
    Consider the rule:
- What can we learn from the calculus?

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \frac{s:\langle R \rangle \varphi}{sRt} \frac{sRt}{t:\varphi}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \frac{s:\neg(\varphi \land \psi)}{s:\neg\psi} \frac{s:\neg\langle R \rangle \varphi}{sRt} \frac{s:\neg(\varphi \land \psi)}{t:\neg\varphi}$$

 $s:\neg\varphi$ 

**S**:φ

s:¬\//

 $s:\neg(\varphi \wedge \psi)$ 

- ► Which similarities / differences with tableaux for PL?
  - ► Does the calculus terminate?
  - What are labels? What are they doing? Can we use them?
  - Is this an algorithm?
  - ► Is it a good algorithm?
- ▶ Did we get it right?
  - Did we get it right in the PL case, to start with?! Consider the rule:
- ► What can we learn from the calculus?
  - Something about models!

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \qquad \frac{s:\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg\varphi}{s:\neg\varphi} \qquad \frac{s:\neg\psi}{sRt} \\ \frac{s:\neg\varphi}{t:\neg\varphi}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\varphi}$$
$$s:\neg\varphi$$
$$s:\neg\psi$$

► Let us see the tableux proof we did before again, for the formula

$$\varphi = \neg \langle R \rangle p \wedge (\langle R \rangle q \wedge \langle R \rangle p)$$

$$s:(\neg\langle R\rangle p \wedge (\langle R\rangle q \wedge \langle R\rangle p))$$

► Let us see the tableux proof we did before again, for the formula


$$\varphi = \neg \langle R \rangle p \wedge (\langle R \rangle q \wedge \langle R \rangle p)$$

$$s: (\neg \langle R \rangle p \wedge (\langle R \rangle q \wedge \langle R \rangle p))$$



Let us see the tableux proof we did before again, for the formula

$$\varphi = \neg \langle R \rangle p \wedge (\langle R \rangle q \wedge \langle R \rangle p)$$



$$s: (\neg \langle R \rangle p \wedge (\langle R \rangle q \wedge \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \wedge \langle R \rangle p)$$

Let us see the tableux proof we did before again, for the formula

$$\varphi = \neg \langle R \rangle p \wedge (\langle R \rangle q \wedge \langle R \rangle p)$$

$$s \left( \begin{array}{c} \neg \langle R \rangle p \\ \langle R \rangle q \\ \langle R \rangle p \end{array} \right)$$

$$s: (\neg \langle R \rangle p \wedge (\langle R \rangle q \wedge \langle R \rangle p))$$


$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \wedge \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

Let us see the tableux proof we did before again, for the formula



$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$


$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

$$sRt$$

$$t: q$$

▶ Let us see the tableux proof we did before again, for the formula



$$s: (\neg \langle R \rangle p \wedge (\langle R \rangle q \wedge \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \wedge \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$


$$sRt$$

$$t: q$$

$$sRu$$

$$u: p$$

▶ Let us see the tableux proof we did before again, for the formula



$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

$$sRt$$


$$t: q$$

$$sRu$$

$$u: p$$

$$t: \neg p$$

▶ Let us see the tableux proof we did before again, for the formula



$$s: (\neg \langle R \rangle p \land (\langle R \rangle q \land \langle R \rangle p))$$

$$s: \neg \langle R \rangle p$$

$$s: (\langle R \rangle q \land \langle R \rangle p)$$

$$s: \langle R \rangle q$$

$$s: \langle R \rangle p$$

$$sRt$$

$$t: q$$

$$sRu$$

$$u: p$$

$$t: \neg p$$

$$u: \neg p$$

# Tree and Finite Model Properties


 Using the rules of the tableaux calculus we only explore finite, tree models.

$$\frac{s:(\varphi \land \psi)}{s:\varphi} \atop s:\psi \qquad \frac{s:\langle R \rangle \varphi}{sRt} \atop t:\varphi}$$

$$\frac{s:\neg(\varphi \land \psi)}{s:\neg\varphi} \qquad \text{for } t \text{ a new label}$$

$$\frac{s:\neg\varphi}{s:\neg\varphi} \qquad \frac{s:\neg\langle R \rangle \varphi}{sRt} \atop t:\neg\varphi}$$

- Using the rules of the tableaux calculus we only explore finite, tree models.
- ► Let's assume that the calculus is correct (you will have to believe me).



- Using the rules of the tableaux calculus we only explore finite, tree models.
- ► Let's assume that the calculus is correct (you will have to believe me).
- ▶ Then the  $\langle R \rangle$ -language
  - cannot say infinite,
  - cannot say non-tree.

$\frac{s:(\varphi \wedge \psi)}{s:\varphi}$ $s:\psi$	$\frac{s:\langle R\rangle\varphi}{sRt}$ $t:\varphi$
$s:\neg(\varphi \wedge \psi)$	for $\overset{'}{t}$ a new label
$\begin{array}{ccc} s: \neg \varphi & s: \neg \psi \\ \hline \frac{s: \neg \neg \varphi}{s: \varphi} & \end{array}$	$\frac{s:\neg\langle R\rangle\varphi}{sRt} = \frac{t:\neg\varphi}{t:\neg\varphi}$

- Using the rules of the tableaux calculus we only explore finite, tree models.
- ► Let's assume that the calculus is correct (you will have to believe me).
- ▶ Then the  $\langle R \rangle$ -language
  - cannot say infinite,
  - cannot say non-tree.

$$\frac{s:(\varphi \wedge \psi)}{s:\varphi} \frac{s:\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg(\varphi \wedge \psi)}{s:\neg\varphi} \text{ for } t \text{ a new label}$$

$$\frac{s:\neg(\varphi \wedge \psi)}{s:\neg\varphi} \frac{s:\neg\langle R \rangle \varphi}{sRt} \\ \frac{s:\neg\varphi}{t:\neg\varphi}$$

**Theorem:** A formula in the  $\langle R \rangle$ -language is satisfialle if and only if it is satisfiable in a finite, tree relational structure.

▶ We introduce a tableaux method to check satisfiability for the language with  $\langle R \rangle$ .

- ▶ We introduce a tableaux method to check satisfiability for the language with  $\langle R \rangle$ .
- ▶ We saw that we can use labels to describe what is going on in each point of a relational structure.

- ▶ We introduce a tableaux method to check satisfiability for the language with  $\langle R \rangle$ .
- ▶ We saw that we can use labels to describe what is going on in each point of a relational structure.
- ► More importantly: we saw that tableaux are a way to sistematically explore relational structures.

- ▶ We introduce a tableaux method to check satisfiability for the language with  $\langle R \rangle$ .
- ▶ We saw that we can use labels to describe what is going on in each point of a relational structure.
- More importantly: we saw that tableaux are a way to sistematically explore relational structures.
- Actually, from the tableaux algorithm we could learn some model properties: we only need to consider finite tree models.

Tableaux Algorithms

#### Tableaux Algorithms

➤ The Tableaux Method is the core algorithm of most current theorem provers for relational languages.

#### Tableaux Algorithms

- ➤ The Tableaux Method is the core algorithm of most current theorem provers for relational languages.
- ➤ You might have heard about description logics. Racer, FaCT++, Pellet are all based on a calculus similar to the one we studied today.

#### Tableaux Algorithms

- ➤ The Tableaux Method is the core algorithm of most current theorem provers for relational languages.
- ➤ You might have heard about description logics. Racer, FaCT++, Pellet are all based on a calculus similar to the one we studied today.
- Fitting's Web page: http: //comet.lehman.cuny.edu/fitting/



#### Tableaux Algorithms

- ➤ The Tableaux Method is the core algorithm of most current theorem provers for relational languages.
- ➤ You might have heard about description logics. Racer, FaCT++, Pellet are all based on a calculus similar to the one we studied today.
- Fitting's Web page: http: //comet.lehman.cuny.edu/fitting/



Fitting, Melvin (1983). Proof Methods for Modal and Intuitionistic Logics. D. Reidel Publishing Co., Dordrecht.



Tree and Finite Model Properties

 Unraveling, the procedure to turn arbitrary models into trees, was introduced by Segerberg.

- Unraveling, the procedure to turn arbitrary models into trees, was introduced by Segerberg.
- Segerberg's Web page: http://www.phil.ucalgary.ca/ philosophy/people/segerberg.html



- Unraveling, the procedure to turn arbitrary models into trees, was introduced by Segerberg.
- ➤ Segerberg's Web page: http://www.phil.ucalgary.ca/ philosophy/people/segerberg.html
- ➤ A more general result about turning things into other things can be proved using bisimulations.



- Unraveling, the procedure to turn arbitrary models into trees, was introduced by Segerberg.
- Segerberg's Web page: http://www.phil.ucalgary.ca/ philosophy/people/segerberg.html
- ▶ A more general result about turning things into other things can be proved using bisimulations.
- van Benthem's Web page: http://staff.science.uva.nl/~johan/





Tree and Finite Model Properties

- Unraveling, the procedure to turn arbitrary models into trees, was introduced by Segerberg.
- Segerberg's Web page: http://www.phil.ucalgary.ca/ philosophy/people/segerberg.html
- ▶ A more general result about turning things into other things can be proved using bisimulations.
- van Benthem's Web page: http://staff.science.uva.nl/~johan/






Segerberg, Krister (1971). An Essay in Classical Modal Logic, Department of Philosophy Uppsala University, Sweden. Uppsala Philosophical Studies.



van Benthem, Johan (1985). Modal Logic and Classical Logic, Bibliopolis.





# **Interesting Links**

### Interesting Links

- ▶ Some provers for the  $\langle R \rangle$  language based on the tableaux algorithm:
  - Racer http://www.racer-systems.com/
  - ► FaCT++ http://owl.man.ac.uk/factplusplus/
  - Pellet http://pellet.owldl.com/
  - ► HTab http://trac.loria.fr/projects/htab/wiki

### Interesting Links

- ▶ Some provers for the  $\langle R \rangle$  language based on the tableaux algorithm:
  - Racer http://www.racer-systems.com/
  - ► FaCT++ http://owl.man.ac.uk/factplusplus/
  - ▶ Pellet http://pellet.owldl.com/
  - ► HTab http://trac.loria.fr/projects/htab/wiki
- ▶ Some based on other algorithms:
  - ► MSpass (translation based) http://www.cs.man.ac.uk/~schmidt/mspass/
  - HyLoRes (resolution based) http://trac.loria.fr/projects/hylores

### The Next Lecture

No Way to Say Warm in French