

LogicS

Lecture #9: We like it Complete and Compact (and have a Soft Spot for Löwenheim-Skolem)

Carlos Areces and Patrick Blackburn

`{carlos.areces,patrick.blackburn}@loria.fr`

INRIA Nancy Grand Est
Nancy, France

NASSLLI 2010 - Bloomington - USA

The last one!

- ▶ I want to discuss soundness and completeness

The last one!

- ▶ I want to discuss **soundness** and **completeness**
Intuitively, they are the way to **synchronize** the semantics of a logic with an inference method like tableaux.

The last one!

- ▶ I want to discuss **soundness** and **completeness**
Intuitively, they are the way to **synchronize** the semantics of a logic with an inference method like tableaux.
They are the way to show that **we got it right**.

The last one!

- ▶ I want to discuss **soundness** and **completeness**
Intuitively, they are the way to **synchronize** the semantics of a logic with an inference method like tableaux.
They are the way to show that **we got it right**.
- ▶ I want to discuss also **Compactness** and **Löwenheim-Skolem**.

The last one!

- ▶ I want to discuss **soundness** and **completeness**
Intuitively, they are the way to **synchronize** the semantics of a logic with an inference method like tableaux.
They are the way to show that **we got it right**.
- ▶ I want to discuss also **Compactness** and **Löwenheim-Skolem**.
Properties that actually **characterize** first order logic:
Lindström Theorem.

The last one!

- ▶ I want to discuss **soundness** and **completeness**
Intuitively, they are the way to **synchronize** the semantics of a logic with an inference method like tableaux.
They are the way to show that **we got it right**.
- ▶ I want to discuss also **Compactness** and **Löwenheim-Skolem**.
Properties that actually **characterize** first order logic:
Lindström Theorem.
- ▶ And we will finish with **Second Order Logic**.

Not called classical logic for nothing

- ▶ Some logicians (notably Quine) view first-order logic as the be-all and end-all of logic.
- ▶ We disagree with this viewpoint, but there is no getting away from a very stubborn fact — first-order logic certainly is *special* (and *beautiful*).
- ▶ Being “first-order logic” is not a matter of notation or symbolism. It’s about something much deeper. It’s about finally being able to get to grips with each and every element in the domains of our models.
- ▶ And as we shall learn today, this has some deep consequences. Today we’re going to take a (rather abstract) look at inference and expressivity in first-order logic.

In today's lecture

- ▶ First we'll look at inference, and discuss the concepts of **soundness** and **completeness**.
- ▶ But then we turn to the heart of the lecture: expressivity. We have clearly gained a lot of expressivity, but interesting gaps remain. We will discuss two results, the **Compactness Theorem** and the **Löwenheim-Skolem Theorem(s)** that pin down the crucial expressive limitations of first-order logic. Indeed, as we shall learn, these results actually **characterize** first-order logic.

To Infinity and Beyond...?!



- The distinction between finiteness and the different degrees of infinity will play an important role in the later part of this lecture.

To Infinity and Beyond...?!



- ▶ The distinction between finiteness and the different degrees of infinity will play an important role in the later part of this lecture.
- ▶ Will we get to infinity? Will we get beyond?

Example: Theory of linear order

We now have a lot of expressive power — enough to state some interesting theories. Here, for example, is a simple theory of linear order:

- ▶ Axiom Irr: $\llbracket x \rrbracket (x : \neg \langle R \rangle x)$
- ▶ Axiom Tran: $\llbracket x \rrbracket \llbracket y \rrbracket (x : \langle R \rangle \langle R \rangle y \rightarrow x : \langle R \rangle y)$
- ▶ Axiom Lin: $\llbracket x \rrbracket \llbracket y \rrbracket (x : y \vee x : \langle R \rangle y \vee y : \langle R \rangle x)$

So if we had a proof system for our first-order language, we could prove some non-trivial theorems:

$$\models \textit{Irr} \wedge \textit{Tran} \wedge \textit{Lin} \rightarrow \llbracket x \rrbracket \llbracket y \rrbracket (x : \langle R \rangle y \rightarrow y : \neg \langle R \rangle x).$$

Proof theory

- ▶ Proof theory is the **syntactic** approach to logic.
- ▶ It attempts to define collections of rules and/or axioms that enable us to generate new formulas from old. That is, it attempts to pin down the notion of inference syntactically.
- ▶ Given some proof system P , we write $\vdash_P \varphi$ to indicate that a formula φ is provable in the the proof system. ($\nvdash_P \varphi$ means that φ is *not* provable in proof system P .).

Many types of proof system

- ▶ Natural deduction
- ▶ Hilbert-style system (often called axiomatic systems)
- ▶ Sequent calculus
- ▶ Tableaux systems
- ▶ Resolution

Why so many different proof systems?

- ▶ Well, one of the most important reasons may simply be that logicians love to play with such systems — and every logician has his or her own favourite pet system!
- ▶ A more serious reason is: different proof systems are typically good for different purposes.
- ▶ In particular, some systems (notably tableau and resolution) are particularly suitable for computational purposes.

But what does all this have to do with semantics and inference?

- ▶ Note: **nothing** we have said so far about proof systems makes any connection with the model-theoretic ideas previously introduced.
- ▶ All we have done is talk about provability and vaguely said that we want to “generate” formulas syntactically. What does this have to do with relational structures and semantics?

But what does all this have to do with semantics and inference?

- ▶ Note: **nothing** we have said so far about proof systems makes any connection with the model-theoretic ideas previously introduced.
- ▶ All we have done is talk about provability and vaguely said that we want to “generate” formulas syntactically. What does this have to do with relational structures and semantics?
- ▶ Answer: we insist on working with proof systems with two special properties, namely **soundness** and **completeness**.

Soundness

- ▶ Recall that we write $\models \varphi$ to indicate that the formula φ is valid (that is, satisfied in all models under all assignments).
- ▶ Recall that we write $\vdash_P \varphi$ to indicate that φ is provable in proof system P .
- ▶ We say that a proof system P is **sound** if and only if
$$\vdash_P \varphi \text{ implies } \models \varphi.$$

Soundness

- ▶ Recall that we write $\models \varphi$ to indicate that the formula φ is valid (that is, satisfied in all models under all assignments).
- ▶ Recall that we write $\vdash_P \varphi$ to indicate that φ is provable in proof system P .
- ▶ We say that a proof system P is **sound** if and only if
$$\vdash_P \varphi \text{ implies } \models \varphi.$$
- ▶ That is, soundness means that syntactic provability implies semantic validity. To put it another way: **P does not produce garbage.**
- ▶ Needless to say, all the standard proof systems are sound.

Proving Soundness

- ▶ Soundness is typically an easy property to prove.
- ▶ Proofs typically have some kind of inductive structure. One shows that if the first part of proof is true in a model, then the rules only let us generate formulas that are also true in a model.

Completeness

- ▶ Recall that we write $\models \varphi$ to indicate that the formula φ is valid (that is, satisfied everywhere in all models).
- ▶ Recall that we write $\vdash_P \varphi$ to indicate that φ is provable in proof system P .
- ▶ We say that a proof system p is **complete** if and only if $\models \varphi$ implies $\vdash_P \varphi$.

Completeness

- ▶ Recall that we write $\models \varphi$ to indicate that the formula φ is valid (that is, satisfied everywhere in all models).
- ▶ Recall that we write $\vdash_P \varphi$ to indicate that φ is provable in proof system P .
- ▶ We say that a proof system p is **complete** if and only if
$$\models \varphi \text{ implies } \vdash_P \varphi.$$
- ▶ That is, completeness means that our proof system is strong enough to prove everything that is provable.
- ▶ To put it another way: **No valid formula is out of reach of our proof system.**
- ▶ The standard proof systems are complete.

Remark

- ▶ Completeness is a **much more informative** property than soundness, and is a lot more difficult to prove.
- ▶ It is typically proved by contraposition. **We show that if some formula is not provable ($\not\vdash \varphi$) then φ is not valid ($\not\models \varphi$).** This is done by building a model for $\neg\varphi$.
- ▶ And our first-order language is very expressive — so it can describe some pretty intricate models (they certainly won't all be trees!).

Soundness and completeness together

- Recall: proof system P is sound if and only if

$$\vdash_P \varphi \text{ implies } \models \varphi$$

- Proof system P is complete if and only if

$$\models \varphi \text{ implies } \vdash_P \varphi$$

- So if a proof system is both sound and complete (which is what we want) we have that:

$$\models \varphi \text{ if and only if } \vdash_P \varphi$$

- That is, syntactic provability and semantic validity coincide.

Expressivity of first-order logic

- ▶ We turn now to the theme of **expressivity of first-order logic**.
- ▶ Our discussion revolves around two famous results: the **Compactness Theorem**, and the **Löwenheim Skolem Theorem(s)**. I'm going to state (but won't prove) these results and discuss some of their consequences.
- ▶ As we said at the start of the lecture, our discussion will have a lot to do with **finiteness**, and the **different grades of infinity**. So before going any further let's remind ourselves what an infinite set is ...

Finite and Infinite

Finite and Infinite

- ▶ The most fundamental infinite set in set theory is the natural numbers \mathbb{N} . This is the set

$$\{0, 1, 2, 3, 4, 5, \dots\}.$$

Finite and Infinite

- ▶ The most fundamental infinite set in set theory is the natural numbers \mathbb{N} . This is the set

$$\{0, 1, 2, 3, 4, 5, \dots\}.$$

- ▶ A set S is **infinite** if there is one-to-one (injective) function from \mathbb{N} to S .

Finite and Infinite

- ▶ The most fundamental infinite set in set theory is the natural numbers \mathbb{N} . This is the set

$$\{0, 1, 2, 3, 4, 5, \dots\}.$$

- ▶ A set S is **infinite** if there is one-to-one (injective) function from \mathbb{N} to S .
- ▶ A set is **finite** if it is not infinite. That is, a set is finite if it is not possible to define such a function. This amounts to saying that it is the same size as some finite set

$$\{1, 2, 3, 4, 5, \dots, n\}$$

.

Infinite Axiom Sets!



- ▶ We already seen a simple set of axioms, namely the three axioms that defined the theory of linear order.
- ▶ But note: **any finite set of axioms can be replaced by one single axiom** — for we simply need to form their conjunction!
- ▶ So, very early in the history of logic, logicians started to consider the consequences of working with infinite sets of axioms. After all, why not? **Remember: Logicians are the Masters of the Universe!**
- ▶ Now, using infinite sets of axioms **does** give us more power. . .

An infinite set of axioms defining infinity!

An infinite set of axioms defining infinity!

- ▶ Try to write down a **single sentence** of first-order logic that is true in every infinite model (and false on every finite one). But don't try too hard, for I promise that you won't find one. . .

An infinite set of axioms defining infinity!

- ▶ Try to write down a **single sentence** of first-order logic that is true in every infinite model (and false on every finite one). But don't try too hard, for I promise that you won't find one. . .
- ▶ But it's easy to write down an **infinite** set of sentences with this property.

An infinite set of axioms defining infinity!

- ▶ Try to write down a **single sentence** of first-order logic that is true in every infinite model (and false on every finite one). But don't try too hard, for I promise that you won't find one. . .
- ▶ But it's easy to write down an **infinite** set of sentences with this property.
- ▶ Let ***AtLeast*₂** be $\langle\langle x \rangle\langle y \rangle\rangle(x:\neg y)$.

An infinite set of axioms defining infinity!

- ▶ Try to write down a **single sentence** of first-order logic that is true in every infinite model (and false on every finite one). But don't try too hard, for I promise that you won't find one. . .
- ▶ But it's easy to write down an **infinite** set of sentences with this property.
- ▶ Let ***AtLeast*₂** be $\langle\langle x \rangle\langle y \rangle\rangle(x:\neg y)$.
- ▶ Let ***AtLeast*₃** be $\langle\langle x \rangle\langle y \rangle\langle z \rangle\rangle(x:\neg y \wedge x:\neg z \wedge y:\neg z)$.

An infinite set of axioms defining infinity!

- ▶ Try to write down a **single sentence** of first-order logic that is true in every infinite model (and false on every finite one). But don't try too hard, for I promise that you won't find one. . .
- ▶ But it's easy to write down an **infinite** set of sentences with this property.
- ▶ Let *AtLeast*₂ be $\langle\langle x \rangle\langle y \rangle\rangle(x:\neg y)$.
- ▶ Let *AtLeast*₃ be $\langle\langle x \rangle\langle y \rangle\langle z \rangle\rangle(x:\neg y \wedge x:\neg z \wedge y:\neg z)$.
- ▶ Following this pattern, we can write down sentences *AtLeast*₄, *AtLeast*₅, *AtLeast*₆, *AtLeast*₇, . . . , *AtLeast*₁₀₀,

An infinite set of axioms defining infinity!

- ▶ Try to write down a **single sentence** of first-order logic that is true in every infinite model (and false on every finite one). But don't try too hard, for I promise that you won't find one. . .
- ▶ But it's easy to write down an **infinite** set of sentences with this property.
- ▶ Let *AtLeast*₂ be $\langle\langle x \rangle\langle y \rangle\rangle(x:\neg y)$.
- ▶ Let *AtLeast*₃ be $\langle\langle x \rangle\langle y \rangle\langle z \rangle\rangle(x:\neg y \wedge x:\neg z \wedge y:\neg z)$.
- ▶ Following this pattern, we can write down sentences *AtLeast*₄, *AtLeast*₅, *AtLeast*₆, *AtLeast*₇, . . . , *AtLeast*₁₀₀,
- ▶ Let **INF** be the set of all these sentences. Then we have that

$\mathcal{M} \models \mathbf{INF}$ iff \mathcal{M} is an infinite model.

Compactness Theorem

Compactness Theorem: Let Σ be an infinite set of first-order sentences. If every finite subset of Σ can be made true (in some model or other) then there is at least one model that simultaneously makes every sentence in Σ true.

We can paraphrase this as follows. To check whether an infinite set of first-order sentences Σ , has a model, we **don't** need to try and find a model that make all the sentences in Σ true at once. If it enough to show that any finite subset of Σ can be made true in some model. For if we can show this, the Compactness Theorem guarantees that there is some model that makes all the (infinitely many) sentences in Σ true all at once.

How do you prove the Compactness Theorem?

Many proofs are known, but two are worth mentioning. . .

- ▶ Actually, one can prove the Compactness Theorem for first-order logic more-or-less simultaneously with the Completeness Theorem. More precisely, completeness for first-order logic is a reasonably simple extension of compactness.
- ▶ And there is another (in a sense more revealing) proof. The ultraproduct construction lets us “multiply together” the finite models into one big model.

A powerful theorem

The Compactness Theorem is central to mathematical model theory:

- ▶ A powerful theorem — and a two-sided one.
- ▶ On the positive side, it allows us to build many interesting and unexpected models (such as non-standard models of arithmetic).
- ▶ And it has negative uses too — it can also show that we cannot define certain things. *And that's the kind use we will put it now.*

Finiteness is not first-order definable

The question we will ask is the following:

*Is there a **single sentence** of first-order logic that defines finiteness?*

That is, is there a single first order sentence (let's call it **fin**) such that:

$$\mathcal{M} \models \mathbf{fin} \text{ iff } \mathcal{M} \text{ is finite}$$

As we shall see, the answer is **no**.

The argument

The argument

- ▶ Suppose there is such a sentence **fin**. We are going to show (with the help of the Compactness Theorem) that this assumption leads to a contradiction.

The argument

- ▶ Suppose there is such a sentence **fin**. We are going to show (with the help of the Compactness Theorem) that this assumption leads to a contradiction.
- ▶ Consider the set of sentences **INF** \cup **fin**

The argument

- ▶ Suppose there is such a sentence **fin**. We are going to show (with the help of the Compactness Theorem) that this assumption leads to a contradiction.
- ▶ Consider the set of sentences **INF** \cup **fin**
- ▶ That is: $\{AtLeast_1, AtLeast_2, AtLeast_3, AtLeast_4, \dots\} \cup \mathbf{fin}$

The argument

- ▶ Suppose there is such a sentence **fin**. We are going to show (with the help of the Compactness Theorem) that this assumption leads to a contradiction.
- ▶ Consider the set of sentences **INF** \cup **fin**
- ▶ That is: $\{AtLeast_1, AtLeast_2, AtLeast_3, AtLeast_4, \dots\} \cup \mathbf{fin}$
- ▶ Claim: every finite subset of this has a model. Why is this?

And now we have a problem

And now we have a problem

- ▶ Since every finite subset of $\mathbf{INF} \cup \mathbf{fin}$ is true, by the Compactness Theorem, there is some model \mathcal{M} that simultaneously makes every sentence in $\mathbf{INF} \cup \mathbf{fin}$

And now we have a problem

- ▶ Since every finite subset of $\mathbf{INF} \cup \mathbf{fin}$ is true, by the Compactness Theorem, there is some model \mathcal{M} that simultaneously makes every sentence in $\mathbf{INF} \cup \mathbf{fin}$
- ▶ Hence, as \mathcal{M} makes \mathbf{INF} true, \mathcal{M} is infinite.

And now we have a problem

- ▶ Since every finite subset of **INF** \cup **fin** is true, by the Compactness Theorem, there is some model \mathcal{M} that simultaneously makes every sentence in **INF** \cup **fin**
- ▶ Hence, as \mathcal{M} makes **INF** true, \mathcal{M} is infinite.
- ▶ But as \mathcal{M} also makes **fin** true, so \mathcal{M} is finite!

And now we have a problem

- ▶ Since every finite subset of **INF** \cup **fin** is true, by the Compactness Theorem, there is some model \mathcal{M} that simultaneously makes every sentence in **INF** \cup **fin**
- ▶ Hence, as \mathcal{M} makes **INF** true, \mathcal{M} is infinite.
- ▶ But as \mathcal{M} also makes **fin** true, so \mathcal{M} is finite!
- ▶ Ooooooooooops!!!!!!!!!!!!

And now we have a problem

- ▶ Since every finite subset of **INF** \cup **fin** is true, by the Compactness Theorem, there is some model \mathcal{M} that simultaneously makes every sentence in **INF** \cup **fin**
- ▶ Hence, as \mathcal{M} makes **INF** true, \mathcal{M} is infinite.
- ▶ But as \mathcal{M} also makes **fin** true, so \mathcal{M} is finite!
- ▶ Ooooooooooops!!!!!!!!!!!!
- ▶ From this contradiction we deduce that **fin** does not exist. That is, there is no sentence of first-order logic that expresses the concept of finiteness.

Countably infinite and bigger...

Countably infinite and bigger...

- ▶ A set S is **countably infinite** if there is one-to-one (injective) and onto (surjective) function from \mathbf{N} to S .

Countably infinite and bigger...

- ▶ A set S is **countably infinite** if there is one-to-one (injective) and onto (surjective) function from \mathbf{N} to S .
- ▶ Countable infinite sets are the smallest infinite sets. Examples of countable infinite sets include: \mathbf{N} , \mathbb{E} (the even numbers), \mathbb{Z} (the integers), and \mathbb{Q} (the fractions).

Countably infinite and bigger...

- ▶ A set S is **countably infinite** if there is one-to-one (injective) and onto (surjective) function from \mathbf{N} to S .
- ▶ Countable infinite sets are the smallest infinite sets. Examples of countable infinite sets include: \mathbf{N} , \mathbb{E} (the even numbers), \mathbb{Z} (the integers), and \mathbb{Q} (the fractions).
- ▶ The set of real numbers \mathbb{R} is **not** countably infinite — it is bigger.

Countably infinite and bigger...

- ▶ A set S is **countably infinite** if there is one-to-one (injective) and onto (surjective) function from \mathbf{N} to S .
- ▶ Countable infinite sets are the smallest infinite sets. Examples of countable infinite sets include: \mathbf{N} , \mathbb{E} (the even numbers), \mathbb{Z} (the integers), and \mathbb{Q} (the fractions).
- ▶ The set of real numbers \mathbb{R} is **not** countably infinite — it is bigger.
- ▶ In general, the power set $\mathcal{P}(S)$ of S is bigger than S .

Countably infinite and bigger...

- ▶ A set S is **countably infinite** if there is one-to-one (injective) and onto (surjective) function from \mathbf{N} to S .
- ▶ Countable infinite sets are the smallest infinite sets. Examples of countable infinite sets include: \mathbf{N} , \mathbb{E} (the even numbers), \mathbb{Z} (the integers), and \mathbb{Q} (the fractions).
- ▶ The set of real numbers \mathbb{R} is **not** countably infinite — it is bigger.
- ▶ In general, the power set $\mathcal{P}(S)$ of S is bigger than S .
- ▶ The universe of sets is huge. It contains infinitely many ever-bigger grades of infinity.

Löwenheim-Skolem Theorems

The Löwenheim Skolem Theorems tell us that first-order logic is completely blind to all these distinctions:

Upward Löwenheim-Skolem Theorem: *If a set of sentences Σ has an infinite model, it has infinite models of all larger infinite cardinalities.*

Downward Löwenheim-Skolem Theorem: *If a set of sentences Σ has an infinite model, it has infinite models of all lower infinite cardinalities. In particular, it always has a countable model.*

It follows that it is not possible to fully describe the real numbers \mathbb{R} (that is, describe them up to isomorphism) using first-order logic. Any description will have many models (indeed models of every infinite cardinality) and a model that (being countable) is too small!

Lindström's Theorem

- ▶ There is a field of logic called **abstract model theory** which works with very abstract definitions of what logics are.
- ▶ There is a celebrated result from abstract model theory called **Lindström's Theorem** which tells us that first-order logic is the **only** logic for which both the Compactness and the Downward Löwenheim-Skolem Theorems hold.
- ▶ That is, if you invent a very strange logical formalism, but can prove that it has these two properties, then you have invented a logic with exactly the same expressive power as first-order logic. You “new” logic, when you get right down to it, is in the business of quantifying over individuals.
- ▶ **And is there really anything else out there to quantify over....?**

Sets, and Relations, and Functions, and

- ▶ Aren't we already quantifying over everything that there is in our models?
- ▶ The answer is **no**. There's a lot more sitting out there in our models, patiently waiting to be quantified.
- ▶ Sure, we're already quantifying over the individuals — but there are **higher-order** entities there too, such as **sets of individuals**, and **relations**.
- ▶ And these logics certainly **do** offer increased expressivity. . .

Transitive closure

- ▶ We already met the concept of the reflexive transitive closure of a relation.
- ▶ There are two (equivalent) ways of defining reflexive transitive closure.
 - ▶ As the relation T on D defined by xTy iff either $x = y$ or there is a finite sequence of elements of D such that $x = d_0$ and

$$d_0 R d_1, d_1 R d_2, \dots, d_{n-1} R d_n.$$

- ▶ As the **smallest** reflexive and transitive relation S (on the domain D) containing an arbitrary relation R ; or
- ▶ Let's try defining this concept with $\llbracket x \rrbracket$ and $\llbracket x \rrbracket \dots$

Let's try...

Let R and S be binary relations

Let's try...

Let R and S be binary relations. It's easy to insist that R is reflexive:

$$\text{Ref}(S) := \llbracket n \rrbracket (n : \langle S \rangle n).$$

Let's try...

Let R and S be binary relations. It's easy to insist that R is reflexive:

$$\text{Ref}(S) := \llbracket n \rrbracket (n : \langle S \rangle n).$$

And it's easy to say that R is transitive:

$$\text{Tran}(S) := \llbracket n \rrbracket \llbracket m \rrbracket (n : \langle S \rangle \langle S \rangle m \rightarrow n : \langle S \rangle n).$$

Let's try...

Let R and S be binary relations. It's easy to insist that R is reflexive:

$$\text{Ref}(S) := \llbracket n \rrbracket (n : \langle S \rangle n).$$

And it's easy to say that R is transitive:

$$\text{Tran}(S) := \llbracket n \rrbracket \llbracket m \rrbracket (n : \langle S \rangle \langle S \rangle m \rightarrow n : \langle S \rangle n).$$

And to say that R is a subrelation of S

$$R \subseteq S := \llbracket n \rrbracket \llbracket m \rrbracket (n : \langle R \rangle m \rightarrow n : \langle S \rangle n).$$

Here's a first try...

So let's put all this together to define transitive reflexive closure:

$$\mathit{Tran}^*(R, S)$$

Here's a first try...

So let's put all this together to define transitive reflexive closure:

$$\mathit{Tran}^*(R, S) \quad := \quad \mathit{Ref}(S)$$

Here's a first try...

So let's put all this together to define transitive reflexive closure:

$$\begin{aligned} Tran^*(R, S) \quad &:= \quad Ref(S) \\ &\quad \wedge \quad Tran(S) \end{aligned}$$

Here's a first try...

So let's put all this together to define transitive reflexive closure:

$$\begin{aligned} \text{Tran}^*(R, S) \quad &:= \quad \text{Ref}(S) \\ &\wedge \text{Tran}(S) \\ &\wedge R \subseteq S \end{aligned}$$

Here's a first try...

So let's put all this together to define transitive reflexive closure:

$$\begin{aligned} \text{Tran}^*(R, S) \quad &:= \quad \text{Ref}(S) \\ &\wedge \text{Tran}(S) \\ &\wedge R \subseteq S \\ &\wedge \textit{S-is-the-smallest-such-relation} \end{aligned}$$

Here's a first try...

So let's put all this together to define transitive reflexive closure:

$$\begin{aligned} Tran^*(R, S) \quad &:= \quad Ref(S) \\ &\wedge Tran(S) \\ &\wedge R \subseteq S \\ &\wedge \textit{S-is-the-smallest-such-relation} \end{aligned}$$

Oh dear...!

No way José!

- ▶ Try as you might, you won't be able to do this
- ▶ And we can prove it using the Compactness Theorem
- ▶ $\{\neg p, [R]\neg p, [R][R]\neg p, [R][R][R]\neg p, \dots, \langle R^* \rangle p\}$
- ▶ Every finite subset has a model. Hence (by Compactness) so does the whole thing. But this is impossible.
- ▶ Hence we cannot define R^* .

No way José!

- ▶ Try as you might, you won't be able to do this
- ▶ And we can prove it using the Compactness Theorem
- ▶ $\{\neg p, [R]\neg p, [R][R]\neg p, [R][R][R]\neg p, \dots, \langle R^* \rangle p\}$
- ▶ Every finite subset has a model. Hence (by Compactness) so does the whole thing. But this is impossible.
- ▶ Hence we cannot define R^* .
- ▶ But as we know, we're free to extend the language.
- ▶ We could just add the $\langle R^* \rangle$ operator — but that was just one example of something we couldn't do.
- ▶ Let's give ourselves the power to quantify over two types of higher order entities: **properties** and **binary relations**.

A second order language

- ▶ $\langle\langle p \rangle\rangle\varphi$, and $\llbracket p \rrbracket\varphi$ express existential and universal quantification over properties.
- ▶ $\langle\langle R \rangle\rangle\varphi$, and $\llbracket R \rrbracket\varphi$ express existential and universal quantification over relations.
- ▶ Semantics? Simply extend what we did in first-order case.

Now we can define reflexive transitive closure. . .

$$\mathit{Tran}^*(R, S)$$

Now we can define reflexive transitive closure. . .

$$\mathit{Tran}^*(R, S) \quad := \quad \mathit{Ref}(S)$$

Now we can define reflexive transitive closure. . .

$$\begin{aligned} \textit{Tran}^*(R, S) \quad &:= \quad \textit{Ref}(S) \\ &\quad \wedge \textit{Tran}(S) \end{aligned}$$

Now we can define reflexive transitive closure. . .

$$\begin{aligned} \textit{Tran}^*(R, S) \quad &:= \quad \textit{Ref}(S) \\ &\wedge \textit{Tran}(S) \\ &\wedge R \subseteq S \end{aligned}$$

Now we can define reflexive transitive closure. . .

$$\begin{aligned} \text{Tran}^*(R, S) \quad &:= \quad \text{Ref}(S) \\ &\wedge \text{Tran}(S) \\ &\wedge R \subseteq S \\ &\wedge \llbracket X \rrbracket (\text{Ref}(X) \wedge \text{Tran}(X) \wedge R \subseteq X \rightarrow S \subseteq X). \end{aligned}$$

What's the Price

- ▶ Loss of Completeness (for standard models)
- ▶ Loss of Compactness. After all:

$$\{\neg p, [R]\neg p, [R][R]\neg p, [R][R][R]\neg p, \dots, \langle R^* \rangle p\}$$

is now an example of a set in which each finite subset has a model, and the complete set doesn't.

- ▶ Loss of Löwenheim-Skolem. (It is easy to define the natural number \mathbb{N} and the integers \mathbb{Z} up to isomorphism.)

What's the Price

- ▶ Loss of Completeness (for standard models)
- ▶ Loss of Compactness. After all:

$$\{\neg p, [R]\neg p, [R][R]\neg p, [R][R][R]\neg p, \dots, \langle R^* \rangle p\}$$

is now an example of a set in which each finite subset has a model, and the complete set doesn't.

- ▶ Loss of Löwenheim-Skolem. (It is easy to define the natural number \mathbb{N} and the integers \mathbb{Z} up to isomorphism.)
- ▶ Which brings us back to the fundamental trade-off, expressivity versus inference/tractability.
- ▶ We've bought serious expressivity — and have lost everything else.

What we covered in the course

- ▶ We've been essentially looking at a menu of logics.
- ▶ But the menu was designed by a Master Chef (Tarski!); the meal is built around the crucial ingredient of relational structures.
- ▶ Relational structures tell us why logic is applicable in semantics (natural language metaphysics) and computer science.
- ▶ Back to a logicist position, but not in traditional sense.
- ▶ Monotheist — but not in terms of logic, rather, in terms of semantics.

Relevant Bibliography



Georg Cantor (1845-1918) created modern set theory, which is the setting for virtually all of modern mathematics. As David Hilbert, the famous mathematician and logician once remarked, “No one shall expel us from the Paradise that Cantor has created.”

Relevant Bibliography



The Completeness Theorem for first-order logic was proved by Kurt Gödel (1906–1978) in his 1928 doctoral thesis. As it turned out, however, this was merely the first of many great results that he was to prove.

Relevant Bibliography



The modern form of the Compactness Theorem seems to trace back to the work of the Russian mathematician Anatoly Maltsev (1909–1967).

Relevant Bibliography



The downward Löwenheim-Skolem Theorem was first proved by Leopold Löwenheim (1878–1957; top picture) in 1915. In 1920 his proof was greatly simplified and generalised by Thoralf Skolem (1887 -1963).