

# Logics for Computation

## Lecture #9: Putting Tiles in an Infinite Bathroom

Carlos Areces and Patrick Blackburn

`{carlos.areces,patrick.blackburn}@loria.fr`

INRIA Nancy Grand Est  
Nancy, France

ESSLLI 2008 - Hamburg - Germany

# The Story so Far

- ▶ We are working with first-order logic.

# The Story so Far

- ▶ We are working with first-order logic.
- ▶ We discussed **soundness** and **completeness**

# The Story so Far

- ▶ We are working with first-order logic.
- ▶ We discussed **soundness** and **completeness**  
Intuitively, they are the way to **synchronize** the semantics of a logic with an inference method like tableaux.

# The Story so Far

- ▶ We are working with first-order logic.
- ▶ We discussed **soundness** and **completeness**  
Intuitively, they are the way to **synchronize** the semantics of a logic with an inference method like tableaux.  
They are the way to show that **we got it right**.

# The Story so Far

- ▶ We are working with first-order logic.
- ▶ We discussed **soundness** and **completeness**  
Intuitively, they are the way to **synchronize** the semantics of a logic with an inference method like tableaux.  
They are the way to show that **we got it right**.
- ▶ We also discussed some properties like **Compactness** and **Löwenheim-Skolem**.

# The Story so Far

- ▶ We are working with first-order logic.
- ▶ We discussed **soundness** and **completeness**  
Intuitively, they are the way to **synchronize** the semantics of a logic with an inference method like tableaux.  
They are the way to show that **we got it right**.
- ▶ We also discussed some properties like **Compactness** and **Löwenheim-Skolem**.
- ▶ We saw that these properties actually **characterize first order logic**: Lindström Theorem.

# The Story so Far

- ▶ We are working with first-order logic.
- ▶ We discussed **soundness** and **completeness**  
Intuitively, they are the way to **synchronize** the semantics of a logic with an inference method like tableaux.  
They are the way to show that **we got it right**.
- ▶ We also discussed some properties like **Compactness** and **Löwenheim-Skolem**.
- ▶ We saw that these properties actually **characterize first order logic**: Lindström Theorem.
- ▶ We've come a long way. Think that on Monday we were in propositional logic!



# What do we do Today

# What do we do Today

- ▶ We'll talk about the satisfiability problem of first-order logic.

# What do we do Today

- ▶ We'll talk about the satisfiability problem of first-order logic.
- ▶ We will argue that the problem is **undecidable**

# What do we do Today

- ▶ We'll talk about the satisfiability problem of first-order logic.
- ▶ We will argue that the problem is **undecidable**
  - ▶ That is, there is no algorithm that can answer for any formula of first order logic, whether the formula has a model or not.

# What do we do Today

- ▶ We'll talk about the satisfiability problem of first-order logic.
- ▶ We will argue that the problem is **undecidable**
  - ▶ That is, there is no algorithm that can answer for any formula of first order logic, whether the formula has a model or not.
- ▶ Actually, we are going to show how to **tile and infinite bathroom**.

---

# Tableaux for First Order Logic

# Tableaux for First Order Logic

- Patrick introduced these two rules yesterday:

$$\frac{s:\langle x \rangle \varphi}{s:\varphi[x \leftarrow n]} (\langle x \rangle) \qquad \frac{s:\neg \langle x \rangle \varphi}{s:\neg \varphi[x \leftarrow o]} (\neg \langle x \rangle)$$

(where  $n$  is a new name)      (where  $o$  is an old name)

# Tableaux for First Order Logic

- ▶ Patrick introduced these two rules yesterday:

$$\frac{s:\langle x \rangle \varphi}{s:\varphi[x \leftarrow n]} (\langle x \rangle) \qquad \frac{s:\neg \langle x \rangle \varphi}{s:\neg \varphi[x \leftarrow o]} (\neg \langle x \rangle)$$

(where  $n$  is a new name)      (where  $o$  is an old name)

- ▶ How comes that this tableaux do not terminate?



# Tableaux for First Order Logic

- ▶ Patrick introduced these two rules yesterday:

$$\frac{s:\langle x \rangle \varphi}{s:\varphi[x \leftarrow n]} (\langle x \rangle) \qquad \frac{s:\neg \langle x \rangle \varphi}{s:\neg \varphi[x \leftarrow o]} (\neg \langle x \rangle)$$

(where  $n$  is a new name)      (where  $o$  is an old name)

- ▶ How comes that this tableaux do not terminate?
- ▶ As we already mentioned,  
The SAT problem of FO is undecidable.
- ▶ Hence, we cannot expect any sound and complete tableaux to also terminate.

# (Un)Decidability

# (Un)Decidability

How can we prove that problem X is **undecidable**?

# (Un)Decidability

How can we prove that problem X is **undecidable**?

One way is

- ▶ Ask somebody (more intelligent than us) to prove that some problem Y is undecidable

# (Un)Decidability

How can we prove that problem X is **undecidable**?

One way is

- ▶ Ask somebody (more intelligent than us) to prove that some problem Y is undecidable
- ▶ Prove that if X would be decidable then Y would be decidable, giving a codification of Y into X.

# (Un)Decidability

How can we prove that problem X is **undecidable**?

One way is

- ▶ Ask somebody (more intelligent than us) to prove that some problem Y is undecidable
- ▶ Prove that if X would be decidable then Y would be decidable, giving a codification of Y into X.

The **halting problem** of Turing machines is the standard example of an undecidable problem. The behaviour of a Turing machine, and the predicate that says that a given turing machine stops on all inputs can be expressed in FO.

# Tiling Problems

# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle




# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side


# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side
- ▶ for example, here we have six different kinds of tiles:



# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side
- ▶ for example, here we have six different kinds of tiles:




- ▶ a simple tiling problem, could be:

*Is it possible to place tiles of the kind we show above on a grid of  $2 \times 4$ , in such a way that we cover the entire grid and that adjacent tiles have the same color on neighboring sides?*

# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side
- ▶ for example, here we have six different kinds of tiles:




- ▶ a simple tiling problem, could be:

*Is it possible to place tiles of the kind we show above on a grid of  $2 \times 4$ , in such a way that we cover the entire grid and that adjacent tiles have the same color on neighboring sides?*




# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side
- ▶ for example, here we have six different kinds of tiles:




- ▶ a simple tiling problem, could be:

*Is it possible to place tiles of the kind we show above on a grid of  $2 \times 4$ , in such a way that we cover the entire grid and that adjacent tiles have the same color on neighboring sides?*



# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side
- ▶ for example, here we have six different kinds of tiles:




- ▶ a simple tiling problem, could be:

*Is it possible to place tiles of the kind we show above on a grid of  $2 \times 4$ , in such a way that we cover the entire grid and that adjacent tiles have the same color on neighboring sides?*




# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side
- ▶ for example, here we have six different kinds of tiles:




- ▶ a simple tiling problem, could be:

*Is it possible to place tiles of the kind we show above on a grid of  $2 \times 4$ , in such a way that we cover the entire grid and that adjacent tiles have the same color on neighboring sides?*




# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side
- ▶ for example, here we have six different kinds of tiles:



- ▶ a simple tiling problem, could be:


*Is it possible to place tiles of the kind we show above on a grid of  $2 \times 4$ , in such a way that we cover the entire grid and that adjacent tiles have the same color on neighboring sides?*






# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side
- ▶ for example, here we have six different kinds of tiles:




- ▶ a simple tiling problem, could be:

*Is it possible to place tiles of the kind we show above on a grid of  $2 \times 4$ , in such a way that we cover the entire grid and that adjacent tiles have the same color on neighboring sides?*




# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side
- ▶ for example, here we have six different kinds of tiles:




- ▶ a simple tiling problem, could be:

*Is it possible to place tiles of the kind we show above on a grid of  $2 \times 4$ , in such a way that we cover the entire grid and that adjacent tiles have the same color on neighboring sides?*




# Tiling Problems

- ▶ A **tiling problem** is a kind of jigsaw puzzle
- ▶ a **tile**  $T$  is a  $1 \times 1$  square, fixed in orientation, with a fixed **color** in each side
- ▶ for example, here we have six different kinds of tiles:



- ▶ a simple tiling problem, could be:

*Is it possible to place tiles of the kind we show above on a grid of  $2 \times 4$ , in such a way that we cover the entire grid and that adjacent tiles have the same color on neighboring sides?*



# Tiling Problems

# Tiling Problems

- ▶ The general form of a tiling problem  
*Given a finite number of kind of tiles  $\mathcal{T}$ , can we cover a given part of  $\mathbb{Z} \times \mathbb{Z}$  in such a way that adjacent tiles have the same color on the neighboring sides?*

# Tiling Problems


- ▶ The general form of a tiling problem  
*Given a finite number of kind of tiles  $\mathcal{T}$ , can we cover a given part of  $\mathbb{Z} \times \mathbb{Z}$  in such a way that adjacent tiles have the same color on the neighboring sides?*
- ▶ In some cases, it is also possible to impose certain conditions on what is considered a correct tiling.

# Tiling Problems

- ▶ The general form of a tiling problem  
*Given a finite number of kind of tiles  $\mathcal{T}$ , can we cover a given part of  $\mathbb{Z} \times \mathbb{Z}$  in such a way that adjacent tiles have the same color on the neighboring sides?*
- ▶ In some cases, it is also possible to impose certain conditions on what is considered a correct tiling.

- ▶ Covering  $\mathbb{N} \times \mathbb{N}$

- ▶ **tiling  $\mathbb{N} \times \mathbb{N}$ :** Given a finite set of tiles  $\mathcal{T}$ , can  $\mathcal{T}$  cover  $\mathbb{N} \times \mathbb{N}$ ?
- ▶ this problem is undecidable (It is equivalent to the halting problem of Turing machines)



---

# Representing Tiling Problems



## Representing Tiling Problems

- Notice that every finite set of tiles  $\mathcal{T} = \{t_1, \dots, t_k\}$  can be represented as two binary relations  $H$ , and  $V$ .

## Representing Tiling Problems

- Notice that every finite set of tiles  $\mathcal{T} = \{t_1, \dots, t_k\}$  can be represented as two binary relations  $H$ , and  $V$ .


We put  $H(t_i, t_j)$  when the right side of  $t_i$  coincide with the left side of  $t_j$ , and similarly for  $V$ .

## Representing Tiling Problems

- ▶ Notice that every finite set of tiles  $\mathcal{T} = \{t_1, \dots, t_k\}$  can be represented as **two binary relations  $H$ , and  $V$** .

We put  $H(t_i, t_j)$  when the right side of  $t_i$  coincide with the left side of  $t_j$ , and similarly for  $V$ .

- ▶ For example, for



we will have

$$H = \{(t_1, t_3), (t_1, t_6), (t_2, t_4), (t_2, t_5), (t_2, t_1), \dots\}$$

$$V = \{(t_1, t_3), (t_1, t_5), (t_1, t_6), (t_2, t_3), (t_2, t_5), \dots\}$$

# Coding a Tiling of the Grid in FO

## Coding a Tiling of the Grid in FO

- **Theorem:** The problem of deciding whether a given FO formula is satisfiable is undecidable.

## Coding a Tiling of the Grid in FO

- **Theorem:** The problem of deciding whether a given FO formula is satisfiable is undecidable.

Let  $\rightarrow$  and  $\uparrow$  be relations, and  $t_1, \dots, t_n$  be propositional symbols.

Let  $\varphi$  be the conjunction of the formulas:

## Coding a Tiling of the Grid in FO

- **Theorem:** The problem of deciding whether a given FO formula is satisfiable is undecidable.

Let  $\rightarrow$  and  $\uparrow$  be relations, and  $t_1, \dots, t_n$  be propositional symbols.

Let  $\varphi$  be the conjunction of the formulas:

1) Total:  $[x]\langle y \rangle (x : \langle R \rangle y)$  for  $R \in \{\rightarrow, \uparrow\}$

## Coding a Tiling of the Grid in FO

- **Theorem:** The problem of deciding whether a given FO formula is satisfiable is undecidable.

Let  $\rightarrow$  and  $\uparrow$  be relations, and  $t_1, \dots, t_n$  be propositional symbols.

Let  $\varphi$  be the conjunction of the formulas:

- 1) **Total:**  $[x]\langle y \rangle (x:\langle R \rangle y)$  for  $R \in \{\rightarrow, \uparrow\}$
- 2) **Functional:**  $[x][y][z](x:\langle R \rangle y \wedge x:\langle R \rangle z \rightarrow y:z)$  for  $R \in \{\rightarrow, \uparrow\}$



## Coding a Tiling of the Grid in FO

- **Theorem:** The problem of deciding whether a given FO formula is satisfiable is undecidable.

Let  $\rightarrow$  and  $\uparrow$  be relations, and  $t_1, \dots, t_n$  be propositional symbols.

Let  $\varphi$  be the conjunction of the formulas:

- 1) **Total:**  $[x]\langle y \rangle (x:\langle R \rangle y)$  for  $R \in \{\rightarrow, \uparrow\}$
- 2) **Functional:**  $[x][y][z](x:\langle R \rangle y \wedge x:\langle R \rangle z \rightarrow y:z)$  for  $R \in \{\rightarrow, \uparrow\}$
- 3) **Commuting:**  $[x][y](x:\langle \rightarrow \rangle \langle \uparrow \rangle y \leftrightarrow x:\langle \uparrow \rangle \langle \rightarrow \rangle y)$

# Coding a Tiling of the Grid in FO

- **Theorem:** The problem of deciding whether a given FO formula is satisfiable is undecidable.

Let  $\rightarrow$  and  $\uparrow$  be relations, and  $t_1, \dots, t_n$  be propositional symbols.

Let  $\varphi$  be the conjunction of the formulas:

- 1) **Total:**  $[x]\langle y \rangle (x:\langle R \rangle y)$  for  $R \in \{\rightarrow, \uparrow\}$
- 2) **Functional:**  $[x][y][z](x:\langle R \rangle y \wedge x:\langle R \rangle z \rightarrow y:z)$  for  $R \in \{\rightarrow, \uparrow\}$
- 3) **Commuting:**  $[x][y](x:\langle \rightarrow \rangle \langle \uparrow \rangle y \leftrightarrow x:\langle \uparrow \rangle \langle \rightarrow \rangle y)$
- 4) **Tiled:**  $[x](x:t_1 \vee \dots \vee x:t_n),$

## Coding a Tiling of the Grid in FO

- **Theorem:** The problem of deciding whether a given FO formula is satisfiable is undecidable.

Let  $\rightarrow$  and  $\uparrow$  be relations, and  $t_1, \dots, t_n$  be propositional symbols.

Let  $\varphi$  be the conjunction of the formulas:

- 1) **Total:**  $[x]\langle y \rangle (x:\langle R \rangle y)$  for  $R \in \{\rightarrow, \uparrow\}$
- 2) **Functional:**  $[x][y][z](x:\langle R \rangle y \wedge x:\langle R \rangle z \rightarrow y:z)$  for  $R \in \{\rightarrow, \uparrow\}$
- 3) **Commuting:**  $[x][y](x:\langle \rightarrow \rangle \langle \uparrow \rangle y \leftrightarrow x:\langle \uparrow \rangle \langle \rightarrow \rangle y)$
- 4) **Tiled:**  $[x](x:t_1 \vee \dots \vee x:t_n),$
- 5) **But not Twice:**  $[x](x:t_i \rightarrow x:\neg t_j)$  for  $i \neq j$ ,

# Coding a Tiling of the Grid in FO

- **Theorem:** The problem of deciding whether a given FO formula is satisfiable is undecidable.

Let  $\rightarrow$  and  $\uparrow$  be relations, and  $t_1, \dots, t_n$  be propositional symbols.

Let  $\varphi$  be the conjunction of the formulas:

- 1) **Total:**  $[x]\langle y \rangle (x:\langle R \rangle y)$  for  $R \in \{\rightarrow, \uparrow\}$
- 2) **Functional:**  $[x][y][z](x:\langle R \rangle y \wedge x:\langle R \rangle z \rightarrow y:z)$  for  $R \in \{\rightarrow, \uparrow\}$
- 3) **Commuting:**  $[x][y](x:\langle \rightarrow \rangle \langle \uparrow \rangle y \leftrightarrow x:\langle \uparrow \rangle \langle \rightarrow \rangle y)$
- 4) **Tiled:**  $[x](x:t_1 \vee \dots \vee x:t_n),$
- 5) **But not Twice:**  $[x](x:t_i \rightarrow x:\neg t_j)$  for  $i \neq j$ ,
- 6) **Horizontal Match:**  $[x][y]((x:t_i \wedge x:\langle \rightarrow \rangle y) \rightarrow (\bigvee_{H(t_i, t_j)} y:t_j)),$

## Coding a Tiling of the Grid in FO

- **Theorem:** The problem of deciding whether a given FO formula is satisfiable is undecidable.

Let  $\rightarrow$  and  $\uparrow$  be relations, and  $t_1, \dots, t_n$  be propositional symbols.

Let  $\varphi$  be the conjunction of the formulas:

- 1) **Total:**  $[x]\langle y \rangle (x:\langle R \rangle y)$  for  $R \in \{\rightarrow, \uparrow\}$
- 2) **Functional:**  $[x][y][z](x:\langle R \rangle y \wedge x:\langle R \rangle z \rightarrow y:z)$  for  $R \in \{\rightarrow, \uparrow\}$
- 3) **Commuting:**  $[x][y](x:\langle \rightarrow \rangle \langle \uparrow \rangle y \leftrightarrow x:\langle \uparrow \rangle \langle \rightarrow \rangle y)$
- 4) **Tiled:**  $[x](x:t_1 \vee \dots \vee x:t_n),$
- 5) **But not Twice:**  $[x](x:t_i \rightarrow x:\neg t_j)$  for  $i \neq j$ ,
- 6) **Horizontal Match:**  $[x][y]((x:t_i \wedge x:\langle \rightarrow \rangle y) \rightarrow (\bigvee_{H(t_i, t_j)} y:t_j)),$
- 7) **Vertical Match:**  $[x][y]((x:t_i \wedge x:\langle \uparrow \rangle y) \rightarrow (\bigvee_{V(t_i, t_j)} y:t_j))$

# Coding a Tiling of the Grid in FO

- **Theorem:** The problem of deciding whether a given FO formula is satisfiable is undecidable.

Let  $\rightarrow$  and  $\uparrow$  be relations, and  $t_1, \dots, t_n$  be propositional symbols.

Let  $\varphi$  be the conjunction of the formulas:

- 1) **Total:**  $[x]\langle y \rangle (x:\langle R \rangle y)$  for  $R \in \{\rightarrow, \uparrow\}$
- 2) **Functional:**  $[x][y][z](x:\langle R \rangle y \wedge x:\langle R \rangle z \rightarrow y:z)$  for  $R \in \{\rightarrow, \uparrow\}$
- 3) **Commuting:**  $[x][y](x:\langle \rightarrow \rangle \langle \uparrow \rangle y \leftrightarrow x:\langle \uparrow \rangle \langle \rightarrow \rangle y)$
- 4) **Tiled:**  $[x](x:t_1 \vee \dots \vee x:t_n),$
- 5) **But not Twice:**  $[x](x:t_i \rightarrow x:\neg t_j)$  for  $i \neq j$ ,
- 6) **Horizontal Match:**  $[x][y]((x:t_i \wedge x:\langle \rightarrow \rangle y) \rightarrow (\bigvee_{H(t_i, t_j)} y:t_j)),$
- 7) **Vertical Match:**  $[x][y]((x:t_i \wedge x:\langle \uparrow \rangle y) \rightarrow (\bigvee_{V(t_i, t_j)} y:t_j))$

Then  $\varphi$  is satisfiable iff  $\mathcal{T}$  covers  $\mathbb{N} \times \mathbb{N}$ .

# Relevant Bibliography I

## Undecidable Problems

- ▶ Alonzo Church invented **lambda calculus** and propose it as **a model for computation**.



# Relevant Bibliography I

## Undecidable Problems

- ▶ Alonzo Church invented **lambda calculus** and propose it as **a model for computation**.
- ▶ He showed then that the problem of satisfiability of first-order logic could not be coded in the lambda calculus. Hence it was **uncomputable**.





# Relevant Bibliography I

## Undecidable Problems

- ▶ Alonzo Church invented **lambda calculus** and propose it as **a model for computation**.
- ▶ He showed then that the problem of satisfiability of first-order logic could not be coded in the lambda calculus. Hence it was **uncomputable**.
- ▶ Here is a list of some of Church's doctoral students: A. Anderson, P. Andrews, M. Davis, L. Henkin, S. Kleene, M. Rabin, B. Rosser, D. Scott, R. Smullyan, and A. Turing.



# Relevant Bibliography I

## Undecidable Problems

- ▶ Alonzo Church invented **lambda calculus** and propose it as **a model for computation**.
- ▶ He showed then that the problem of satisfiability of first-order logic could not be coded in the lambda calculus. Hence it was **uncomputable**.
- ▶ Here is a list of some of Church's doctoral students: A. Anderson, P. Andrews, M. Davis, L. Henkin, S. Kleene, M. Rabin, B. Rosser, D. Scott, R. Smullyan, and A. Turing.



Church, Alonzo (1956). *Introduction to Mathematical Logic*. The Princeton University Press.



# Relevant Bibliography II

## Undecidable Problems

- ▶ Alan Turing invented Turing Machines and Computer Science.



# Relevant Bibliography II

## Undecidable Problems

- ▶ Alan Turing invented **Turing Machines** and **Computer Science**.
- ▶ He showed that the halting problem could not be decided by a Turing Machine.



# Relevant Bibliography II

## Undecidable Problems

- ▶ Alan Turing invented **Turing Machines** and **Computer Science**.
- ▶ He showed that the halting problem could not be decided by a Turing Machine.
- ▶ And that the behavior of a Turing Machine can easily be described in first-order logic, providing an alternative proof that the satisfiability problem of first-order logic is undecidable.



# Relevant Bibliography II

## Undecidable Problems

- ▶ Alan Turing invented **Turing Machines** and **Computer Science**.
- ▶ He showed that the halting problem could not be decided by a Turing Machine.
- ▶ And that the behavior of a Turing Machine can easily be described in first-order logic, providing an alternative proof that the satisfiability problem of first-order logic is undecidable.



Turing, Alan (1936), *On Computable Numbers, with an Application to the Entscheidungsproblem*. Proceedings of the London Mathematical Society, Series 2, Vol.42, pp 230–265, [http://www.thocp.net/biographies/papers/turing\\_oncomputablenumbers\\_1936.pdf](http://www.thocp.net/biographies/papers/turing_oncomputablenumbers_1936.pdf)