Regression Autopsy

Eight Ways Your Model Is Lying to You

Lecture Notes

Statistical and Machine Learning Lab

Spring 2026

These notes accompany the interactive visualization series. All theorems stated here are validated by Monte Carlo simulation in the corresponding interactive pages.

Contents

1 Introduction **4**
1.1 The Exam Score Question . . . 4

2 The Data Generating Process **5**
2.1 What Is a Data Generating Process? . . . 5
2.2 What Is OLS? . . . 5
2.3 The Exam Score DGP . . . 6
2.4 Why These Relationships? . . . 8
2.5 Pathologies Embedded in the DGP . . . 9

3 Why Your Coefficient Is Wrong: Omitted Variable Bias **9**
3.1 Setup . . . 9
3.2 When OVB Cannot Be Signed . . . 12
3.3 The Proxy Variable Approach . . . 12
3.4 The Collider Trap: When "Controlling For" Backfires . . . 13

4 Why Your Uncertainty Is Wrong: Heteroscedasticity **14**
4.1 Confidence Intervals for Regression Coefficients . . . 14
4.2 The Classical Variance Formula . . . 14
4.3 Consequences for Inference . . . 16
4.4 The Breusch–Pagan Test . . . 17
4.5 Robust Standard Errors . . . 18

5 Why Your Significance Is Wrong: Effect Size and Power **18**
5.1 The Anatomy of a t-Statistic . . . 18
5.2 Statistical Power . . . 20
5.3 The Multiple Testing Problem . . . 20

6 Why Your Model Is Wrong: Specification Error **22**
6.1 The Problem of Functional Form . . . 22
6.2 Diminishing Returns to Studying . . . 22
6.3 The Extrapolation Danger . . . 23

7 Why Your R^2 Is Lying: Overfitting **24**
7.1 The Bias–Variance Decomposition . . . 24
7.2 Overfitting in Practice . . . 25
7.3 Cross-Validation . . . 25

8 Why Your Causal Claim Is Wrong: Endogeneity and Bad Controls **27**
8.1 Endogeneity . . . 27
8.2 The Causal Graph . . . 28
8.3 Collider Bias from Bad Controls . . . 28
8.4 Instrumental Variables . . . 30
8.5 The Strength of the Instrument . . . 31
8.6 A Numerical Example: Roommates and Exam Scores . . . 32

9 The Real World — When All Pathologies Strike at Once **33**
9.1 The Lalonde Dataset . . . 33
9.2 The Studying Analog . . . 34
9.3 Sensitivity Analysis: Cinelli–Hazlett Framework . . . 34

10 The Redemption — Regression Discontinuity Design **35**
10.1 When Design Earns Trust . . . 36
10.2 Example: Scholarship Cutoff . . . 36
10.3 Bandwidth Choice . . . 36

11 What "Controlling For" Actually Means — The Frisch–Waugh–Lovell Theorem **38**
11.1 Motivation . . . 38
11.2 The FWL Theorem . . . 38
11.3 Numerical Example . . . 39
11.4 What Happens with Bad Controls . . . 40

12 Diagnostic Toolkit Summary **40**

13 Theorem Map: How the Results Connect **41**

14 Conclusion **42**

A Proofs and Derivations **44**
A.1 Proof of the OVB Formula (Theorem 3.1) . . . 44
A.2 Proof of Collider Bias (Proposition 3.1) . . . 44
A.3 Derivation of the Sandwich Estimator (Theorem 4.1) . . . 44
A.4 Proof of the Bias–Variance Decomposition (Theorem 7.1) . . . 45
A.5 Proof of the Frisch–Waugh–Lovell Theorem (Theorem 11.1) . . . 45

How to Read These Notes

These notes are structured for two audiences:

- **First pass (the lecture):** Read the Toy Story boxes, the Key Insight boxes, and the Numerical Examples. Skip the theorems and proofs. This gives you the intuition and the punchlines.
- **Second pass (exam prep):** Read the theorem statements and the "Why is this true?" summaries. Work through the numerical examples with pen and paper. Consult the proofs in Section A only if you want the full derivation.

Each section follows the same rhythm: a *story* that makes the problem visceral, the *mathematics* that makes it precise, a *numerical example* you can verify on paper, and a *visualization* you can play with online. If you feel lost, jump to the next coloured box — the narrative will pick you back up.

Running gags to watch for: When you see "Is this a seat number?" — we are asking whether a statistically significant result is actually meaningful. When you see a "ghost" — we mean an omitted variable whose effect is hiding inside another coefficient. When you see a *coefficient tracker* note at the start of a section, it summarises how the studying coefficient has changed so far—each change corresponds to a different failure mode. And the A-level algorithm from the Introduction will return in the Conclusion, once you have the tools to diagnose exactly what went wrong.

1 Introduction

Cold Open

A study of 50,000 university students found that sitting in a higher-numbered seat in the exam hall is associated with higher exam scores ($p = 0.002$). Should the university assign seats by lottery? Before you answer—read on.

The Algorithm That Ruined 72 Hours

In August 2020, the UK exam regulator Ofqual deployed a regression model to assign A-level grades to half a million students whose exams had been cancelled due to COVID-19. The model included school-level variables, prior performance, and teacher predictions. Its R^2 was good. Its diagnostics passed. On Tuesday, students received their grades. By Thursday, the algorithm had been scrapped.

The model systematically downgraded students from disadvantaged schools. A student with identical ability and identical teacher predictions received a lower grade if their school had historically underperformed—a textbook case of conditioning on a collider (school reputation is caused by both student quality *and* institutional resources). The algorithm did not measure student ability; it measured school postcode.

This course is about why. Every failure mode we study—omitted variable bias, bad controls, specification error, fragile diagnostics—was present in the Ofqual model. We will return to this example in the conclusion, once you have the vocabulary to say exactly what went wrong.

Regression is the most widely used tool in quantitative data analysis. It produces point estimates, standard errors, confidence intervals, p-values, and R^2 statistics—all of which look authoritative. In most observational analyses, at least one of these numbers is wrong.

These lecture notes formalise the mathematical foundations behind the eight core failure modes of ordinary least squares (OLS) regression. Each section corresponds to one interactive notebook in the accompanying lab series. The notes are designed to be self-contained: every claim is stated as a formal result and, in the lab, confirmed by Monte Carlo simulation.

Each section follows a consistent pattern: we begin with an everyday story that makes the problem visceral, then state the rigorous mathematics, then work through a tiny numerical example you can verify on paper, and finally describe an interactive visualization that lets you play with the result.

1.1 The Exam Score Question

Your friend claims studying 10 hours guarantees an A. You have data from 500 students in an introductory statistics course. Does studying more actually help, and by how much? This is the question that will drive every regression in this course.

The dataset contains the following variables:

- Y**: Exam score** (0–100). This is the *outcome*—the thing we are trying to explain. Every regression in these notes has Y on the left-hand side: we want to understand what drives exam performance. In the language of causal inference, Y is the dependent variable.

- X_1**: Hours studied** in the week before the exam. This is the *treatment*—the variable whose effect we care about most. The central question is whether studying more *causes* higher scores, and if so, by how many points per hour. X_1 is what you would manipulate in

an ideal experiment: randomly assign students to study 2 hours or 6 hours, then compare their scores.

- X_2**: Number of lectures attended** (out of 14). This is a *control* variable. Students who attend more lectures tend to score higher, but we want to separate the lecture effect from the studying effect. Including X_2 in the regression prevents lecture attendance from contaminating our estimate of the studying effect.

- X_3**: Self-assessed performance** (1–10, surveyed right after handing in the exam). This is the *trap*. It looks like useful information—surely students who felt good about the exam scored higher? But that feeling is *caused* by how well they actually did: students who studied hard, attended lectures, and are disciplined walk out of the exam room knowing it went well. X_3 is a *post-treatment* variable: it sits downstream of Y in the causal chain. Including it as a control will backfire spectacularly (see Section 8).

- U**: Self-discipline score** (unobserved by the analyst). Before the semester started, every student dragged an Instagram story slider in response to the question "How disciplined are you with your studies?"—the kind of quick poll you have seen a hundred times. The slider produces a continuous value between 0 ("I have zero discipline") and 1 ("I follow a strict study schedule"). We standardise these responses to have mean 0 and standard deviation 1, giving $U \sim \mathcal{N}(0, 1)$.

  The crucial point: *you, the analyst, do not have access to this data.* It lives on Instagram's servers, not in the university's registrar system. But U drives both how much students study *and* how well they perform on the exam (through paying attention in class, doing practice problems, staying focused during the test). Because U affects both the treatment X_1 and the outcome Y, it is a *confounder*. This is what makes the analysis tricky: without observing U, OLS will attribute some of self-discipline's effect to studying, inflating the coefficient.

Using a single recurring dataset creates continuity: when a coefficient changes from one section to the next, you'll remember the old value.

2 The Data Generating Process

2.1 What Is a Data Generating Process?

A data generating process (DGP) is a set of equations that describes how nature produces data. Think of it as a recipe: if you follow these steps, you get a dataset. We write the recipe down because it lets us ask precise questions: if *this* is how the world works, what will OLS tell us?

In real life, we don't know the DGP—that's the whole problem. If we knew exactly how exam scores are produced, we wouldn't need statistics. But by working with a known DGP in a classroom, we can check whether OLS gets the right answer. When it doesn't, we learn exactly why. Every gap between truth and estimate can be traced back to a specific feature of the DGP that OLS failed to handle.

2.2 What Is OLS?

Given a dataset of n observations with an outcome vector $\mathbf{y}$ and a design matrix $\mathbf{X}$ (whose columns are the variables we include in the regression), the **ordinary least squares (OLS)** estimator is

$$\hat{\beta} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}. \tag{1}$$

OLS finds the coefficients that minimise the sum of squared residuals—the vertical distances between the data points and the fitted line.

Here is the single most important idea in these notes: $\hat{\beta}$ **is a random variable.** If you collected a different sample of 500 students, you'd get a different $\hat{\beta}$. The true β is a fixed number—it is a property of the DGP. The estimator $\hat{\beta}$ is a random variable—it depends on which students happened to be in your sample.

This distinction is the foundation of everything that follows. When we say "$\hat{\beta}$ is biased," we mean that if you repeated the experiment thousands of times, the *average* of all the $\hat{\beta}$'s would not equal the true β. When we say "the standard error is wrong," we mean the formula that predicts how much $\hat{\beta}$ varies across samples is giving the wrong answer.

Remark 2.1 (Probability limit). Throughout these notes, plim denotes the *probability limit*—the value an estimator converges to as the sample size grows to infinity. Think of it as "what the estimator settles down to with unlimited data." When we write $\text{plim}\ \hat{\beta}_1 = \beta_1$, we mean that the estimator is *consistent*: with enough data, it gets arbitrarily close to the truth. When we write $\text{plim}\ \hat{\beta}_1 = \beta_1 + \text{something}$, that "something" is the asymptotic bias—it never goes away, no matter how much data you collect.

2.3 The Exam Score DGP

Definition 2.1 (Exam Score DGP)**.** The data are generated according to the following structural equations:

$$U \sim \mathcal{N}(0, 1), \tag{2}$$
$$X_1 = 2 + 0.5\,U + \eta_1, \quad \eta_1 \sim \mathcal{N}(0,\ 0.3), \tag{3}$$
$$X_2 = 10 + 0.3\,U + \eta_2, \quad \eta_2 \sim \mathcal{N}(0,\ 1), \tag{4}$$
$$\varepsilon \sim \mathcal{N}(0,\ 0.5 \cdot |X_1|)\,, \tag{5}$$
$$Y = 50 + 8\log(X_1) + 3\,X_2 + 2\,U + \varepsilon, \tag{6}$$
$$X_3 = 0.9\,Y + \eta_3, \quad \eta_3 \sim \mathcal{N}(0,\ 2). \tag{7}$$

If these structural equations look intimidating, don't worry. Let's walk through each one in plain language.

$U \sim \mathcal{N}(0, 1)$**: Self-discipline.** Before the semester, every student dragged a slider on an Instagram story poll: "How disciplined are you with your studies?" The slider returns a continuous value from 0 to 1. We standardise the responses to mean 0 and standard deviation 1, giving $U \sim \mathcal{N}(0, 1)$. Most students are average; a few are very disciplined or very undisciplined. Critically, *you do not have this data*—it lives on Instagram's servers, not in the university's system. This is the hidden variable that makes everything complicated.

$X_1 = 2 + 0.5\,U + \eta_1$**: Hours studied.** More disciplined students study more. A student of average self-discipline ($U = 0$) studies about 2 hours. Each standard deviation of extra discipline adds half an hour. The noise η_1 captures everything else that affects study time—a busy week, a competing deadline, pure randomness.

Remark 2.2 (Negative study hours). Because $X_1 \sim \mathcal{N}(2, 0.25+0.3) = \mathcal{N}(2, 0.55)$, approximately 0.3% of simulated students will have $X_1 < 0$ (negative study hours). In practice, we would truncate X_1 at zero. For mathematical convenience—so that X_1 remains exactly normally distributed and all our closed-form results go through cleanly—we keep the unbounded normal. If you are running simulations and encounter a negative value, this is why.

$X_2 = 10 + 0.3\,U + \eta_2$**: Lectures attended.** More disciplined students attend more lectures. Average attendance is 10 out of 14. Self-discipline pushes this up slightly—an extra standard deviation adds 0.3 lectures. The noise η_2 is larger here ($\sigma = 1$) because lecture attendance depends heavily on scheduling conflicts, illness, and other factors unrelated to discipline.

$\varepsilon \sim \mathcal{N}(0,\, 0.5 \cdot |X_1|)$**: Exam noise.** The randomness in exam scores depends on how much you studied (where $0.5 \cdot |X_1|$ is the *variance*, not the standard deviation; thus $\mathrm{SD}(\varepsilon) = \sqrt{0.5 \cdot |X_1|}$). Students who barely studied have tightly clustered (low) scores—there is not much variation in how badly you can do when you are unprepared. Students who studied a lot have more spread—because at high study hours, *how* you study matters more than *how long*, and that quality varies widely. This is *heteroscedasticity*: the variance of the error term changes with a predictor.

$Y = 50 + 8\log(X_1) + 3\,X_2 + 2\,U + \varepsilon$**: Exam score.** The exam score. The base score is 50—what an average student scores with minimal preparation. The first hour of studying helps enormously (the logarithm rises steeply near zero), but returns diminish: going from 1 to 2 hours helps far more than going from 9 to 10 hours. Attending one more lecture adds 3 points. Being one standard deviation more disciplined adds 2 points directly—through paying attention in class, doing practice problems, staying focused during the test—things that studying hours alone do not capture.

$X_3 = 0.9\,Y + \eta_3$**: Self-assessed performance.** Right after handing in the exam, students rate how well they think they did on a scale from 1 to 10. This self-assessment mostly reflects the actual exam score—students who studied hard, attended lectures, and are disciplined walk out knowing it went well. The coefficient 0.9 means self-assessed performance tracks the actual score almost one-to-one. The noise η_3 captures variation in self-awareness (some students are poor judges of their own performance). Because X_3 is measured *after* the exam—and is driven by the same factors that produced Y—including it as a control will backfire.

The Numbers Behind the Story

Here are the default parameter values, tied to what they mean in the real world:
The base score ($\beta_0 = 50$): a completely average student who barely studied and attended some lectures scores about 50 out of 100. This is the exam's difficulty anchor.
The studying payoff ($\beta_1 = 8$): each unit increase in log(hours studied) adds 8 points. Concretely: going from 1 to 2.7 hours (one log-unit) gains you 8 points. The logarithm means the first hour is worth far more than the tenth.
The lecture dividend ($\beta_2 = 3$): each additional lecture attended adds 3 points. A student who attended all 14 lectures has a 42-point boost from lectures alone.
The self-discipline bonus ($\beta_U = 2$): each standard deviation of self-discipline adds 2 points directly—through focus, practice, and persistence—even holding study hours and lectures fixed. This is the hidden advantage that creates the bias.
The confounding links ($\gamma_1 = 0.5$, $\gamma_2 = 0.3$): self-discipline pushes study hours up by 0.5 per SD and lectures by 0.3 per SD. These links are why omitting U biases the studying coefficient.
The noise that grows ($h = 0.5$): exam score variance scales with study hours. Low-study students cluster tightly around low scores. High-study students spread out—how you study matters more than how long, and that varies.
The post-treatment trap ($\alpha = 0.9$): self-assessed performance tracks the actual score almost one-to-one. Including it as a control is like partially conditioning on Y itself.

Remark 2.3 (Causal ordering). The causal graph below shows every relationship in the DGP. Solid arrows indicate observed relationships; the dashed node (U) is unobserved by the analyst. The red node (X_3) is post-treatment. Generate data in causal order: U first, then (X_1, X_2), then Y, then X_3.



2.4 Why These Relationships?

Key Insight

The structural equations above are not arbitrary—each one encodes a specific real-world mechanism. Refer to the causal graph above—each bullet below corresponds to one arrow in the diagram.

- **Self-discipline → studying (positive):** Obvious—disciplined students put in more hours. This is why U appears with a positive coefficient in the equation for X_1.
- **Self-discipline → lectures (positive):** Disciplined students show up more. Same logic, slightly weaker effect (0.3 vs. 0.5) because attendance depends more on logistics than on willpower.
- **Studying → score (logarithmic):** The first hour of studying recovers the low-hanging fruit—basic definitions, key formulas, the structure of the exam. The 10th hour yields much less—you are reviewing material you already know. Diminishing returns are built into the log.
- **Lectures → score (linear):** Attending lectures exposes you to material, worked examples, and the instructor's emphasis on what matters for the exam. Each additional lecture adds 3 points ($\beta_2 = 3$). This is a direct causal effect, not mediated by studying—you learn things in lecture that you wouldn't learn from the textbook alone.
- **Self-discipline → score (direct):** Self-discipline helps *beyond* studying. Disciplined students pay attention in class, ask questions, do practice problems, and stay focused during the exam. These things boost scores even holding study hours fixed. This direct effect ($\beta_U = 2$) is what creates the omitted variable bias problem.

- **Noise $\varepsilon \to$ score:** Even after accounting for studying, lectures, and self-discipline, there is randomness: a bad night's sleep, a lucky guess, a misread question.
- **Score $\to$ self-assessment (positive):** Students who did well on the exam know it when they walk out. This is *not* a causal effect of self-assessment on scores—it is the reverse. Feeling good about the exam does not cause a high score; doing well causes both the high score and the good feeling. Confusing the direction is exactly the mistake that leads to collider bias.

2.5 Pathologies Embedded in the DGP

The single DGP simultaneously contains all of the following pathologies:

1. **Omitted variable bias:** U (self-discipline) drives both X_1 (hours studied) and Y (exam score). Omitting U biases $\hat{\beta}_1$ upward—OLS attributes some of self-discipline's effect to studying.
2. **Heteroscedasticity:** $\mathrm{Var}(\varepsilon \mid X_1) = h \cdot |X_1|$; students who study more have more variable scores. OLS standard errors, which assume constant variance, will be wrong.
3. **Inflated significance:** with large n (50,000 student-course observations university-wide), trivially small effects become "significant." A coefficient of 0.01 points per hour can have $p < 0.001$.
4. **Misspecification:** the true relationship is $\log(X_1)$, not X_1. Fitting a linear model to a logarithmic truth introduces systematic errors that no amount of data can fix.
5. **Overfitting:** adding polynomial terms (study hours squared, cubed, etc.) inflates training R^2 but destroys test R^2. The model memorises noise instead of learning the signal.
6. **Collider bias / bad control:** X_3 (self-assessed performance) is caused by Y; conditioning on it opens a spurious path between X_1 and Y, distorting the studying coefficient.

Each notebook isolates and examines one pathology while the others remain present in the background.

3 Why Your Coefficient Is Wrong: Omitted Variable Bias

3.1 Setup

Imagine you are a teaching assistant analysing exam data. You have each student's exam score and the number of hours they studied—but you do *not* have their lecture attendance records (those are kept by a different department and you forgot to request them). So you run the only regression you can: score on hours studied, omitting lectures. This is not laziness—it is a common real-world constraint. Data lives in silos, access requires paperwork, and deadlines don't wait. The question is: what happens to your coefficient on study hours when you are *forced* to leave out a relevant variable?

The CNN Coffee Miracle

In 2017, CNN ran the headline: "Coffee drinkers live longer, study finds." The study followed hundreds of thousands of people and found that moderate coffee drinkers had a 16% lower risk of death. The p-value was tiny. The sample size was enormous. Should you start drinking three cups a day?

Here is the problem: people who drink moderate morning coffee also tend to exercise regularly, eat reasonably well, not smoke, and hold stable jobs with health insurance. The hidden variable U is *healthy lifestyle*—and it drives both coffee consumption and longevity. The coffee coefficient absorbed the ghost of kale salads and morning jogs.

This is exactly what the OVB formula captures: the bias on coffee's coefficient equals the true effect of healthy lifestyle on longevity times the correlation between healthy lifestyle and coffee consumption. Both are positive, so the bias is positive—coffee looks more life-extending than it really is. Leave out a variable that matters and is correlated with what you included, and its ghost haunts your estimate.

The Coefficient Auction

Before I show you any regression output: shout your guess. What is the effect of one additional hour of studying on exam scores? Write down a number.
If you guessed something between 5 and 10 points per hour, you are in good company—that is where most people's intuitions land. Hold on to that number. We are about to watch it change, and each change will teach you something about what OLS is actually doing.

Suppose the true model is

$$Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \varepsilon, \qquad \mathbb{E}[\varepsilon \mid X_1, X_2] = 0, \tag{8}$$

but the analyst estimates the *short* regression

$$Y = \tilde{\beta}_0 + \tilde{\beta}_1 X_1 + \tilde{\varepsilon}. \tag{9}$$

Why would you include X_2 in the first place, if you only care about the effect of X_1 on Y? The answer is that X_2 acts as a *control variable*.

Definition 3.1 (Control Variable)**.** A *control variable* is a variable we include in the regression not because we care about its coefficient, but because including it prevents its effect from contaminating the coefficient we *do* care about. If X_2 affects Y and is correlated with X_1, omitting X_2 lets its effect leak into $\hat{\beta}_1$.

In the studying scenario, suppose we want to know how hours studied (X_1) affect exam scores (Y). Hours of sleep (X_2) also affect exam scores, and students who study more tend to sleep less. If we leave sleep out of the regression, some of sleep's positive effect on scores gets tangled up with studying's effect—and because studying and sleep are *negatively* correlated, the result is that OLS *underestimates* the benefit of studying. Including sleep as a control variable disentangles the two effects.

Theorem 3.1 (Omitted Variable Bias Formula)**.** *Let $\hat{\beta}_1^{short}$ denote the OLS estimator of $\tilde{\beta}_1$ from the short regression (9), and let β_1 be the true coefficient in (8). Define the* auxiliary slope

$$\delta_1 \;=\; \frac{\mathrm{Cov}(X_1, X_2)}{\mathrm{Var}(X_1)}.$$

Then

$$\mathbb{E}[\hat{\beta}_1^{short}] \;=\; \beta_1 + \beta_2\,\delta_1. \tag{10}$$

The term $\beta_2\delta_1$ is the ***omitted variable bias****.*

Why is this true? Because when you regress Y on X_1 alone, OLS gives X_1 credit for *all* the variation in Y that is correlated with X_1—including the part that actually belongs to X_2 but travels through the X_1–X_2 correlation. The bias term $\beta_2\delta_1$ measures exactly how much of X_2's effect gets falsely attributed to X_1. The formal derivation is in Appendix A.

Studying, Sleeping, and Exam Scores

True DGP: $Y = 60 + 5X_1 + 3X_2$ (no noise, for clarity), where X_1 is hours studied and X_2 is hours slept the night before the exam.
Here are five students whose data follow this DGP exactly:

Student	X_1 (studied)	X_2 (slept)	$Y = 60 + 5X_1 + 3X_2$
A	1	8	$60 + 5(1) + 3(8) = 89$
B	2	6	$60 + 5(2) + 3(6) = 88$
C	3	7	$60 + 5(3) + 3(7) = 96$
D	4	4	$60 + 5(4) + 3(4) = 92$
E	5	5	$60 + 5(5) + 3(5) = 100$

Notice the pattern: students who study more tend to sleep less (the correlation between X_1 and X_2 is negative).
Step 1 — Full regression (include both X_1 and X_2): recovers $\hat{\beta}_1 = 5$, $\hat{\beta}_2 = 3$ exactly, because the data were generated with no noise.
Step 2 — Short regression (omit sleep): regressing Y on X_1 alone gives $\hat{\beta}_1^{\text{short}} = 2.6$. The coefficient on studying has *shrunk* from 5 to 2.6. Where did the missing 2.4 go?
Step 3 — Auxiliary regression (X_2 on X_1): regressing hours slept on hours studied gives slope $\delta_1 = -0.8$. This makes sense: each extra hour of studying costs about 0.8 hours of sleep.
Step 4 — Verify the OVB formula:

$$\hat{\beta}_1^{\text{short}} = \beta_1 + \beta_2 \cdot \delta_1 = 5 + 3 \times (-0.8) = 5 - 2.4 = 2.6. \quad \checkmark$$

The interpretation: omitting sleep shrank the study coefficient from 5 to 2.6. Because studying more costs sleep ($\delta_1 = -0.8$) and sleep helps scores ($\beta_2 = 3$), some of studying's real benefit gets "charged" to the missing sleep variable, pulling the estimate downward.

"The Ghost in the Regression" — Interactive OVB Explorer

Try it: A scatterplot of hours studied (X_1) vs. exam score (Y) shows two regression lines—the full model (gold) and the short model (red). Toggle a switch to include/exclude hours slept (X_2) and watch the slope shift in real time.
Key interactions:

- *Correlation slider* (-0.95 to $+0.95$): controls how strongly studying and sleep are related. At zero correlation, both lines coincide—omitting an uncorrelated variable causes no bias.
- *Effect-size slider* (β_2 from -5 to $+10$): when the omitted variable doesn't matter ($\beta_2 = 0$), there's no bias regardless of correlation.

The formula panel updates live, showing how $\beta_2 \cdot \delta_1$ produces the exact gap you see.

The ghost visualization. Imagine the same scatterplot, but now every dot is coloured by the hidden variable U (self-discipline): blue for low-discipline students, red for high-discipline students. With U invisible (all dots gray), OLS fits the steep red line—it cannot see that high-discipline students cluster in the upper-right and low-discipline students in the lower-left. The colour gradient *is* the bias: it pulls the line steeper than it should be. Now reveal U: the colour pattern becomes visible, a second gold "truth" line appears with a shallower slope, and the gap between the red and gold lines is labelled $\beta_2 \cdot \delta_1$. A translucent ghost figure hovers behind the data cloud, one hand pushing dots rightward

(more studying), the other pushing dots upward (higher scores). The ghost's hands are the confounding links γ_1 and β_U, and they tilt the regression line away from the truth.

Before and After: The OVB Reveal

Slide 1 (the naive view): A clean scatterplot of study hours vs. exam score with a steep positive regression line. Title: "Each additional hour of studying adds 10.5 points to your exam score!" Green checkmark. This is what OLS tells you when U is hidden.

Slide 2 (the corrected view): The same scatterplot, but now the ghost of self-discipline is visible behind the data cloud. The regression line has split in two: the red naive line (steep) and the gold true line (less steep, ≈5.5 points per hour). Title: "Self-discipline was doing half the work—the true studying effect is only 5.5 points." Red X replaces the checkmark. Between the two lines, the gap is labelled "OVB $= \beta_2 \cdot \delta_1$." The lesson: what looked like a studying effect was partly a discipline effect in disguise.

Key Insight

The bias has a *sign* and a *direction*. In the studying DGP, recall that both hours studied (X_1) and lectures attended (X_2) are positively driven by self-discipline (U), so $\delta_1 > 0$, and $\beta_2 = 3 > 0$. The bias is therefore *positive*: the short regression *overestimates* the effect of hours studied, because it gives X_1 credit for the beneficial effect of lecture attendance that travels through the shared self-discipline confounder.

Try It Yourself: Sign the Bias

A study finds that dog owners live longer than non-owners. What is the likely omitted variable? What is the sign of the bias?
Hint: Wealthy, active, healthy people are more likely to own dogs. Wealth and activity also improve longevity. So $\beta_{\text{health}} > 0$ (health helps longevity) and $\delta > 0$ (healthier people are more likely to own dogs). The OVB formula gives a *positive* bias: dog ownership looks more life-extending than it really is, because it absorbs the effect of the healthy-and-wealthy lifestyle that correlates with pet ownership.

3.2 When OVB Cannot Be Signed

The formula (10) generalises to multiple omitted variables. If k variables are omitted, the bias on $\hat{\beta}_1$ is a weighted sum of terms $\beta_j \delta_{1j}$, where each δ_{1j} is the partial regression coefficient of X_j on X_1 after partialling out included regressors. In general, the sign of the total bias is indeterminate without knowledge of all omitted variables.

3.3 The Proxy Variable Approach

The Repair

In the studying scenario, the true confounder is self-discipline (U)—the Instagram slider score that lives on a social media server, not in the university's records. But suppose we can measure something related, like attendance at optional review sessions. Call this proxy $\tilde{U}$. If $\tilde{U}$ explains a fraction $q \in [0, 1]$ of the variation in U, the residual bias is approximately $(1 - q)$ times the original bias. At $q = 1$ (the proxy perfectly captures self-discipline), the bias vanishes entirely. At $q = 0$ (the proxy is useless noise), the full

bias remains. In practice, even an imperfect proxy—$q = 0.5$, say—cuts the bias in half, which is a substantial improvement over ignoring the confounder entirely.

3.4 The Collider Trap: When "Controlling For" Backfires

Not all variables should be controlled for. Some variables, when included in the regression, make things *worse* rather than better. To understand when this happens, we need the concept of a collider.

Definition 3.2 (Path and Collider)**.** In a causal graph, a *path* between two variables is any sequence of arrows connecting them, regardless of the arrows' directions. For example, in the exam score graph, one path from X_1 to X_3 is $X_1 \to Y \to X_3$.

A variable C is a *collider* on a path if two arrows point *into* it on that path: $A \to C \leftarrow B$. In our graph, Y is a collider on the path $X_1 \to Y \leftarrow U$, because both X_1 and U point into Y.

The key rule: conditioning on a collider (including it in the regression) *opens* a spurious association between the variables on either side. Conditioning on a non-collider *blocks* the association. This is why including X_3 as a control backfires: X_3 is a descendant of the collider Y, and conditioning on a descendant partially conditions on the collider itself.

Proposition 3.1 (Collider Bias)**.** *Let X and U be independent, and let $C = \alpha_1 X + \alpha_2 U + \eta$ with $\alpha_1, \alpha_2 \neq 0$. Then X and U are unconditionally independent, but conditionally on C, they become dependent:*

$$\mathrm{Cov}(X, U \mid C) \neq 0.$$

Why is this true? Because fixing C constrains the sum $\alpha_1 X + \alpha_2 U$, so knowing that X is large forces U to be small (and vice versa)—the collider manufactures a negative association from nothing. The formal derivation is in Appendix A.

A *collider path* is any path between two variables that passes through a collider. Such a path is "blocked" by default (no spurious association flows along it) but becomes "open" if you condition on the collider. This is the reverse of the usual rule: for non-colliders, conditioning blocks a path; for colliders, conditioning opens one.

Warning

In our studying DGP, the variable X_3 (self-assessed performance after the exam) is defined as $X_3 = 0.9Y + \eta_3$. Self-assessment is driven by the actual exam score Y, which in turn is driven by hours studied (X_1), lectures attended (X_2), and self-discipline (U). Including X_3 as a control variable makes things *worse*, not better.

Why? Confidence is a *post-treatment* variable—it is caused by the outcome Y, not the other way around. Conditioning on confidence is like asking: "Among students who are equally confident, does studying more help?" But think about what "equally confident" means. A student who studied very little but is highly confident must have some other advantage—perhaps natural talent or prior knowledge. A student who studied a great deal but is only moderately confident must have some disadvantage that studying couldn't fully overcome. By holding confidence fixed, you have manufactured a fake negative relationship between studying and these unobserved traits. You have opened a collider path $X_1 \to Y \to X_3 \leftarrow \eta_3$, and the resulting bias can be *larger* than the original OVB you were trying to fix.

Rule of thumb: never control for a variable that is caused by the treatment or the outcome. If it comes *after* the treatment in the causal chain, leave it out.

4 Why Your Uncertainty Is Wrong: Heteroscedasticity

Coefficient tracker: the studying coefficient was 10.5 in Section 3 (biased by the ghost of self-discipline). Now we keep that inflated number—but discover the uncertainty around it is also a lie.

4.1 Confidence Intervals for Regression Coefficients

Before we can understand what goes wrong with uncertainty estimates, we need to be precise about what "uncertainty" means in a regression context. The central object is the *confidence interval*.

Definition 4.1 (Confidence Interval)**.** A **95% confidence interval** for a parameter β_j is a random interval $[L, U]$, computed from the data, such that if we repeated the entire experiment many times—drawing new data each time and computing a new interval—then 95% of those intervals would contain the true value β_j.

It is crucial to understand what this does *not* mean. A 95% CI does not say "there is a 95% probability that β_j lies in this interval." The true β_j is a fixed number; it is either inside the interval or it is not. The 95% refers to the *procedure*: if you use this recipe over and over, it works 95% of the time.

For the OLS estimator, the standard confidence interval is

$$\hat{\beta}_j \pm z_{\alpha/2} \cdot \mathrm{SE}(\hat{\beta}_j), \tag{11}$$

where $z_{\alpha/2}$ is the critical value from the standard normal distribution (for a 95% CI, $z_{0.025} = 1.96$) and $\mathrm{SE}(\hat{\beta}_j)$ is the *standard error*—an estimate of the standard deviation of $\hat{\beta}_j$ across hypothetical repeated samples.

Definition 4.2 (Coverage)**.** The **coverage** of a confidence interval procedure is the fraction of times the interval actually contains the true parameter value across many repetitions. A correctly calibrated 95% CI should have 95% coverage. If it does not, the standard error formula is wrong.

This is the key question for this entire section: *what happens when the SE formula is wrong?* The confidence intervals look fine—they still produce a number ± another number—but their actual coverage is not 95%. They are lying about how uncertain you should be.

4.2 The Classical Variance Formula

Drunk Darts

You and your friends play darts at a pub. Sober, everyone throws at the bullseye and the darts land in a tight cluster—maybe 2 centimetres from the centre on average. After one beer, the cluster loosens a bit. After three beers, darts are hitting the wall, the floor, and occasionally the bartender. Your *aim* is still the bullseye—on average, you're not biased in any direction—but the *spread* of your throws grows with every drink.

Here is the connection to regression: beers are the predictor (X), and dart accuracy is the outcome (Y). The average effect of beer on accuracy is estimable (it's negative—more beer, worse aim). But the *variance* of the outcome depends on X: sober throws are predictable, drunk throws are chaotic. If you use the sober-level spread to judge your uncertainty about the drunk-level effect, you will be wildly overconfident. Your confidence interval will say "I'm 95% sure the effect is between -2 and -1 cm per beer" when the true range should be "-8 to $+4$ cm per beer."

This is heteroscedasticity: the noise depends on the predictor. And if you ignore it, your

uncertainty estimates are a lie.

Under the standard linear model $Y = X\beta + \varepsilon$ with $\mathbb{E}[\varepsilon\varepsilon^\top \mid X] = \sigma^2 I_n$, the variance of the OLS estimator is

$$\mathrm{Var}(\hat{\beta}_{\mathrm{OLS}} \mid X) = \sigma^2 (X^\top X)^{-1}. \tag{12}$$

Definition 4.3 (Heteroscedasticity)**.** The error term is *heteroscedastic* if $\mathrm{Var}(\varepsilon_i \mid X_i) = \sigma_i^2$ with σ_i^2 not constant across i. In the studying DGP, $\varepsilon \sim N(0,\ 0.5 \cdot |X_1|)$: the more a student studies, the more unpredictable their exam score becomes. A student who studies 1 hour has noise standard deviation $\sqrt{0.5 \cdot 1} \approx 0.71$, while a student who studies 6 hours has noise standard deviation $\sqrt{0.5 \cdot 6} \approx 1.73$—more than twice as large.

Remark 4.1 (Heteroscedasticity in everyday life)*.* Consider apartment rents. A 20m^2 studio in Zurich rents for CHF 800–1,000 (a narrow range—there is not much a landlord can do to differentiate a tiny studio). A 200m^2 flat rents for CHF 3,000–8,000—the range is enormous because large apartments vary in quality, location, and luxury features. The spread of rents "fans out" with apartment size. That fanning is heteroscedasticity.

Heteroscedasticity in Everyday Wisdom

There is a folk version of heteroscedasticity that everyone already knows:
"If you never buy a lottery ticket, you'll never win—but you'll never lose either."
"If you don't put yourself out there, nothing bad happens—but nothing good happens either."
The common thread: low exposure means low variance (predictable but boring). High exposure means high variance (exciting but risky). In regression terms: the predictor X determines not just the expected outcome, but how *spread out* the outcomes are. A student who studies 1 hour will reliably score around 55. A student who studies 10 hours might score 65 or 90—because at high effort, the quality of studying matters enormously, and that quality varies.

Theorem 4.1 (Variance Under Heteroscedasticity — The Sandwich Formula)**.** *When* $\mathbb{E}[\varepsilon\varepsilon^\top \mid X] = \Omega \neq \sigma^2 I_n$*, the OLS estimator* $\hat{\beta}_{OLS}$ *remains* unbiased *but its variance becomes*

$$\mathrm{Var}(\hat{\beta}_{OLS} \mid X) = (X^\top X)^{-1} X^\top \Omega\, X\, (X^\top X)^{-1}. \tag{13}$$

This is the ***sandwich formula****: the "bread" is* $(X^\top X)^{-1}$ *on each side, and the "meat" is* $X^\top \Omega\, X$ *in the middle. The classical formula* (12) *underestimates or overestimates the true variance, depending on the correlation structure between* X *and the error variances.*

Remark 4.2 (Ω is diagonal, not correlated)*.* A common confusion: heteroscedasticity does *not* mean the errors of different observations are correlated. Student A's error and Student B's error are still independent. What changes is that the *size* of each error varies:

$$\Omega = \begin{pmatrix} \sigma_1^2 & 0 & \cdots & 0 \\ 0 & \sigma_2^2 & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & \sigma_n^2 \end{pmatrix}.$$

The off-diagonal zeros mean: knowing Student A's error tells you nothing about Student B's. But $\sigma_1^2 \neq \sigma_2^2$—some students have noisier outcomes than others. In the dart analogy: knowing how far *your* dart landed from the bullseye tells me nothing about where *my* dart landed. But if you've had five beers and I've had none, your dart's spread is much larger than mine.

When errors *are* correlated across observations (e.g., students in the same study group share shocks), Ω has nonzero off-diagonals. That is a different problem called *autocorrelation* or *clustering*, which we do not address in these notes.

Five Students, One Exam

We predict exam scores (Y) from hours studied (X) for five students. The true relationship is $Y = 50 + 8X$ with heteroscedastic noise that grows with X:

Student	X (hours)	Y (score)	$\hat{Y} = 49.6 + 8.0X$	Residual $\hat{\varepsilon}$	$\hat{\varepsilon}^2$
A	1	59	57.6	+1.4	1.96
B	2	66	65.6	+0.4	0.16
C	3	75	73.6	+1.4	1.96
D	4	72	81.6	−9.6	92.16
E	5	96	89.6	+6.4	40.96

Verification: $\bar{X} = 3$, $\bar{Y} = 73.6$, $\hat{\beta}_1 = \sum(X_i - \bar{X})(Y_i - \bar{Y}) / \sum(X_i - \bar{X})^2 = 80/10 = 8.0$, and $\hat{\beta}_0 = 73.6 - 8.0 \times 3 = 49.6$.
Low-X students (A, B) have average $\hat{\varepsilon}^2 = 1.06$. High-$X$ students (D, E) have average $\hat{\varepsilon}^2 = 66.56$—**over 60 times larger**. The classical SE treats all five students as equally informative about the slope. But high-study students are wildly noisier, so the classical confidence interval is a lie: it claims more precision than the data actually support.

Predict Before You Peek

You have just seen that low-X students have tiny residuals and high-X students have enormous residuals. The classical 95% confidence interval ignores this and uses a single pooled estimate of the noise. Before reading further: what fraction of these nominal 95% confidence intervals do you think actually contain the true coefficient? Write down your guess.

"The Funnel of False Confidence" — Coverage Simulator

Try it: A scatterplot of hours studied vs. exam score shows data fanning out (the funnel). Two confidence bands overlay the regression line: classical (constant width, blue) and robust (varying width, green).
Key interaction: Click "Draw 100 new samples." Watch 100 confidence intervals cascade in, coloring green if they contain the truth and red if they miss. With heteroscedasticity, classical CIs achieve only ≈82% coverage while robust CIs hit the promised 95%.
Break it: Set sample size to $n = 20$ and watch the robust SEs *perform worse* than classical—the repair has a minimum sample-size cost.

4.3 Consequences for Inference

If the classical standard errors are used under heteroscedasticity:

- Confidence intervals have incorrect coverage. In a simulation of the studying DGP, the nominal 95% CI achieves only ≈82% coverage: out of every 100 intervals you construct, only about 82 contain the true coefficient, not the 95 you were promised.
- t-statistics and p-values are unreliable. A p-value of 0.03 might really correspond to a true rejection probability of 0.10—you think you have a significant result, but your evidence is much weaker than it appears.

- The coefficient estimates themselves are *not* biased—only the uncertainty is wrong. OLS still estimates the right number on average; it just gives you a dangerously wrong answer to "how sure are you?"

4.4 The Breusch–Pagan Test

Remark 4.3 (The chi-squared distribution). The **chi-squared distribution** with p degrees of freedom, written χ^2_p, is the distribution of the sum of p independent squared standard normal variables: if $Z_1, \ldots, Z_p \overset{\text{iid}}{\sim} \mathcal{N}(0,1)$, then $Z_1^2 + \cdots + Z_p^2 \sim \chi^2_p$. It is always non-negative, right-skewed, and becomes approximately normal for large p. We use it here because the Breusch–Pagan test statistic follows a χ^2_p distribution under the null hypothesis of homoscedasticity.

Remark 4.4 (R^2: How much does X explain?). The **coefficient of determination** R^2 measures the fraction of the total variation in the outcome that is explained by the predictors. If $R^2 = 0.80$, the predictors explain 80% of the variation; the remaining 20% is unexplained noise. An R^2 close to zero means the predictors have almost no explanatory power.

Before we can test for heteroscedasticity, we need to be clear about what a test statistic is in general.

Definition 4.4 (Test Statistic)**.** A *test statistic* is a single number computed from the data that measures how far the data deviates from what we would expect under the null hypothesis H_0. The key property: we must know (or be able to approximate) its distribution under H_0, so we can compute a p-value. Large values of the test statistic mean the data looks "surprising" if H_0 were true; small values mean the data looks "normal" under H_0.

Now we can state the specific test for heteroscedasticity.

The logic. If errors are homoscedastic, the squared residuals $\hat{\varepsilon}_i^2$ should be unrelated to X—knowing how much a student studied should tell you nothing about the *size* of the prediction error. If we fit a linear model with the squared residuals $\hat{\varepsilon}_i^2$ as the outcome and the original predictors X as the inputs and find a strong relationship, that is evidence against homoscedasticity.

What "regress squared residuals on regressors" means. Take the residuals from your main regression (exam score on hours studied). Square each one—this gives you a measure of how big the error was for each student, ignoring its sign. Now run a *new* regression with these squared residuals as the outcome and the original hours studied as the predictor. If this auxiliary regression has a high R^2, then hours studied predicts the error size—i.e., the errors are heteroscedastic.

Definition 4.5 (Breusch–Pagan Test)**.**

1. Estimate the main regression $Y = X\beta + \varepsilon$ by OLS. Compute residuals $\hat{\varepsilon}_i$.
2. Run a linear regression of $\hat{\varepsilon}_i^2$ on X (the same regressors as the main regression). Compute the R^2 of this auxiliary regression.
3. Under H_0: homoscedasticity, the test statistic nR^2 from this auxiliary regression follows a χ^2_p distribution asymptotically, where p is the number of regressors (excluding the constant).
4. Reject H_0 if nR^2 exceeds the critical value from the χ^2_p distribution.

Why nR^2? The R^2 from the auxiliary regression measures how much of the variation in $\hat{\varepsilon}_i^2$ is explained by X. Under H_0 (homoscedasticity), R^2 should be close to zero—X should have no power to predict the error size. Multiplying by n converts this proportion into a test statistic with a known distribution. The result that $nR^2 \sim \chi^2_p$ under H_0 is a standard asymptotic result from the theory of auxiliary regressions: when the true relationship is zero, the explained variation, scaled by n, converges to a chi-squared random variable.

Why we do not need to know σ^2. We do not need to know the true error variance. We only need to test whether the variance *changes* with X. The R^2 measures exactly that: the fraction of variance in $\hat{\varepsilon}_i^2$ that X explains. Whether the overall level of $\hat{\varepsilon}_i^2$ is large or small does not matter—what matters is whether it varies systematically with the regressors.

Contrast with the t-statistic. The t-statistic for testing a coefficient (which we will meet in Section 5) requires estimating σ via $s = \sqrt{\mathrm{RSS}/(n-p)}$, where $\mathrm{RSS} = \sum_{i=1}^n \hat{\varepsilon}_i^2$ is the **residual sum of squares**—the total squared prediction error. The Breusch–Pagan test avoids this entirely: it works through R^2, which is scale-free. This is why we can detect heteroscedasticity without knowing the baseline variance level.

4.5 Robust Standard Errors

The Repair

The **White–Huber–Eicker** ("sandwich") standard errors are computed as

$$\widehat{\mathrm{Var}}(\hat{\beta}_{\mathrm{OLS}}) = (X^\top X)^{-1} \left(\sum_{i=1}^n \hat{\varepsilon}_i^2 \, x_i x_i^\top \right) (X^\top X)^{-1},$$

where $\hat{\varepsilon}_i$ are the OLS residuals. These standard errors are *consistent under heteroscedasticity of unknown form*: you do not need to know or specify how the variance changes with X. The estimator "lets the data speak" by using each observation's own squared residual $\hat{\varepsilon}_i^2$ as a plug-in estimate for σ_i^2.
With robust SEs, the 95% CI recovers ≈95% coverage in the studying DGP simulation.

Why does this work? Because each observation's own squared residual $\hat{\varepsilon}_i^2$ serves as a plug-in estimate for its unknown variance σ_i^2, letting the data speak for itself without requiring any assumption about how the variance changes with X. The formal derivation is in Appendix A.

Warning

Robust standard errors perform poorly in small samples. When $n = 20$, the sandwich estimator can be *more biased* than the classical estimator, because each individual $\hat{\varepsilon}_i^2$ is a noisy estimate of σ_i^2, and with few observations the noise dominates. The crossover point depends on the severity of heteroscedasticity, but as a rough guide, $n \geq 50$ is typically needed for reliable sandwich SEs.

5 Why Your Significance Is Wrong: Effect Size and Power

Coefficient tracker: the studying coefficient was 10.5 (biased) in Section 3, and 10.5 ± 1.2 (wrong CI) in Section 4. Now we ask: does the effect even matter? The ghost of self-discipline inflated the number; heteroscedasticity made us overconfident about it. This section adds a third betrayal.

5.1 The Anatomy of a t-Statistic

The Seat Number That "Predicts" Your Grade

The university records each student's seat number in the exam hall—a number from 1 to 200, assigned when you walk in. Seat number has absolutely *zero* effect on your exam score. It is pure noise. No one seriously believes that sitting in seat 147 makes you smarter

than sitting in seat 12.
But you run the regression anyway, on 50,000 student-exam observations collected over 10 years. The output:

Variable	Coefficient	SE	p-value
Seat number	0.003	0.001	0.002

The coefficient is 0.003—each seat adds three *thousandths* of a point. Over 200 seats, the total difference between seat 1 and seat 200 is 0.6 points on a 100-point exam. Completely meaningless. Yet the p-value screams $p = 0.002$: "highly significant!"
How can a variable with zero true effect be "highly significant"? Because with 50,000 observations, the standard error shrinks to 0.001, and even a coefficient of 0.003 is three standard errors from zero. The t-statistic doesn't measure importance—it measures "how many standard errors is the coefficient from zero." With enough data, even the tiniest random correlation becomes "statistically significant."
The lesson: **statistical significance tells you the effect is not exactly zero. It says nothing about whether the effect matters.**

Stand Up If Significant

Classroom exercise: Everyone who thinks the seat number effect ($p = 0.002$) is "statistically significant," stand up.
(Most of the room stands.)
Now: stay standing if you think the seat number effect is *important enough to act on*—important enough that you would switch seats, or that the university should redesign its seating policy.
(Watch the room sit down in waves as they reconsider.)
Look around. Almost everyone stood up for "significant." Almost no one stayed up for "important." That gap—between standing and sitting—is the entire point of this section. Your legs just taught you the difference between statistical significance and practical importance.

Key Insight

From now on, whenever you see a "significant" result, ask yourself: *"Is this a seat number?"* That is: is the effect statistically distinguishable from zero only because the sample is enormous, while the actual magnitude is too small to matter? This question will serve you well for the rest of these notes—and for the rest of your career.

The seat number example illustrates a general phenomenon. Let us see why it happens mathematically.

The t-statistic for testing $H_0 : \beta_j = 0$ is

$$t = \frac{\hat{\beta}_j}{\mathrm{SE}(\hat{\beta}_j)}. \tag{14}$$

Why the t-statistic? We want to test whether $\beta_j = 0$. The raw estimate $\hat{\beta}_j$ is not enough because its scale depends on the units of X and Y. If we measure hours studied in minutes instead of hours, $\hat{\beta}_j$ changes by a factor of 60, but the evidence for or against $\beta_j = 0$ has not changed at all. Dividing by $\mathrm{SE}(\hat{\beta}_j)$ standardises the estimate: the result is a dimensionless number that measures "how many standard errors away from zero is our estimate?" Under H_0,

this ratio follows a t-distribution with $n-p$ degrees of freedom (or approximately $N(0,1)$ for large n). We use the t-distribution rather than the normal distribution because we *estimate* σ rather than knowing it—the extra uncertainty from estimating σ inflates the tails slightly, making extreme values a bit more likely than under the normal. With large n, the estimate of σ becomes very precise, the extra tail inflation vanishes, and the t-distribution converges to the normal.

Since $\mathrm{SE}(\hat{\beta}_j) = O(1/\sqrt{n})$, for any fixed $\beta_j \neq 0$:

$$|t| \to \infty \quad \text{as } n \to \infty.$$

Key Insight

With enough data, *every* non-zero effect becomes statistically significant, no matter how small. As we saw in the seat number example above, with $n = 50{,}000$ observations the standard error shrinks so much that even a coefficient of 0.003 becomes "highly significant" ($p = 0.002$)—despite being completely meaningless in practice. Statistically detectable, practically irrelevant.

5.2 Statistical Power

Definition 5.1 (Power)**.** The **power** of a test is the probability of correctly rejecting H_0 when H_0 is false:

$$\text{Power} = P(|t| > c_\alpha \mid \beta_j \neq 0)\,,$$

where c_α is the critical value at significance level α. In plain language: power is the probability of detecting a real effect when it exists. A test with low power will miss real effects most of the time; a test with high power will reliably find them.

Proposition 5.1 (Power Increases with n and Effect Size)**.** *For a two-sided t-test at level α with true coefficient β_j and standard deviation σ,*

$$\textit{Power} \approx \Phi\left(\frac{|\beta_j|}{\sigma/\sqrt{n}} - z_{\alpha/2}\right),$$

where Φ is the standard normal CDF and $z_{\alpha/2}$ is the upper $\alpha/2$ quantile. Power increases with both $|\beta_j|$ (effect size) and $\sqrt{n}$ (sample size).

This formula reveals two levers for increasing power: you can look for bigger effects (larger $|\beta_j|$), or you can collect more data (larger n). It also shows why under-powered studies are dangerous: with small n and a modest effect, Power can be as low as 10–20%, meaning you will miss the real effect 80–90% of the time.

5.3 The Multiple Testing Problem

Proposition 5.2 (Expected False Positives Under Multiple Testing)**.** *If m independent hypothesis tests are conducted at level α, and* all *null hypotheses are true, then*

$$\mathbb{E}[\textit{number of false positives}] = m\alpha.$$

Proof. Define indicator variables $I_j = \mathbf{1}(\text{test } j \text{ falsely rejects})$ for $j = 1, \ldots, m$. Since each $H_{0,j}$ is true and the test has level α, we have $P(I_j = 1) = \alpha$ and thus $\mathbb{E}[I_j] = \alpha$. The total number of false positives is $\sum_{j=1}^m I_j$. By linearity of expectation:

$$\mathbb{E}\left[\sum_{j=1}^m I_j\right] = \sum_{j=1}^m \mathbb{E}[I_j] = m\alpha.$$

Note: linearity of expectation holds regardless of whether the tests are independent or dependent. The result $\mathbb{E}[\text{false positives}] = m\alpha$ is therefore completely general—it requires no assumption about the dependence structure between tests. □

With $m = 1{,}000{,}000$ tests at $\alpha = 0.05$, we expect 50,000 false positives. In the studying context: suppose you test whether each of 200 classroom features (room temperature, lighting level, chair type, wall colour, distance to the cafeteria, ...) predicts exam scores. Even if *none* of these features matter, you expect $200 \times 0.05 = 10$ "significant" results by pure chance.

The Repair

The **Bonferroni correction** tests each hypothesis at level α/m, controlling the family-wise error rate at α. For genome-wide association studies ($m \approx 10^6$), the corrected threshold is $\alpha/m = 5 \times 10^{-8}$. In our 200-classroom-feature example, the corrected threshold is $0.05/200 = 0.00025$—any "significant" result that only cleared the 0.05 bar disappears under this stricter standard. Always report effect sizes and confidence intervals alongside p-values.

Remark 5.1 (The Replication Crisis). The multiple testing problem is not a theoretical curiosity—it is at the heart of the *replication crisis* that has shaken psychology, medicine, and economics since 2011. The Open Science Collaboration (2015) attempted to replicate 100 published psychology studies. Only 36 produced significant results the second time around. Among the casualties: the famous "power posing" study (Carney et al., 2010), which claimed that standing in an expansive pose for two minutes raises testosterone and lowers cortisol. The original had $p = 0.045$ with $n = 42$. Multiple replications with larger samples found no effect. The reasons are exactly what this section teaches: p-hacking (running many tests and reporting only the significant ones), underpowered studies (small n, large effect needed), and the base rate problem (most hypotheses tested are probably false, so even with $\alpha = 0.05$, a large fraction of "discoveries" are false positives).

The lesson for your own work: a single $p < 0.05$ result from an observational study is the *beginning* of an investigation, not the end. Always ask: what is the effect size? Could this survive a pre-registered replication? Is this a seat number?

The Keyboard That Saves 0.1 Seconds

A new keyboard layout saves an average of $\beta = 0.1$ seconds per email ($\sigma = 2$ seconds). Is this "significant"?

Sample size n	$\text{SE} = \sigma/\sqrt{n}$	$t = 0.1/\text{SE}$	Significant?
5	0.894	0.11	No
50	0.283	0.35	No
5,000	0.028	3.54	Yes ($p < 0.001$)

The effect never changed—it was always 0.1 seconds. Only the sample size grew. With 5,000 emails, saving one tenth of a second becomes "highly significant" ($p < 0.001$). Statistical significance tells you "the effect is not zero." It says nothing about whether the effect matters.

"The Magnifying Glass That Can't Measure" — Sample Size Simulator

Try it: A slider controls sample size n (50 to 100,000). Three linked panels show the $|t|$-statistic rising, the p-value plummeting, and the confidence interval narrowing—all while the effect size stays fixed at a tiny value.
The aha moment: The dashboard goes from gray "not significant" to screaming red "HIGHLY SIGNIFICANT ($p < 0.001$)" while the actual effect label never changes. Significance is a property of n, not of importance.

6 Why Your Model Is Wrong: Specification Error

Coefficient tracker: 10.5 (biased), 10.5 ± 1.2 (wrong CI), significant but a seat number? Now the ghost strikes again—the studying coefficient was wrong not just because of omitted variables, but because we assumed the relationship was a straight line when it is actually a curve.

6.1 The Problem of Functional Form

The Straight Ruler on a Winding Road

You're driving up a mountain on a road that curves back and forth. A friend asks you to describe the road, but you only have a straight ruler. You lay it on the map across the section you've driven so far, and it gives a decent summary—"on average, the road goes uphill at about this angle." But then you extend that straight line past where you've actually been, and it says you should be 500 feet in the air by now, flying off a cliff. The ruler was the *best* straight summary of the curvy road you had data for, but using it to predict beyond that range was a disaster.
This is exactly what the linear projection under misspecification gives you: OLS finds the best-fitting straight line through curved reality, which can look reasonable in-sample but diverges catastrophically when you extrapolate.

Suppose the true relationship is $Y = f(X) + \varepsilon$ for some nonlinear function f, but we fit the linear model $Y = X\beta + \varepsilon$. What does OLS actually recover?

Theorem 6.1 (Linear Projection Under Misspecification)**.** *If $Y = f(X) + \varepsilon$ with $\mathbb{E}[\varepsilon \mid X] = 0$, and we estimate $Y = X\beta + u$ by OLS, then the OLS estimator converges to*

$$\beta^* = \arg\min_{\beta} \mathbb{E}\left[(Y - X\beta)^2\right],$$

which is the best linear approximation to $f(X)$ in the mean-squared-error sense. This β^* equals the true β only if f is linear.

In the Exam Score DGP, the true relationship between study hours and exam score includes the term $8\log(X_1)$—a logarithmic, not linear, function of hours studied. A linear model in X_1 provides a reasonable approximation within the observed data range but diverges catastrophically when extrapolating.

6.2 Diminishing Returns to Studying

Why is the relationship logarithmic? Think about what happens when you study. The first hour of studying recovers the low-hanging fruit: you review the key definitions, skim the formula sheet, and recall the structure of the exam. The second hour solidifies that foundation. But by the

tenth hour, you are reviewing material you already know. Each additional hour buys you less and less.

Let us make this concrete. Consider a simplified version of our DGP with no controls and no noise: $Y = 50 + 8\log(X_1)$, where X_1 is hours studied and Y is the exam score.

The Diminishing Returns of Study Hours

We compute the true score at several study-hour values and compare with what a linear model would predict. Suppose the linear model is fit on students who studied between 1 and 16 hours and yields $\hat{Y} = 52.5 + 1.6\,X_1$ (a reasonable OLS fit to the log curve over this range).

Hours (X_1)	**True** $Y = 50 + 8\log(X_1)$	**Linear** $\hat{Y} = 52.5 + 1.6\,X_1$	**Error**	**Marginal gain**
1	50.0	54.1	+4.1	—
2	55.5	55.7	+0.2	+5.5
4	61.1	58.9	−2.2	+5.5
8	66.6	65.3	−1.3	+5.5
16	72.2	78.1	+5.9	+5.5

The "Marginal gain" column shows the true score increase from doubling study hours. Under the log model, every doubling adds the same $8\log(2) \approx 5.5$ points—but this means going from 1 to 2 hours gives the same boost as going from 8 to 16 hours. The absolute gain per additional hour shrinks rapidly: the jump from hour 1 to hour 2 is worth 5.5 points, but the jump from hour 9 to hour 10 is worth only $8(\log 10 - \log 9) \approx 0.9$ points. Within the observed range, the linear model's errors look tolerable—off by a few points at most. The real disaster happens outside this range.

6.3 The Extrapolation Danger

Predict Before You Peek

A student claims to have studied 30 hours for the exam. Based on the linear model $\hat{Y} = 52.5 + 1.6\,X_1$, what score would you predict? Write it down before reading further.

Key Insight

A student who claims to have studied 30 hours in one week—the linear model predicts $\hat{Y} = 52.5 + 1.6 \times 30 = 100.5$, a score of over 100 on a 100-point exam. This is not merely inaccurate; it is *logically impossible*. The log model predicts $50 + 8\log(30) \approx 77.2$. Still high, but plausible.

At 50 hours (a student who claims to have done nothing but study for an entire week), the linear model predicts 132.5. The log model predicts 81.3. One answer is absurd; the other is at least defensible.

The problem is not that the linear model is a bad model. Within the data range, it performs respectably. The problem is that the linear model makes an implicit promise about the future: it says that every additional hour of studying adds the same number of points, forever. The logarithmic model makes a different promise: each hour adds less than the last. The second promise is closer to reality, so the logarithmic model degrades gracefully outside the data range while the linear model degrades catastrophically.

The Repair

Fit the correctly specified model by transforming the predictor:

$$Y = \beta_0 + \beta_1 \log(X_1) + \beta_2 X_2 + \varepsilon.$$

Extrapolation accuracy improves dramatically. The log transformation captures the diminishing returns that are baked into the true DGP.
However, even the log model is a local approximation—at extreme extrapolation, all parametric models eventually fail. A student who studies 1,000 hours does not score $50 + 8\log(1000) \approx 105$; fatigue, burnout, and the hard ceiling of 100 points intervene. The model is better, not perfect.

"The Comfort Zone" — Extrapolation Explorer

Try it: Drag a prediction point along the x-axis (hours studied). Within the data range (1–16 hours), the background is calm blue and the error bar is tiny. Drag past the data boundary: the background turns progressively red, the error bar explodes, and the warning escalates from "Mild extrapolation" to "CATASTROPHIC: prediction is fiction."
Toggle: Show the correctly specified log model alongside the line. The log model tracks truth even far out—the contrast is stark.

7 Why Your R^2 Is Lying: Overfitting

Coefficient tracker: 10.5 (biased), wrong CI, significant-but-trivial, wrong functional form. Now we try to fix the shape by adding polynomial terms—and the ghost of self-discipline watches as a degree-15 polynomial achieves $R^2 = 0.97$ on training data and negative on test data. The model did not learn; it memorised.

7.1 The Bias–Variance Decomposition

The Three Dartboard Players

Three friends play darts. The first always throws at the same wrong spot—consistently two inches left of the bullseye. Every throw lands in the same cluster, just in the wrong place. That's high bias, low variance. The second friend aims perfectly and averages a bullseye, but her throws scatter wildly all over the board. That's low bias, high variance. The third friend accepts that the dartboard sits on a wobbly wall, so even a perfect throw gets jostled randomly. That wobble is irreducible noise—nobody can beat it. The best player is the one who aims at the right spot AND throws consistently, finding the sweet spot between the first two friends' mistakes.
This is exactly what the bias-variance decomposition says: total prediction error equals systematic mis-aim (bias squared) plus scatter in your throws (variance) plus the wobble you can never control (irreducible noise).

Theorem 7.1 (Bias–Variance Decomposition)**.** *For any estimator $\hat{f}$ of the regression function f, the expected prediction error at a point x_0 decomposes as*

$$\mathbb{E}\Big[(Y - \hat{f}(x_0))^2\Big] = \underbrace{\Big(f(x_0) - \mathbb{E}[\hat{f}(x_0)]\Big)^2}_{Bias^2} + \underbrace{\mathrm{Var}(\hat{f}(x_0))}_{Variance} + \underbrace{\sigma_\varepsilon^2}_{Irreducible\ noise}. \tag{15}$$

Why is this true? Because the prediction error splits cleanly into three independent pieces: the noise cross-term vanishes (since ε is independent of $\hat{f}$ and has mean zero), and the remaining squared error decomposes into a systematic part (bias squared) and a random part (variance). The formal derivation is in Appendix A.

7.2 Overfitting in Practice

Adding polynomial terms to a regression always increases training R^2 (since R^2 is monotonically non-decreasing in the number of regressors). In the exam score dataset, we can fit polynomials of increasing degree to predict Y from X_1. Here is what happens:

Degree	**Training R^2**	**Test R^2**
1	0.71	0.69
3	0.82	0.78
7	0.91	0.63
15	0.97	< 0

The Polynomial Beauty Contest

Quick poll: Which model would you deploy? If you picked degree 15, you just hired a model that memorises the exam answers but cannot think for itself. A training R^2 of 0.97 is not a sign of understanding—it is a sign of memorisation. The degree-3 model "knows less" but *generalises* better.

A degree-15 polynomial achieves $R^2 = 0.97$ on training data but predicts *worse than the sample mean* on held-out data. What happened? The polynomial has 16 free parameters. With enough flexibility, it can chase every bump and dip in the training data—including bumps that are pure noise. On new data, those noise-driven wiggles point in the wrong direction, and predictions become absurdly wrong.

This is the core trap: a model that "explains" 97% of the variation in your data might be explaining mostly noise. Training R^2 always goes up with complexity; test R^2 eventually crashes.

7.3 Cross-Validation

How do you choose the right level of complexity without a test set? You simulate having one.

Definition 7.1 (k-Fold Cross-Validation)**.** Partition the data into k equally sized folds. For each fold j, train the model on all data except fold j and compute the prediction error on fold j. The CV estimate of prediction error is

$$\mathrm{CV}(k) = \frac{1}{k} \sum_{j=1}^{k} \mathrm{MSE}_j \,.$$

Cross-Validation: Picking the Right Model

We have 20 students' data and two candidate models: a linear model ($Y = a + bX_1$) and a log model ($Y = a + b \log X_1$). We use 5-fold CV:

Fold	Linear MSE	Log MSE
1	42.3	28.1
2	38.7	25.9
3	51.2	31.4
4	44.1	27.8
5	39.8	26.3
Average	**43.2**	**27.9**

The log model wins on every single fold—its average MSE is 35% lower. This is not a coincidence: the log model matches the true DGP (Y depends on $\log X_1$), while the linear model systematically misses the diminishing-returns curve. Cross-validation detected the misspecification that training R^2 alone could not.

The idea is that each fold takes a turn being the "held-out" data. By averaging across folds, you get an honest estimate of how well the model predicts data it has never seen. In the exam score dataset, cross-validation correctly identifies the log-linear model as the best specification, outperforming all polynomial models.

The Repair

Using cross-validation, the log-linear model (which matches the true DGP) achieves the lowest CV error, outperforming all polynomial models. The right model family matters more than the degree of flexibility. A degree-15 polynomial has enormous capacity but wastes it memorising noise. The log-linear model has minimal capacity but uses it on the right shape.

Warning

Cross-validation itself can overfit if the number of candidate model specifications searched is large. The best CV R^2 inflates with the breadth of the search. If you try 500 different model specifications and pick the one with the best CV score, that score is optimistically biased. This is sometimes called "researcher degrees of freedom"—the more choices you make, the more chances you have to get lucky.

Three Models, Three Training Sets

We simplify the exam score setting to isolate the bias-variance tradeoff. Suppose the true relationship is $Y = 2X_1$ (every hour of studying adds exactly 2 points, with some noise). We want to predict the score of a student who studied 6 hours. The true score is $Y = 2 \times 6 = 12$.

We draw three small training sets and fit three models: Model A (constant—just predict the average), Model B (straight line), and Model C (degree-4 polynomial).

	Model A (constant)	**Model B** (line)	**Model C** (degree-4)
Prediction (Set 1)	7	10.8	8.0
Prediction (Set 2)	7	13.2	16.0
Prediction (Set 3)	7	12.2	−13.0
Average prediction	7.0	12.1	3.7
Bias^2	25.0	≈ 0	69.4
Variance	0.0	1.0	149.6
Total MSE	**25.0**	**1.0**	**219.0**

Model A always predicts 7 regardless of the training data—zero variance, but it misses the truth by 5 points every time (bias$^2 = 25$). Model B is nearly unbiased (average prediction ≈ 12) and has modest variance. Model C achieves zero training error on every dataset, yet its predictions swing from +16 to −13. The line wins decisively. Perfect in-sample fit amplifies noise into wild out-of-sample predictions.

"The U-Shaped Trap" — Bias-Variance Explorer

Try it: A slider controls polynomial degree (1 to 15). The left panel shows a stacked area chart of Bias^2 + Variance + Noise forming the classic U-shaped MSE curve. The right panel overlays 20 fitted curves from different training sets.
The aha moment: At degree 1, twenty nearly identical lines all miss the true curve (stable but wrong). At degree 15, twenty polynomials pass through their training points perfectly but look completely different from each other—wild oscillations everywhere (unstable and absurd). Press "Animate" and watch the sweet spot at degree 3–4.

8 Why Your Causal Claim Is Wrong: Endogeneity and Bad Controls

Coefficient tracker: 10.5 (biased), wrong CI, significant-but-trivial, wrong shape, memorised noise. Now we try "controlling for" post-exam confidence—and the ghost of self-discipline does not just haunt the estimate, it destroys it. The studying coefficient collapses toward zero. Another ghost, another way to be wrong.

8.1 Endogeneity

The "Hot Guys Are Jerks" Study

In 2016, a study at Brunel University found that men with more attractive bodies were more selfish and less egalitarian. SheKnows.com announced: "Really hot guys are also really not nice, says science." Gossip blogs celebrated: science had confirmed what everyone knew from dating. But had it? By the end of this section, you will see that the entire correlation could be an artifact of who you are looking at.

Why Every Movie Star's Spouse Seems Boring

Among famous actors, their spouses seem to be unusually un-funny or un-attractive. Does fame make you marry dull people? No—you're only looking at people who are already famous. People become famous either by being extremely talented or by being extremely

good-looking (or both). Once you filter for "is famous," the talented-but-plain people and the gorgeous-but-untalented people get mixed together, creating a fake tradeoff that doesn't exist in the general population. By restricting your attention to a group defined by the outcome of the very thing you're studying, you manufactured a relationship from thin air.
This is exactly what collider bias describes: when you control for a variable that is *caused by* the things you're studying, you create a spurious association that didn't exist before—more controls made the answer worse, not better.

The comedian Pete Davidson—who has dated Ariana Grande, Kate Beckinsale, and Kim Kardashian—is a walking embodiment of this paradox. He once told Page Six that people were obsessed with his dating life "because I'm ugly." But Davidson is famous for being extremely funny, not for his looks. The "mismatch" is manufactured by the fame filter: you only noticed Davidson because he cleared the talent threshold, so his looks seem below average *relative to his famous partners*. Among non-famous comedians, nobody would bat an eye.

Definition 8.1 (Endogeneity)**.** A regressor X_j is *endogenous* if $\mathrm{Cov}(X_j, \varepsilon) \neq 0$. This can arise from omitted variables, measurement error, or simultaneous causation (feedback loops).

In the exam score DGP, studying hours (X_1) are endogenous because self-discipline (U) drives both studying and scores. The structural equation for scores is $Y = 50 + 8\log(X_1) + 3X_2 + 2U + \varepsilon$, but the analyst does not observe U. When we regress Y on X_1, the OLS estimate of "the effect of studying" includes the effect of self-discipline traveling through the study-hours channel. Disciplined students study more *and* score higher for other reasons; OLS cannot tell these apart. It lumps both effects into a single inflated coefficient.

8.2 The Causal Graph

The causal structure of the exam score DGP is:

$$U \longrightarrow X_1 \longrightarrow Y \longrightarrow X_3, \qquad U \longrightarrow Y, \qquad U \longrightarrow X_2 \longrightarrow Y.$$

In words:

- Self-discipline (U) drives studying (X_1), which drives exam scores (Y).
- Self-discipline also drives scores directly (through focus, practice habits, attentiveness).
- Self-discipline drives lecture attendance (X_2), which drives scores.
- Exam scores drive self-assessed performance (X_3).

The arrow from Y to X_3 is what makes confidence a *post-treatment* variable. Confidence does not cause high scores; being well-prepared causes both high scores and high confidence. Including X_3 as a control conditions on a descendant of Y—a dangerous move, as we will see next.

8.3 Collider Bias from Bad Controls

Proposition 8.1 (Collider Bias)**.** *If $X_3 = \alpha Y + \eta_3$ (i.e., X_3 is a post-treatment variable), then the regression of Y on (X_1, X_3) yields a biased estimate of the causal effect of X_1 on Y, even if X_1 is otherwise exogenous. The bias arises because conditioning on a descendant of Y partially conditions on Y itself.*

To understand why, think about what "controlling for confidence" means. Among students with the same confidence level, those who studied fewer hours must have had something else going for them—higher self-discipline, better background knowledge, or just luck. By holding confidence fixed, you are implicitly selecting on a combination of studying and these other factors, creating a spurious association between studying and the error term.

Key Insight

You might think: "I'll just include every variable in my dataset as a control. That way I can't have omitted variable bias." This reasoning is wrong for two reasons:

1. **You can only control for variables you observe.** If the true confounder U is unmeasured, no amount of other controls helps—unless they happen to be good proxies for U. In our DGP, self-discipline is hidden. Including hours studied, lectures attended, and confidence does not capture self-discipline directly.

2. **Some observed variables are post-treatment.** Including confidence (X_3) does not remove bias—it *introduces* new bias. The variable X_3 sits downstream of the outcome Y in the causal graph. Conditioning on it opens a path that was previously blocked, allowing spurious associations to flow.

The right set of controls depends on the *causal structure*, not on what happens to be in your spreadsheet. Adding a variable is not free; it can make things worse.

Warning

The advice "control for everything available" is wrong. Controlling for post-treatment variables or colliders *introduces* bias rather than removing it. Variable selection for causal inference requires understanding the causal structure, not just statistical association.

Talent, Luck, and the Fame Filter

People have Talent (0 or 1) and Luck (0 or 1), independently assigned. A person is Famous if they have Talent OR Luck (or both).

Type	Talent	Luck	Famous?
I	0	0	No
II	0	1	Yes
III	1	0	Yes
IV	1	1	Yes

Full population: $P(\text{Talent} = 1) = 1/2$, $P(\text{Luck} = 1) = 1/2$, and $P(\text{both}) = 1/4 = 1/2 \times 1/2$. Talent and Luck are **independent**. Correlation $= 0$.
Among the Famous only (drop Type I): If someone is talented (Types III, IV), $P(\text{Lucky}) = 1/2$. If not talented (Type II only), $P(\text{Lucky}) = 1$. Suddenly, knowing a famous person is talented makes them *less* likely to be lucky!
The observation "famous people are either talented OR lucky but rarely both" is not evidence that the universe is fair—it is a mathematical artifact of selecting on a collider.

"The Paradox of Conditioning" — Collider Explorer

Try it: A scatterplot of Talent vs. Luck for 1,000 people. All dots are gray; no correlation visible. Toggle "Condition on Fame" and watch: most dots fade, the highlighted subset reveals a clear negative slope. A DAG panel alongside animates: a box appears around the Fame node, and a dashed red "spurious association" arc draws between Talent and Luck.
Slider: Fame threshold—from 10th to 90th percentile. At low thresholds (most peo-

ple famous), the spurious correlation is weak. At high thresholds (top 5%), it becomes extreme.

8.4 Instrumental Variables

When X is endogenous, OLS is stuck. No matter how large your sample, the estimate converges to the wrong value. The probability limit of $\hat{\beta}_{\text{OLS}}$ includes a bias term that depends on $\text{Cov}(X, U)$—and this term does not shrink with the sample size. You could collect data on every student in every university in the world, and OLS would still give you the wrong answer. We need a fundamentally different strategy.

The idea behind instrumental variables is to find a source of variation in X that is "clean"—uncontaminated by the confounder U. If we can isolate this clean variation and look at how Y responds to it, we recover the causal effect.

In the exam score context: studying hours are contaminated by self-discipline. But suppose dorm room assignments at the university are random, and your roommate's study habits influence how much *you* study (peer effects). If your roommate is a diligent studier, you tend to study more; if your roommate plays video games all evening, you study less. Crucially, your roommate's study hours are correlated with *your* study hours (relevance), but your roommate's habits have no direct effect on *your* exam score other than through influencing how much you study (exclusion). Your roommate does not take the exam for you, does not whisper answers during the test, and does not grade your paper. This is the instrument.

Definition 8.2 (Instrument)**.** A variable Z is a valid *instrument* for X in the equation $Y = X\beta + \varepsilon$ if:

1. **Relevance:** $\text{Cov}(Z, X) \neq 0$. The instrument must actually affect the endogenous variable. A roommate's study hours must influence your study hours.
2. **Exclusion:** $\text{Cov}(Z, \varepsilon) = 0$. The instrument affects Y *only* through X. Your roommate's hours affect your score only because they change how much you study, not through any other channel.

Try It Yourself: Name That Instrument

You want the causal effect of study hours (X_1) on exam score (Y), but self-discipline (U) confounds the relationship. For each candidate instrument below, evaluate whether it satisfies *relevance* (does it affect study hours?) and *exclusion* (does it affect exam scores **only** through study hours?).

(a) **The student's IQ score.** Relevance: plausible—smarter students may study differently. Exclusion: **fails.** IQ directly affects exam scores through faster comprehension, better reasoning, and test-taking ability, independent of study hours. IQ is a confounder, not an instrument.

(b) **Whether the library was open late that week** (determined by a random maintenance schedule). Relevance: yes—when the library closes early, students have fewer hours available to study. Exclusion: **plausible.** The maintenance schedule is random, so it is unrelated to student ability or discipline. Library closures affect exam scores only by limiting study time. *This is a valid instrument.*

(c) **The student's Instagram screen time.** Relevance: plausible—more screen time likely means fewer study hours. Exclusion: **fails.** Screen time reflects self-discipline, which directly affects exam scores through focus, persistence, and class engagement. The instrument is contaminated by the very confounder we are trying to bypass.

The lesson: a valid instrument must provide variation in X that is "clean"—uncontaminated by any backdoor path to Y. Options (a) and (c) fail because they are entangled with the confounder U. Only (b), which is driven by an external random event, passes both conditions.

Theorem 8.1 (IV Consistency)**.** *If Z is a valid instrument, the IV estimator*

$$\hat{\beta}_{IV} = (Z^\top X)^{-1} Z^\top Y \tag{16}$$

is consistent for β:

$$\text{plim } \hat{\beta}_{IV} = \text{plim } \frac{Z^\top Y/n}{Z^\top X/n} = \frac{\text{Cov}(Z, Y)}{\text{Cov}(Z, X)} = \beta.$$

What does consistency mean, and why do we want it? An estimator is *consistent* if it converges to the true parameter value as the sample size grows to infinity. OLS, when X is endogenous, is *not* consistent—it converges to the wrong value, forever. Collecting ten times more data does not help; the bias is baked into the method, not the sample. IV is consistent: with enough data, it gets arbitrarily close to the true causal effect.

This is why we tolerate IV's larger variance. In finite samples, IV estimates are noisier than OLS—the confidence intervals are wider, the point estimate jumps around more from sample to sample. But IV is noisy around the *right* answer, while OLS is precise around the *wrong* answer. A noisy estimate of the right thing beats a precise estimate of the wrong thing, at least when your goal is to learn the truth.

8.5 The Strength of the Instrument

Not all instruments are equally useful. An instrument that barely moves X is almost as bad as no instrument at all.

Definition 8.3 (F-Statistic of the First Stage)**.** The F-statistic from the first-stage regression (X on Z and controls) measures how strongly the instrument Z predicts the endogenous variable X. Formally, it tests H_0: the coefficient on Z in the first stage is zero.

A large F means the instrument has strong predictive power for X. The rule of thumb: $F > 10$ indicates a sufficiently strong instrument. When $F < 10$, the instrument is "weak": it barely moves X, so the IV ratio $\mathrm{Cov}(Z, Y)/\mathrm{Cov}(Z, X)$ becomes unstable (you are dividing by something close to zero). Weak instruments produce wildly variable estimates with enormous confidence intervals—potentially more biased than OLS.

To see why weak instruments are dangerous, think about the IV formula: $\hat{\beta}_{\mathrm{IV}} = \mathrm{Cov}(Z, Y)/\mathrm{Cov}(Z, X)$. If $\mathrm{Cov}(Z, X)$ is close to zero, you are dividing a small number by an even smaller number. Small random fluctuations in the denominator get amplified into huge swings in the estimate. In the extreme case where Z is completely unrelated to X, you are dividing noise by noise—the result is meaningless.

Warning

Weak instruments ($\mathrm{Cov}(Z, X) \approx 0$) cause the IV estimator to be highly variable and unreliable. When the first-stage F-statistic is below 10, IV confidence intervals are extremely wide and the estimator may be more biased than OLS. A weak instrument is not just useless—it is actively harmful, because it gives you false confidence that you have solved the endogeneity problem when you have not.

8.6 A Numerical Example: Roommates and Exam Scores

Roommate Study Hours as an Instrument

We want the causal effect of study hours (X_1) on exam score (Y). The problem: self-discipline (U) boosts both. We use roommate's average study hours (Z) as an instrument. Roommates are randomly assigned, so Z is independent of self-discipline.
Suppose the true DGP (simplified) is:

$$\begin{aligned} X_1 &= 1 + Z + U, \\ Y &= 50 + 3\,X_1 + 2\,U + \varepsilon. \end{aligned}$$

The true causal effect of one extra hour of studying is $\beta_1 = 3$ points.

Student	Z (roommate hrs)	U (self-discipline)	$X_1 = 1 + Z + U$	$Y = 50 + 3X_1 + 2U$	ε
A	1	2	4	66	0
B	2	-1	2	54	-2
C	3	0	4	62	0
D	4	-2	3	55	-4
E	5	1	7	73	2

The noise ε values are chosen so that $\mathrm{Cov}(Z, \varepsilon) = 0$ (the exclusion restriction holds exactly in this example).
Why OLS is biased. OLS regresses Y on X_1 directly:

$$\hat{\beta}_{\mathrm{OLS}} = \frac{\mathrm{Cov}(X_1, Y)}{\mathrm{Var}(X_1)} = \frac{30.5}{3.5} \approx 8.71.$$

The true effect is 3, but OLS says 8.71—nearly three times too large! The reason: disciplined students (high U) both study more and score higher. OLS attributes the self-discipline premium to studying.

Why IV works. IV uses only the variation in X_1 that comes from the instrument Z:

$$\hat{\beta}_{\text{IV}} = \frac{\text{Cov}(Z, Y)}{\text{Cov}(Z, X_1)} = \frac{6.0}{2.0} = 3.0.$$

Exactly the true causal effect! The instrument isolates the "clean" variation in study hours—the part driven by who your roommate happens to be, not by your own self-discipline. By looking at how scores respond to this clean variation, IV strips away the contamination.

Verification. We can check the instrument validity conditions:

- *Relevance:* $\text{Cov}(Z, X_1) = 2.0 \neq 0$. Roommate hours do predict your study hours. ✓
- *Exclusion:* $\text{Cov}(Z, U) = 0$ by random assignment, and $\text{Cov}(Z, \varepsilon) = 0$ by construction. Your roommate's habits affect your score only through your own studying. ✓

Remark 8.1. In this numerical example, $\hat{\beta}_{\text{OLS}} = 8.71$ while $\beta_{\text{true}} = 3$. The bias is $+5.71$—OLS nearly triples the true effect. This is not a small distortion. If a university used the OLS estimate to decide how many study hours to mandate, it would wildly overestimate the benefit and perhaps impose unreasonable requirements. Getting the causal effect right is not an academic exercise; it has real policy consequences.

"The Backdoor and the Detour" — IV Strength Simulator

Try it: Three linked panels show the first stage (Z vs X_1, roommate hours vs own study hours), reduced form (Z vs Y, roommate hours vs exam score), and structural (X_1 vs Y) relationships. The structural panel overlays OLS (blue, biased), truth (gold), and IV (green) lines.

Key slider: Instrument strength (0.01 to 0.95). With a strong instrument, IV hugs the truth. Drag toward zero: the first-stage scatter becomes noise, the F-statistic drops below 10 (turns red), and the IV sampling distribution *explodes*. A weak instrument is worse than no instrument at all.

9 The Real World — When All Pathologies Strike at Once

Coefficient tracker: 10.5 (biased), wrong CI, significant-but-trivial, wrong shape, memorised noise, destroyed by bad controls. Now we meet the tutoring regression: coefficient = −3. The ghost is not just present—it flipped the sign. The ghost is running the show.

9.1 The Lalonde Dataset

The previous sections studied one pathology at a time in a clean simulated world. Real data is not so polite—every pathology arrives simultaneously, and none announces itself.

In the lab series, this section transitions from simulated data to the Lalonde (1986) dataset, a benchmark in causal inference. The question: does a job training programme increase earnings?

- **Observational estimate (OLS):** negative coefficient—training appears to *reduce* earnings.
- **Experimental estimate (RCT):** positive and large (≈$1,800).
- The two estimates do not even have the same sign.

All standard diagnostics pass: VIFs are low, Breusch–Pagan is non-significant, robust standard errors barely change. The problem is not in the model—it is in the *data*. The observational and experimental groups differ systematically on unobservables.

9.2 The Studying Analog

The Lalonde story has a direct parallel in our exam score world.

Does Tutoring Help?

The university offers free tutoring sessions to any student who signs up. You collect data on 500 students: whether they attended tutoring, and their final exam score. A simple OLS regression says tutoring *decreases* scores by 3 points.

Should you shut down the tutoring programme? Of course not. The problem is self-selection: students who sign up for tutoring are predominantly those who are struggling—they have low self-discipline (U), missed lectures (X_2), and studied fewer hours (X_1). These are precisely the students who would have scored poorly anyway. The tutoring coefficient is negative not because tutoring hurts, but because weak students are the ones who seek it out.

Now imagine an experiment: you randomly assign half the students to attend tutoring regardless of whether they would have chosen it. The experimental estimate shows tutoring *raises* scores by 8 points. The observational estimate had the wrong sign because it confused "who selects into tutoring" with "what tutoring does."

This is the observational researcher's nightmare. You cannot diagnose the problem by staring at the regression output. The model looks fine. The standard errors are well-behaved. The R^2 is respectable. Yet the central conclusion is not just imprecise—it points in the wrong direction entirely.

9.3 Sensitivity Analysis: Cinelli–Hazlett Framework

When you cannot run an experiment, you can at least ask: *how strong would a hidden confounder need to be to explain away my result?* The Cinelli–Hazlett framework formalises this question.

Remark 9.1 (Partial R^2). The **partial** R^2 of a variable X with respect to an outcome Y, after controlling for other variables W, measures how much *additional* variation X explains beyond what W already explains. Formally: regress Y on W to get residuals $\tilde{Y}$, regress X on W to get residuals $\tilde{X}$, then the partial R^2 is the ordinary R^2 from regressing $\tilde{Y}$ on $\tilde{X}$. A partial R^2 of 0.10 means X explains 10% of the variation in Y that the other variables left on the table. This is the natural measure for the Cinelli–Hazlett framework, because it captures how much "unexplained work" a confounder would need to do.

Definition 9.1 (Robustness Value)**.** The **robustness value** (RV) is the minimum strength of association (measured in partial R^2) that an unobserved confounder would need to have with both the treatment and the outcome to explain away the observed result.

Formally, define

$$\mathrm{RV}_{q,\alpha} = \min\{r : \text{adjusted estimate crosses } q \cdot \hat{\beta} \text{ at level } \alpha\}.$$

A small RV indicates fragility: a weak confounder suffices to nullify the result. A large RV indicates robustness: only a confounder with a very strong relationship to both treatment and outcome could overturn the finding.

In our tutoring example, the observational estimate that "tutoring hurts" has a tiny robustness value. A confounder (like self-discipline) with even modest partial R^2 with both tutoring attendance and exam score would flip the sign—which is exactly what happens when you account for self-selection in the experimental design.

How Fragile Is the Tutoring Estimate?

The observational regression of exam score on tutoring attendance (with controls for prior GPA and attendance) yields $\hat{\beta}_{\text{tutor}} = -4.2$ with SE $= 1.8$ and $t = -2.33$ ($p = 0.02$). The estimate is "significant" — tutoring appears to *hurt*.
How strong would an unobserved confounder need to be to explain this away?
Using the Cinelli–Hazlett framework, we compute the **robustness value**: RV $= 0.04$. This means: a confounder that explains just 4% of the residual variance in *both* tutoring attendance and exam scores would be enough to reduce the coefficient to zero.
Is 4% a lot? Compare it to the observed covariates:

Covariate	**Partial R^2 with tutoring**	**Partial R^2 with score**
Prior GPA	0.12	0.18
Lecture attendance	0.08	0.09

Both observed covariates have partial R^2 values far above the robustness value of 0.04. An unobserved confounder weaker than lecture attendance would suffice to flip the sign. The result is **extremely fragile**: self-discipline (U), which we know exists but cannot measure, almost certainly exceeds this threshold.
Contrast with the experimental estimate: The randomised trial gave $\hat{\beta}_{\text{tutor}} = +1{,}800$. The observational estimate was not just wrong — it had the wrong *sign*. The robustness value warned us: the result was one weak confounder away from flipping.

Key Insight

The Lalonde dataset—and its tutoring analog—teaches the hardest lesson in applied statistics: a well-fitting model with clean diagnostics can still give you an answer that is not just wrong, but *directionally* wrong. Diagnostics check the model's internal consistency. Only the research design can tell you whether the estimate is credible. When in doubt, compute the robustness value. If a plausible confounder could flip your result, treat the finding with extreme scepticism.

Remark 9.2 (The COVID Smoking Paradox). A striking modern example of collider bias in the wild: early in the COVID-19 pandemic, data from Chinese hospitals seemed to show that smokers were *protected* against COVID-19—only 8% of hospitalised COVID patients were smokers, versus 26% in the general population. Headlines announced a "smoker's paradox." The explanation? Hospitalisation is a collider: smokers were already in hospitals for smoking-related illness. Among hospitalised patients, smokers appeared less likely to have COVID—a completely spurious protective effect. Griffith et al. (2020) demonstrated this formally using UK Biobank data. The lesson: whenever you condition on a variable that is caused by both the treatment and the outcome, expect to see ghosts.

10 The Redemption — Regression Discontinuity Design

Remark 10.1 (Potential outcomes). The **potential outcomes** framework imagines two versions of each individual: Y_1 (the outcome they *would* experience with treatment) and Y_0 (the outcome they *would* experience without treatment). The causal effect for individual i is $Y_{1i} - Y_{0i}$. The fundamental problem of causal inference is that we can never observe both Y_1 and Y_0 for the same person—each individual is either treated or not, so one potential outcome is always missing. RDD, introduced next, cleverly sidesteps this problem by comparing individuals who are nearly identical but fall on opposite sides of a treatment cutoff.

Coefficient tracker: 10.5 (biased), wrong CI, significant-but-trivial, wrong shape, memorised, destroyed by bad controls, wrong sign. Eight ways to be wrong. Now—for the first time—the coefficient earns our trust. The ghost of self-discipline cannot haunt a design where treatment is assigned by a cutoff, not by choice.

10.1 When Design Earns Trust

After eight sections of everything going wrong, you might be wondering: is regression ever trustworthy? The answer is yes—but only when the *research design* earns that trust. This section is the redemption arc.

Regression works brilliantly when the world provides a natural experiment. The **regression discontinuity design** (RDD) exploits a sharp threshold to identify causal effects—no randomisation required.

The Kid Who Scored 69 vs. The Kid Who Scored 70

Your school says anyone who scores 70 or above on the final exam gets into the honours class next year. The kid who scored 70 and the kid who scored 69 are basically the same student—maybe one got lucky on one question, maybe the other had a headache. They studied the same amount, have the same ability, live in the same neighbourhoods. But one gets the honours class and the other does not. If you come back a year later and the 70-kid is doing noticeably better, that gap cannot be explained by the kids being different—they were virtually identical. The jump in outcomes right at the cutoff *is* the real effect of the honours class itself.

This is exactly what the regression discontinuity design measures: the difference in outcomes just above versus just below the cutoff, where people are essentially identical, gives you the causal effect without a randomised experiment.

Definition 10.1 (Sharp RDD)**.** Suppose treatment is assigned deterministically by a *running variable* R at a cutoff c:

$$D_i = \mathbf{1}(R_i \geq c).$$

Under continuity of potential outcomes at c, the **local average treatment effect** at the cutoff is

$$\tau_{\text{RDD}} = \lim_{r \downarrow c} \mathbb{E}[Y \mid R = r] - \lim_{r \uparrow c} \mathbb{E}[Y \mid R = r]. \tag{17}$$

Theorem 10.1 (RDD Identification)**.** *If $\mathbb{E}[Y_0 \mid R = r]$ and $\mathbb{E}[Y_1 \mid R = r]$ are continuous at $r = c$, then*

$$\tau_{RDD} = \mathbb{E}[Y_1 - Y_0 \mid R = c].$$

The design eliminates confounding locally *at the cutoff because units just above and just below the threshold are nearly identical in all characteristics.*

10.2 Example: Scholarship Cutoff

Students with exam scores just above a threshold receive scholarships; those just below do not. Near the cutoff, the two groups are essentially identical—they studied similar amounts, attended similar numbers of lectures, and have similar (unobserved) self-discipline levels. The jump in outcomes at the threshold identifies the causal effect of the scholarship.

10.3 Bandwidth Choice

The RDD estimator faces a bias–variance tradeoff controlled by the *bandwidth* h:

- **Narrow bandwidth:** low bias (observations are close to the cutoff), high variance (few observations).
- **Wide bandwidth:** low variance (more observations), high bias (observations far from the cutoff may not be comparable).

Think of it this way: the student who scored 69.5 is almost certainly comparable to the student who scored 70.5. But is the student who scored 40 comparable to the student who scored 90? Almost certainly not. The bandwidth determines how far from the cutoff you are willing to look. Optimal bandwidth selection methods (e.g., Imbens–Kalyanaraman or Calonico–Cattaneo–Titiunik) balance these two forces automatically.

Scholarship Scores and Outcomes

Students who score ≥ 50 on an entrance exam receive a scholarship. Data generated from $Y = 60 + 0.5R + 5D$ (true treatment effect $\tau = 5$):

Student	R (exam score)	Scholarship?	Y (outcome)
1	43	No	81.5
2	45	No	82.5
3	47	No	83.5
4	49	No	84.5
5	51	Yes	90.5
6	53	Yes	91.5
7	55	Yes	92.5
8	57	Yes	93.5

Naïve above-vs-below comparison: $\bar{Y}_{\text{above}} - \bar{Y}_{\text{below}} = 92.0 - 83.0 = 9.0$. But this includes the baseline trend (higher-scoring students do better anyway).
Fit lines on each side and extrapolate to the cutoff ($R = 50$): below gives $\hat{Y}(50) = 85.0$, above gives $\hat{Y}(50) = 90.0$. The RD estimate: $\hat{\tau} = 90.0 - 85.0 = 5.0$—exactly the true treatment effect.

The Repair

The RDD does not require controlling for confounders. The design itself ensures comparability near the cutoff. When the autopsy report is run on the RDD estimate, all indicators are green: the regression is trustworthy because the *design* is trustworthy. This is why causal inference researchers love RDDs: the credibility comes from the structure of the problem, not from statistical adjustments you hope are sufficient.

"The Cliff" — Bandwidth Explorer

Try it: A scatterplot of Y vs. exam score R with a vertical cutoff line. Two local regression lines meet at the cutoff; the gap is the treatment effect. A bandwidth slider (2 to 50) controls how many points are used: narrow bandwidth shows a clean jump with wide confidence intervals, wide bandwidth smooths the jump away.
Break it: Drag the cutoff away from the true value (50) and watch the "treatment effect" vanish—a built-in falsification test.

11 What "Controlling For" Actually Means — The Frisch–Waugh–Lovell Theorem

Coefficient tracker: we have seen the studying coefficient be inflated, misshapen, memorised, destroyed, and sign-flipped. Every time someone said "controlling for," something happened to that number. This section finally explains what "controlling for" actually means—mechanically, algebraically, with no ambiguity.

11.1 Motivation

"Controlling for X_2" is one of the most commonly used phrases in empirical research. It is also one of the most commonly misunderstood. When we say "the coefficient on X_1, controlling for X_2," we do *not* mean "holding X_2 constant." We do not mean "looking only at observations where X_2 has the same value." We mean something much more specific, and the Frisch–Waugh–Lovell (FWL) theorem tells us exactly what.

We mean: the coefficient from regressing the part of Y unexplained by X_2 on the part of X_1 unexplained by X_2. Both sides get residualised. This is not a thought experiment—it is an algorithm you can execute step by step.

Is It the Coaching or Just the New Shoes?

Your friend claims her new running coach made her faster. But she also bought expensive new running shoes at the same time. You have data on 50 runners: each runner's speed improvement (Y), coaching hours (X_1), and shoe quality score (X_2).

How do you isolate the coach's contribution? Here are the three concrete steps:

Step 1: Remove what shoes explain about speed. You have speed data and shoe data for all 50 runners. Regress speed improvement on shoe quality: this tells you "for each point of shoe quality, runners got X seconds faster." Now compute each runner's residual: their ACTUAL speed improvement minus what shoes alone would predict. Call these residuals $\tilde{Y}$. These are the speed improvements that shoes *cannot* explain.

Step 2: Remove what shoes explain about coaching. Regress coaching hours on shoe quality: maybe the coach recommended the shoes, so runners with better shoes also got more coaching. Compute residuals: each runner's ACTUAL coaching hours minus what shoe quality would predict. Call these $\tilde{X}_1$. These are the coaching hours that have nothing to do with shoes.

Step 3: Compare the leftovers. Regress $\tilde{Y}$ on $\tilde{X}_1$. The slope is the coach's true contribution to speed, cleaned of the shoe effect.

"But wait," you might ask, "how can I subtract out the shoe effect if I haven't measured it yet?" You HAVE measured it—you have shoe quality scores for all 50 runners. The regression in Step 1 IS the measurement. You are not subtracting a hypothetical; you are subtracting a prediction computed from your actual data. FWL is not a thought experiment; it is an algorithm you can execute step by step on a spreadsheet.

11.2 The FWL Theorem

Theorem 11.1 (Frisch–Waugh–Lovell)**.** *Consider the regression $Y = X_1\beta_1 + X_2\beta_2 + \varepsilon$. Let $M_2 = I - X_2(X_2^\top X_2)^{-1}X_2^\top$ be the residual-maker matrix that projects orthogonally to the column space of X_2. Then*

$$\hat{\beta}_1 = (X_1^\top M_2 X_1)^{-1} X_1^\top M_2 Y. \tag{18}$$

That is, $\hat{\beta}_1$ from the full regression is identical *(to numerical precision) to the coefficient obtained by:*

1. *Regressing Y on X_2 and taking residuals: $\tilde{Y} = M_2 Y$.*

2. *Regressing X_1 on X_2 and taking residuals:* $\tilde{X}_1 = M_2 X_1$.
3. *Regressing $\tilde{Y}$ on $\tilde{X}_1$ (no intercept).*

Why is this true? Because the partitioned normal equations let you solve for $\hat{\beta}_1$ by substituting out $\hat{\beta}_2$, and the algebra naturally produces the residual-maker matrix M_2 on both sides—showing that "controlling for X_2" is literally projecting X_2 out of both Y and X_1. The formal derivation is in Appendix A.

Key Insight

"Controlling for" means residualising—stripping out the linear relationship with the control variables from *both* sides of the equation. Not just the outcome. Not just the predictor. Both. The partial regression plot (also called an added-variable plot) visualises this: it plots $\tilde{Y}$ against $\tilde{X}_1$. The slope of this scatterplot *is* $\hat{\beta}_1$ from the full multiple regression.

Remark 11.1. FWL is not just a theoretical curiosity. When you read a paper that says "controlling for age and income, education increases wages by…", the author is (perhaps unknowingly) describing exactly the FWL procedure: strip age and income out of both sides, then measure what is left.

11.3 Numerical Example

Exercise, Sleep, and Mood

True DGP: $Y = 2 + 4.684\,X_1 + 1.45\,X_2 + \varepsilon$.
We want the effect of exercise (X_1, hours) on mood (Y, score 1–20), controlling for sleep (X_2, hours). Here is our data:

Person	X_1 (exercise)	X_2 (sleep)	Y (mood)
A	1	2	3
B	2	4	7
C	3	6	10
D	5	8	18
E	4	10	12

Step 1: Regress Y on X_2. The fitted slope is 1.45. Now compute each person's residual: their actual mood minus what sleep alone predicts. Notice: this step REQUIRES having data on both Y and X_2, which we do.
Residuals: $\tilde{Y} = (-1.2,\ -0.1,\ 0.0,\ +5.1,\ -3.8)$.
Step 2: Regress X_1 on X_2. The fitted slope is 0.45. Residuals: each person's actual exercise hours minus what sleep would predict. Again, we use our observed data on X_1 and X_2—nothing hypothetical here.
Residuals: $\tilde{X}_1 = (-0.2,\ -0.1,\ 0.0,\ +1.1,\ -0.8)$.
Step 3: Regress $\tilde{Y}$ on $\tilde{X}_1$. Slope $= 8.9/1.9 = \mathbf{4.684}$.
Full multiple regression (Y on X_1 and X_2): coefficient on $X_1 = \mathbf{4.684}$.
They match—not approximately, but to every decimal place. "Controlling for sleep" literally means: remove everything sleep can explain from *both* mood and exercise, then see what is left.

11.4 What Happens with Bad Controls

The FWL theorem does not just explain what "controlling for" means when you get it right. It also reveals, with painful clarity, what happens when you control for the wrong variable.

Key Insight

When you "control for" X_3 (the post-exam self-assessment), FWL tells you what you are actually doing: you are removing the linear relationship between self-assessment and BOTH the outcome (exam score) and the predictor (study hours). Because self-assessment is driven by the actual exam score—it is essentially $0.9 \cdot Y$ plus noise—stripping it out removes genuine signal, not noise. You are asking: "among students who rated their performance the *same*, does studying more help?" But students with the same self-assessment who studied different amounts must differ on something else (like self-discipline) in exactly the offsetting direction. The coefficient on study hours collapses toward zero—"controlling for" a post-treatment variable did not clean your estimate, it destroyed it.

"The Double Residual" — FWL Animator

Try it: Three panels. Panel 1: a 3D scatter of (X_1, X_2, Y) with the regression plane. Panel 2: two mini-scatters showing the residualisation (residual bars appear as regressions are fit). Panel 3: the partial regression plot—residuals vs. residuals.
The aha moment: Click "Animate." Watch regression lines fit in Panel 2, residual bars appear, then the residual dots "fly" into Panel 3 and reassemble into a new scatter. The slope reads 4.684. Check the multiple regression output: $\beta_1 = 4.684$. Exact match.

12 Diagnostic Toolkit Summary

Coefficient tracker: the studying coefficient has been 10.5 (biased), 10.5 ± 1.2 (wrong CI), "significant" but trivial (seat number), 1.6 per hour (wrong shape), $R^2 = 0.97$ *(memorised), near zero (destroyed by bad control),* -3 *(wrong sign), 3.0 (IV—finally correct), and 5.0 (RDD—correct by design). Nine versions of the same question. The diagnostics below would have flagged at most two of these problems.*

Think of these diagnostics as the instrument panel in a cockpit. Each gauge checks one specific condition—but all gauges reading green does not mean the flight plan is correct.

The following diagnostics are used across all notebooks:

Diagnostic	Tests For	Interpretation
Residuals vs. Fitted	Nonlinearity, heteroscedasticity	Residuals should show no pattern; funnel shape indicates heteroscedasticity
Normal Q–Q plot	Non-normality of residuals	Points should lie on the 45° line
Scale–Location	Heteroscedasticity	$\sqrt{\lvert\text{standardised residuals}\rvert}$ should be flat against fitted values
Residuals vs. Leverage	Influential observations	Points with high leverage *and* large residuals (Cook's $d > 1$) are influential. Cook's distance measures how much the fitted values change when observation i is deleted: $d_i = \frac{(\hat{Y}-\hat{Y}_{(i)})^\top(\hat{Y}-\hat{Y}_{(i)})}{p\cdot\hat{\sigma}^2}$.
VIF	Multicollinearity	$\text{VIF}_j > 10$ suggests problematic collinearity. The *Variance Inflation Factor* for predictor j is $\text{VIF}_j = 1/(1-R_j^2)$, where R_j^2 is the R^2 from regressing X_j on all other predictors. A VIF of 10 means that collinearity inflates the variance of $\hat{\beta}_j$ by a factor of 10.
Breusch–Pagan	Heteroscedasticity	H_0: homoscedasticity; reject if $p < \alpha$
Durbin–Watson	Autocorrelation	Statistic near 2 indicates no autocorrelation
Jarque–Bera	Non-normality	H_0: residuals are normally distributed

Warning

In the tutoring example (Section 9), every diagnostic passed—VIFs low, Breusch–Pagan non-significant, robust standard errors barely changed—yet the OLS estimate had the wrong sign. The model was internally consistent: residuals looked fine, no multicollinearity, no heteroscedasticity. But the *research design* was fatally flawed. Self-selection into tutoring created omitted variable bias that no diagnostic can detect.

Diagnostics check whether the model's assumptions about the error term are satisfied. They cannot check whether you have the right variables in the model or whether the treatment assignment mechanism is ignorable. That is a question about the world, not about the residuals.

13 Theorem Map: How the Results Connect

Coefficient tracker — the full history:

Section	Coefficient	What was wrong
3 (OVB)	10.5	Inflated by the ghost of self-discipline
4 (Heteroscedasticity)	10.5 ± 1.2	CI too narrow (wrong SEs)
5 (Significance)	0.003*** (seat number)	Meaningless but "significant"
6 (Specification)	1.6 per hour (linear)	Wrong shape—log truth, linear fit
7 (Overfitting)	degree-15 poly, $R^2 = 0.97$	Memorising noise
8 (Bad controls)	≈ 0 (controlling for confidence)	Destroyed by collider
9 (Real world)	-3 (tutoring)	Wrong sign—the ghost is running the show
8 (IV)	**3.0** (via roommate)	**Finally correct**
10 (RDD)	**5.0** (scholarship cutoff)	**Correct by design**

Nine versions of the same question. Seven wrong answers. Two right ones. The two right ones did not come from better statistics—they came from better *design*.

Before the final summary, let us step back and see how all nine results connect. Each theorem in this course relaxes an assumption that the previous one relied on, forming a chain of progressively more honest conclusions.

NB	Theorem	Generalisation Relationship
1	Omitted Variable Bias	Foundation: bias from omitting a relevant variable
2	Sandwich Variance	Relaxes $\mathrm{Var}(\varepsilon) = \sigma^2 I$; OVB is a special case of endogeneity
3	Power Analysis	$t = \hat{\beta}/\,\mathrm{SE}(\hat{\beta})$; builds on correct SEs from NB 2
4	Linear Projection	Best linear approx. to $f(X)$; relaxes f linear
5	Bias–Variance Tradeoff	$\mathrm{MSE} = \mathrm{Bias}^2 + \mathrm{Var} + \sigma^2$; generalises specification error
6	IV Consistency	If $\mathbb{E}[\varepsilon X] \neq 0$ but $\mathbb{E}[\varepsilon Z] = 0$; generalises OVB repair
7	Sensitivity Analysis	Robustness value quantifies fragility of any estimate
8	RDD Identification	Continuity at cutoff $\Rightarrow$ causal identification
1.5	Frisch–Waugh–Lovell	What "controlling for" means algebraically

Read the table from top to bottom and you see a story: each theorem relaxes one assumption that the previous theorem relied on. Classical regression assumes everything is fine. Each section showed you one way it is not—and the next section's theorem is the response.

The meta-lesson: each theorem relaxes an assumption that justified the previous conclusion. Classical variance assumes homoscedasticity—the sandwich estimator does not. The linear model assumes the true function is linear—the linear projection result tells you what OLS estimates when it is not. Omitted variable bias is a special case of endogeneity—instrumental variables handle the general case.

The chain has a direction. You cannot meaningfully discuss power (NB 3) without first understanding correct standard errors (NB 2). You cannot appreciate the robustness value (NB 7) without understanding what confounding looks like (NB 1). And the FWL theorem (NB 1.5) retroactively clarifies every "controlling for" claim you made along the way.

In the exam score DGP, each theorem answers a different version of the same question: "what is the true effect of studying on exam scores?" OVB tells you the naïve estimate is too high. Heteroscedasticity tells you the uncertainty around that estimate is wrong. Specification error tells you the functional form is wrong. IV consistency tells you how to fix the confounding. RDD tells you how a scholarship cutoff can identify causal effects without any of these corrections. And FWL tells you what it actually means when you "control for lectures attended."

14 Conclusion

Remember the Ofqual algorithm from the Introduction? The one that decided half a million students' futures on a Tuesday and was scrapped by Thursday? You now have the vocabulary to say exactly what went wrong:

- It had **omitted variable bias**—it used school-level averages instead of individual ability. The ghost of student potential was hiding inside the school-reputation coefficient.

- It had **specification error**—it assumed linear relationships between prior results and predicted grades.
- It had **bad controls**—it conditioned on school characteristics that were downstream of socioeconomic status, a collider.
- And no one ran the **sensitivity analysis** that would have revealed how fragile the estimates were.

Every failure mode we studied was present. All gauges were green. The plane flew to the wrong destination. And half a million students paid the price.

These notes have formalised the eight core failure modes of OLS regression demonstrated in the Regression Autopsy lab series:

1. **Omitted variable bias** shifts your coefficient—OLS attributes self-discipline's effect to study hours, inflating the estimated return to studying.
2. **Heteroscedasticity** corrupts your standard errors—students who study more have more variable exam scores, and the default SE formula does not account for this.
3. **Large samples** make trivial effects "significant"—with 50,000 student records, a 0.01-point improvement per study hour achieves $p < 0.001$.
4. **Misspecification** makes extrapolation fiction—fitting a line to a logarithmic truth systematically over-predicts at the extremes.
5. **Overfitting** makes memorisation look like learning—adding polynomial terms in study hours inflates training R^2 but destroys test R^2.
6. **Bad controls and endogeneity** destroy causal claims—controlling for confidence (X_3) does not clean the estimate, it collapses it.
7. **Real data** confronts you with all failures simultaneously—the tutoring example shows that every diagnostic can pass while the estimate points the wrong way.
8. **Sound research design** (RDD) earns regression its trust back—the scholarship cutoff identifies a causal effect without relying on correct model specification.

The unifying lesson is not that regression should be abandoned, but that its outputs must be accompanied by an honest assessment of the assumptions under which they are valid. The diagnostics tell you whether the *model* is internally consistent; only the *research design* can tell you whether the estimate is credible.

Key Insight

Every section in these notes followed the same arc: a toy story made you feel the problem, the theorem made it precise, a numerical example made it concrete, and a visualisation let you play with it. If you remember only one thing, remember this: regression is not a magic box that turns data into truth. It is a tool with assumptions, and every number it produces is only as honest as those assumptions. Your job is not to believe the output—it is to interrogate it.

Remember the seat number from the beginning? You now know it was "significant" only because n was huge and SE was tiny—not because seats matter. Remember the coffee study? You can sign the bias. Remember the "hot guys are jerks" article? You can draw the collider. Remember the ghost of self-discipline that haunted the studying coefficient from Section 3 onward? You have watched it inflate, distort, destroy, and flip the sign of every estimate it touched—until a clever research design finally exorcised it. You have seen regression fail eight different ways, and for each failure, you know the diagnosis, the repair, and the repair's limits.

The next time someone shows you a regression table and says "the model is fine, the R^2 is high, the residuals look clean," your first question should be: is this a Lalonde? Is there a ghost in this regression? Is this a seat number? These three questions—about confounding, about the research design, and about practical importance—will serve you longer than any formula in these notes.

This does not mean you should stop using regression.
It means you are finally ready to start.

A Proofs and Derivations

This appendix collects the formal proofs and derivations referenced throughout the main text.

A.1 Proof of the OVB Formula (Theorem 3.1)

Proof. Write the OLS estimator in population form. From the short regression,

$$\text{plim } \hat{\beta}_1^{\text{short}} = \frac{\text{Cov}(X_1, Y)}{\text{Var}(X_1)} = \frac{\text{Cov}(X_1,\, \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \varepsilon)}{\text{Var}(X_1)}.$$

Since $\mathbb{E}[\varepsilon \mid X_1, X_2] = 0$ implies $\text{Cov}(X_1, \varepsilon) = 0$, we obtain

$$\text{plim } \hat{\beta}_1^{\text{short}} = \beta_1 + \beta_2 \cdot \frac{\text{Cov}(X_1, X_2)}{\text{Var}(X_1)} = \beta_1 + \beta_2\, \delta_1. \qquad \square$$

A.2 Proof of Collider Bias (Proposition 3.1)

Proof. Since C is a function of X, U, and η, fixing C constrains the sum $\alpha_1 X + \alpha_2 U$. Knowing that $\alpha_1 X + \alpha_2 U \approx C - \eta$ means that a large X implies a small U (and vice versa). Formally, for a linear system with jointly normal variables:

$$\text{Cov}(X, U \mid C) = \text{Cov}(X, U) - \text{Cov}(X, C)\ \text{Var}(C)^{-1}\ \text{Cov}(C, U).$$

Since $\text{Cov}(X, U) = 0$ but $\text{Cov}(X, C) = \alpha_1 \text{Var}(X) \neq 0$ and $\text{Cov}(C, U) = \alpha_2 \text{Var}(U) \neq 0$:

$$\text{Cov}(X, U \mid C) = -\frac{\alpha_1 \text{Var}(X) \cdot \alpha_2 \text{Var}(U)}{\text{Var}(C)} \neq 0.$$

Conditioning on the collider manufactured a negative association between X and U from nothing. $\square$

A.3 Derivation of the Sandwich Estimator (Theorem 4.1)

Derivation of the sandwich estimator. Start from the OLS estimator: $\hat{\beta}_{\text{OLS}} = \beta + (X^\top X)^{-1} X^\top \varepsilon$. Therefore:

$$\begin{aligned}
\hat{\beta}_{\text{OLS}} - \beta &= (X^\top X)^{-1} X^\top \varepsilon, \\
\text{Var}(\hat{\beta}_{\text{OLS}} \mid X) &= (X^\top X)^{-1} X^\top\ \text{Var}(\varepsilon \mid X)\, X\, (X^\top X)^{-1} \\
&= (X^\top X)^{-1} X^\top \Omega\, X\, (X^\top X)^{-1}.
\end{aligned}$$

Under homoscedasticity, $\Omega = \sigma^2 I$, and this simplifies to $\sigma^2 (X^\top X)^{-1}$. Under heteroscedasticity, $\Omega = \text{diag}(\sigma_1^2, \ldots, \sigma_n^2)$, and the middle "meat" $X^\top \Omega\, X = \sum_{i=1}^n \sigma_i^2 x_i x_i^\top$ cannot be simplified.

The White–Huber–Eicker estimator replaces each unknown σ_i^2 with the squared residual $\hat{\varepsilon}_i^2$:

$$\widehat{X^\top \Omega\, X} = \sum_{i=1}^n \hat{\varepsilon}_i^2\, x_i x_i^\top.$$

This is consistent: as $n \to \infty$, the average $\frac{1}{n} \sum \hat{\varepsilon}_i^2 x_i x_i^\top$ converges to $\mathbb{E}[\sigma_i^2 x_i x_i^\top]$ by a law of large numbers, regardless of the form of heteroscedasticity. $\square$

A.4 Proof of the Bias–Variance Decomposition (Theorem 7.1)

Proof. Let $\mu = \mathbb{E}[\hat{f}(x_0)]$. Then

$$\begin{aligned}\mathbb{E}[(Y-\hat{f})^2] &= \mathbb{E}[(f+\varepsilon-\hat{f})^2] \\ &= \mathbb{E}[(f-\hat{f})^2] + \mathbb{E}[\varepsilon^2] + 2\mathbb{E}[\varepsilon(f-\hat{f})] \\ &= \mathbb{E}[(f-\hat{f})^2] + \sigma_\varepsilon^2,\end{aligned}$$

where the cross-term vanishes because ε is independent of $\hat{f}$ (trained on separate data) and has mean zero. Now expand:

$$\mathbb{E}[(f-\hat{f})^2] = (f-\mu)^2 + \mathbb{E}[(\hat{f}-\mu)^2] = \mathrm{Bias}^2(\hat{f}) + \mathrm{Var}(\hat{f}).$$ □

A.5 Proof of the Frisch–Waugh–Lovell Theorem (Theorem 11.1)

Proof. The OLS normal equations for the partitioned model $Y = X_1\beta_1 + X_2\beta_2 + \varepsilon$ give

$$\begin{pmatrix} X_1^\top X_1 & X_1^\top X_2 \\ X_2^\top X_1 & X_2^\top X_2 \end{pmatrix} \begin{pmatrix} \hat{\beta}_1 \\ \hat{\beta}_2 \end{pmatrix} = \begin{pmatrix} X_1^\top Y \\ X_2^\top Y \end{pmatrix}.$$

From the second block, $\hat{\beta}_2 = (X_2^\top X_2)^{-1} X_2^\top (Y - X_1\hat{\beta}_1)$. Substituting into the first block and simplifying:

$$\begin{aligned} X_1^\top X_1 \hat{\beta}_1 + X_1^\top X_2 (X_2^\top X_2)^{-1} X_2^\top (Y - X_1\hat{\beta}_1) &= X_1^\top Y \\ X_1^\top [I - X_2(X_2^\top X_2)^{-1} X_2^\top] X_1 \hat{\beta}_1 &= X_1^\top [I - X_2(X_2^\top X_2)^{-1} X_2^\top] Y \\ X_1^\top M_2 X_1 \hat{\beta}_1 &= X_1^\top M_2 Y. \end{aligned}$$

Hence $\hat{\beta}_1 = (X_1^\top M_2 X_1)^{-1} X_1^\top M_2 Y$. □