PAPER

Boost-range scaling of the fluctuations of the Benincasa–Dowker action and its consequences for Everpresent Λ

Carlos Echeverría-Arrondo

Instituto de Diseño para la Fabricación y Producción Automatizada, MALTA Consolider Team, Universitat Politècnica de València, 46022, València, Spain

E-mail: cecharr@idf.upv.es

Keywords: causal sets, Benincasa–Dowker action, cosmological constant, Everpresent Lambda, fluctuations

Abstract

Everpresent Λ is a causal-set inspired proposal in which the cosmological constant fluctuates as $\delta\Lambda \propto V^{-1/2}$, where V is the spacetime volume of the past of the observer, an exponent inherited from the Poisson fluctuations of the number of causal-set elements. Moradi, Yazdi and Zilhão asked whether these fluctuations could instead be sourced by the Benincasa–Dowker (BD) action, and gave a formalism for its variance without evaluating it. We evaluate it, analytically and with Poisson sprinklings in four dimensions. The leading term, the Poisson noise in the number of nearly null intervals, grows with the range of Lorentz boosts that fits inside the region and scales exactly as $(\rho V)^{2-2/d}$. In causal diamonds, truncated past light cones and past cones of spatially flat FLRW spacetimes reaching the big bang, the remaining correlation terms are a constant fraction of this term over the simulated range (ρV up to 12 800 for the diamond and 6400 for the cones), which bounds a residual power law in their ratio to $|\delta| \lesssim 0.1$. Extrapolating, the asymptotic exponent is $\Delta S \propto (\rho V)^{3/4}$, so that $\delta\Lambda \propto V^{-1/4}$, rather than the Poisson value; the geometry only fixes the amplitude, through a boost-range integral that we compute. For the smeared action the correlations do not grow faster than the leading term within a smaller range, and the physically relevant limit of strong smearing remains an extrapolation. Reading the action fluctuation as a fluctuation of the cosmological term, we build zero-mean stochastic $\Lambda(t)$ histories with this scaling, the measured temporal correlation and a free amplitude, and confront them with Pantheon+SH0ES supernovae, DESI DR2 baryon acoustic oscillations and Planck distance priors. None of 10^4 histories fits the data as well as ΛCDM ($\Delta\chi^2_{\rm min} = +124$ for the BD scaling with all three probes; +26 already without the CMB), and the exclusion is robust against the memory of the process and across a family of Ornstein–Uhlenbeck dynamics.

1 Introduction

Causal set theory replaces the spacetime continuum by a locally finite partially ordered set, whose elements are related by the causal order and whose number counts spacetime volume [1, 2]. A heuristic argument by Sorkin [3, 4], made precise in the Everpresent Λ model [5, 6, 7, 8], connects this discreteness with the cosmological constant. In unimodular gravity Λ is conjugate to the spacetime volume V; if V is only defined up to the Poisson fluctuation $\delta N = \sqrt{N}$ of the number of elements, then Λ fluctuates by $\delta\Lambda \sim V^{-1/2} \sim H^2$. The resulting dark energy density tracks the critical density at all epochs, which would explain why Ω_Λ is of order one today without fine tuning.

The exponent $-1/2$ is essential to the model. If $\delta\Lambda \propto V^{p-1}$ with $V \sim H^{-4}$, then $\delta\Lambda/H^2 \propto H^{2-4p}$, and only $p = 1/2$ gives an *everpresent* Ω_Λ. In the phenomenological realisation of Das, Nasiri and Yazdi (Model 1 of [7]), this exponent is imposed by construction: each causal-set element contributes an independent random variable to the action term ΛV. The authors note that "the true causal set mechanism that gives rise to Everpresent Λ is not fully understood" [7].

A natural candidate for a mechanism is the causal-set action itself. The Benincasa–Dowker (BD) action [9, 10] is a discretisation of the Einstein–Hilbert action, built from counts of intervals, whose mean reproduces the Einstein–Hilbert action in the continuum limit, up to boundary terms [11, 12]. Moradi, Yazdi and Zilhão (MYZ) [13] developed a formalism for the variance of this action under Poisson sprinkling and raised the question of whether its fluctuations could be the origin of Everpresent Λ. They expressed the variance as a set of integrals over pairs of intervals but

did not evaluate them. To our knowledge, the variance of the BD action and its scaling with the number of elements have not been computed since.

In this paper we compute them and follow the consequences through to cosmological data. Our main results are:

1. The *same-pair* term of the variance, the Poisson noise in the number of nearly null intervals, grows exactly as $(\rho V)^{2-2/d}$, with a prefactor set by a boost-range integral G of the region that we compute in closed form or by quadrature and verify up to $\rho V = 10^8$ (section 3).

2. In all the regions we simulate (causal diamonds, truncated past light cones and past cones of flat FLRW spacetimes reaching the big bang), the remaining correlation terms are a constant fraction of the same-pair term within the measured range ($\rho V \leq 12\,800$), for the local action and, within a smaller range of $\epsilon\rho V$, for the smeared one (section 4). The asymptotic exponent of ΔS is therefore consistent with 3/4 in four dimensions, not with the Poisson value 1/2.

3. If the action fluctuation is read as a fluctuation of the cosmological term, $\delta\Lambda \propto V^{-1/4}$ (section 5). We build stochastic $\Lambda(t)$ histories with this scaling and with the temporal correlation that we measure along nested past cones, and confront them with supernova, BAO and CMB data. They are excluded, as is Model 1 in our implementation, and the exclusion is robust against the dynamical assumptions we can vary (section 6).

The mechanism behind the first result is well known in another guise: the non-compactness of the Lorentz group makes the number of links, or of intervals of any fixed small cardinality, grow without bound in a sprinkled causal set [14], and the mean abundances of small intervals in flat regions are known exactly [15]. The same non-compact rapidity range biases the statistics of links [16]. What is new here is its role in the *fluctuations* of the action: we show that it controls their scaling, quantify the amplitude through G in the regions relevant for cosmology, and follow the consequences for a fluctuating cosmological constant.

2 The BD action and its variance

2.1 Sprinklings and the action

A Poisson sprinkling of density ρ into a region M of a d-dimensional Lorentzian manifold produces a causal set whose number of elements N is a Poisson variable with mean ρV, $V = \mathrm{vol}(M)$. For a related pair $y \prec x$ let $n(y,x)$ be the number of elements strictly between them, and let N_k be the number of pairs with $n = k$. We use the BD action in the form of a sum over related pairs,

$$S = a_N N + \sum_{y \prec x} f\big(n(y,x)\big)\,, \tag{1}$$

which is the form analysed by MYZ [13]. For the *local* action, $f(n) = a_n$ for $n < n_d$ and $f = 0$ otherwise, with

$$d=2: \quad (a_N; a_0, a_1, a_2) = (1; -2, 4, -2), \qquad d=4: \quad (a_N; a_0, \ldots, a_3) = \frac{4}{\sqrt{6}}(1; -1, 9, -16, 8), \tag{2}$$

in four dimensions these are the coefficients of [9], normalised so that the mean action tends to $\frac{1}{2\ell^2}\int R\,\mathrm{d}V$ (plus boundary terms [11, 12]), with $\ell = \rho^{-1/4}$; MYZ use a different global constant. In two dimensions they are those of [17], for which the mean action of a flat causal diamond tends to 1, its Euler characteristic. The *smeared* action of Sorkin [18] and Dowker and Glaser [10] introduces a non-locality parameter $\epsilon \in (0,1]$, with $\epsilon = (\ell/\ell_k)^d$ for a non-locality length ℓ_k. It has the same form (1) with

$$a_N = \epsilon^{2/d} a_N^{(1)}, \qquad f_\epsilon(n) = \epsilon^{1+2/d} \sum_{k<n_d} a_k^{(1)} \binom{n}{k} \epsilon^k (1-\epsilon)^{n-k}, \tag{3}$$

where $a^{(1)}$ are the local coefficients (2); the overall factor $\epsilon^{2/d}$ comes from $\ell_\epsilon^2 = \ell^2/\epsilon^{2/d}$, and the additional ϵ in f_ϵ from the normalisation of the smeared operator [10]. For $\epsilon = 1$ the local action is recovered. The binomial weight is the probability that a pair with n elements in between keeps exactly k of them after thinning each with probability ϵ, so the smeared action can be written with "thinned" abundances $\tilde{N}_k = \sum_{y\prec x} \binom{n}{k}\epsilon^k(1-\epsilon)^{n-k}$, and the MYZ formalism applies to them with minor modifications (appendix B).

2.2 Means and the Mecke formula

All the mean values we need follow from the Mecke formula for Poisson processes [19]. For any function h,

$$\Big\langle \sum_{y\prec x} h\big(n(y,x)\big)\Big\rangle = \rho^2 \int_M \mathrm{d}V_x \int_{M\cap J^-(x)} \mathrm{d}V_y\, \mathbb{E}\, h\big(\mathrm{Poisson}(\rho\,\mathrm{vol}\, I(y,x))\big), \tag{4}$$

where $I(y,x)$ is the causal interval. In a flat causal diamond the inner integral depends on x only through the volume of its past within the diamond, and (4) reduces to two nested one-dimensional quadratures. We also need the covariance of N with the pair sum $F=\sum_{y\prec x} f(n)$. Differentiating (4) with respect to the intensity gives (appendix A)

$$\mathrm{Cov}(N,F) = \rho\,\frac{\mathrm{d}\langle F\rangle}{\mathrm{d}\rho}\,, \tag{5}$$

which for the local action reduces to the identity $\mathrm{Cov}(N,N_k)=(k+2)L_k-(k+1)L_{k+1}$ of MYZ, with $L_k=\langle N_k\rangle$.

2.3 The MYZ decomposition

The variance of (1) is

$$(\Delta S)^2 = a_N^2\,\rho V + 2a_N\,\mathrm{Cov}(N,F) + \mathrm{Var}(F)\,, \tag{6}$$

and MYZ decompose $\mathrm{Var}(F)$ according to how the two pairs (y_1,x_1), (y_2,x_2) in F^2 share their endpoints:

$$\mathrm{Var}(F) = J^1 + J^2 + J^3 + 2J^4 + J^6_{\mathrm{c}}\,. \tag{7}$$

Here $J^1=\langle\sum_{y\prec x} f(n)^2\rangle$ is the *same-pair* term; J^2 and J^3 collect distinct pairs sharing their past or future endpoint; J^4 collects chains $y_1\prec x_1=y_2\prec x_2$ (MYZ's J^5, the chain with the opposite orientation, equals J^4 by symmetry and is absorbed in the factor 2); and J^6_{c} is the connected part of the contribution of pairs with four distinct endpoints, which is non-zero only when the two intervals overlap. MYZ write all terms as integrals over configurations of points, with the joint distribution of the two interval cardinalities given by the volume of the intersection of the two intervals.

We evaluated the MYZ integrals by Monte Carlo with an importance sampler adapted to nearly null intervals. In four dimensions the intersection volume of two causal intervals has no closed form; we computed it by quadrature, using Green's theorem for the intersection of discs at fixed time and fixed coordinate along one axis. In two dimensions the prediction agrees with direct sprinklings at five densities $\rho V = 25$–400 ($|z|\le 1.56$, 4000 sprinklings each) and, for the smeared action with $\epsilon = 0.5, 0.25$, at ten more points ($|z|\le 1.72$). In four dimensions the agreement holds for $\rho V = 25, 50, 100$, but the Monte Carlo error grows quickly with density (23% at $\rho V = 100$). This makes the full MYZ integrals impractical as a scaling tool, and in the rest of the paper we follow a different strategy: we compute the terms that dominate asymptotically *exactly*, and we measure the remaining terms on sprinklings.

3 The same-pair term and the boost range

Consider a nearly null pair, with small proper time τ between its endpoints. Its interval has volume $c_d\tau^d$ ($c_4=\pi/24$), so its cardinality is a Poisson variable of mean $\mu=\rho c_d\tau^d$ that stays of order one however large the region is. What grows with the size L of the region is the number of such pairs. Writing $y = x - \tau(\cosh\chi, \sinh\chi\,\hat{\boldsymbol{n}})$, the measure of the past hyperboloid is $\tau^{d-1}\sinh^{d-2}\chi\,\mathrm{d}\tau\,\mathrm{d}\chi\,\mathrm{d}\Omega$, and the pair fits inside the region as long as its rapidity satisfies $\tau\mathrm{e}^{\chi}/2 \lesssim \ell(x,\hat{\boldsymbol{n}})$, where $\ell(x,\hat{\boldsymbol{n}})$ is the extent, in time, of the past null ray from x in the direction $\hat{\boldsymbol{n}}$ that remains inside the region. The rapidity integral then gives a factor $(\ell/\tau)^{d-2}$, which is the volume of the available range of boosts. Inserting this in (4) and changing variables to μ gives, for $d=4$,

$$\Big\langle \sum_{y\prec x} h(n)\Big\rangle \simeq (\rho V)^{3/2}\,\frac{G/V^{3/2}}{8\sqrt{c_4}}\sum_{n\ge 0} h(n)\,\frac{\Gamma(n+\frac12)}{n!}\,, \qquad G=\int_M \mathrm{d}V_x \int_{S^2}\mathrm{d}\Omega\;\ell(x,\hat{\boldsymbol{n}})^2\,, \tag{8}$$

and in general dimension the exponent is $2-2/d$, with a weight $\Gamma(n+2/d)/n!$. Two special cases matter.

Two dimensions. For $d=2$ the boost volume is $\ln(\ell/\tau)$ and the weight is $\Gamma(n+1)/n!=1$, so that $\langle\sum h\rangle = \rho V\ln(\rho V)\sum_n h(n) + O(\rho V)$. The variance is Poissonian up to a logarithm, $(\Delta S)^2 \simeq b\,\rho V\ln\rho V$ with $b=\sum_n f(n)^2$ ($b=24$ for the local action).

The cross term. For $h(n) = f(n)$ the leading term of (8) vanishes. For the local action in $d = 4$, $\sum_k a_k \Gamma(k + \frac{1}{2})/(k!\sqrt{\pi}) \propto -1 + \frac{9}{2} - 6 + \frac{5}{2} = 0$. This is the same cancellation that makes the mean action finite in flat space, and by (5) it implies that $\mathrm{Cov}(N, F)$ grows only like ρV.

Therefore, if the correlation terms in (7) do not grow faster than J^1, the variance is dominated by

$$J^1 \propto (\rho V)^{2-2/d} \sum_n f(n)^2 \, \frac{\Gamma(n + 2/d)}{n!} \quad \Longrightarrow \quad \Delta S \propto (\rho V)^{1-1/d}, \tag{9}$$

that is, $\Delta S \propto (\rho V)^{3/4}$ in four dimensions. The region enters only through G, the integrated squared length of the past null rays: the boost range available to nearly null intervals. The exponent is a direct consequence of Lorentz invariance together with a cut-off set by the size of the region, the same non-compactness that makes the number of links unbounded in a sprinkling [14]; it cannot be changed by a finite rescaling of the density.

Why the correlation terms should follow J^1. We have no proof that the other terms in (7) grow no faster than J^1, but there are two heuristic reasons to expect it. First, J^2, J^3 and J^4 are sums of products of the row and column sums of the matrix of pair weights (appendix B), e.g. $\sum_x f(y, x)$ for a fixed y. The boost-enhanced part of such a sum is proportional to $\sum_n f(n)\Gamma(n + \frac{1}{2})/n!$, which vanishes by the cancellation just described, so these terms cannot acquire a second factor of the boost range. Second, J^6_c only receives contributions from pairs of intervals that overlap. Two nearly null intervals overlap appreciably only if they lie along nearly the same null direction with comparable rapidity, so for each interval the measure of correlated partners is finite, and J^6_c inherits the boost range once, like J^1. In two dimensions this is borne out: the connected term grows with the same logarithm as J^1. The numerical tests of the next section probe this expectation directly.

The approach to the asymptotic regime is slow. The exact J^1 of the local action in the four-dimensional diamond, computed from (4) without Monte Carlo, has a local exponent $\frac{1}{2}\,\mathrm{d}\ln J^1/\mathrm{d}\ln \rho V$ that decreases from 1.26 at $\rho V \approx 35$ to 0.96 at $\rho V \approx 570$, 0.88 at $\rho V \approx 2300$ and 0.77 at $\rho V \approx 3 \times 10^5$. Exponents measured at small densities, such as the $\Delta S \propto (\rho V)^{1.2}$ we find for $\rho V \leq 100$, are therefore pre-asymptotic.

4 Numerical tests

4.1 Method

For each region we compute J^1, $\langle F \rangle$ and $\mathrm{Cov}(N, F)$ exactly from (4) and (5), measure the total variance on independent Poisson sprinklings, and define the correlation *remainder*

$$r = \frac{\mathrm{Var}(F) - J^1}{J^1} = \frac{J^2 + J^3 + 2J^4 + J^6_c}{J^1}, \tag{10}$$

the relative size of all correlation terms. The asymptotic exponent (9) holds if r tends to a constant. Our criterion is that the slope of a weighted linear fit of r against $\ln \rho V$ be compatible with zero at 4σ. It was fixed before the corresponding simulations, except for the causal diamond, where it was fixed after a first set of five densities and before the larger ones. The point at $\rho V = 12\,800$, the FLRW cones at $\rho V = 6400$ and the tests described as additional below were added, with their criteria fixed in advance, after an independent review of a first draft. A slope compatible with zero is a weak statement on its own, so we also quote the bound it implies on a residual power law, $r \propto (\rho V)^{\delta}$: to first order the slope equals $\delta\,\bar{r}$. Such a law would change the asymptotic exponent to $p = 3/4 + \delta/2$ once the correlations dominate. Each sprinkling is an exact, independent sample of the ensemble, so no thermalisation is involved. The statistical error of a variance estimated from M sprinklings is $\mathrm{std}(d^2)/\sqrt{M}$, with $d = S - \bar{S}$.

4.2 Causal diamond

Figure 1 shows the four-dimensional diamond with the local action, for $\rho V = 25$ to 6400 (500 to 6000 sprinklings per density). The total variance follows the exact J^1 with a constant offset, and the local exponent of the total variance decreases with ρV in step with that of J^1, from 1.24 ± 0.04 to 0.72 ± 0.05; on these sprinklings alone, $r = 0.317 \pm 0.014$ with slope $+0.006 \pm 0.009$. As an additional test we combined them with the independent sprinklings of the smeared-action runs below (at $\epsilon = 1$, $\rho V = 400$–6400) and with 500 new sprinklings at $\rho V = 12\,800$. The remainder at the largest density is $r = 0.38 \pm 0.10$, compatible with the earlier mean (0.6σ), and over the whole range $r = 0.324 \pm 0.012$, i.e. $(\Delta S)^2 \simeq 1.32\,J^1$, with slope $+0.007 \pm 0.008$ (1.0σ, $\chi^2 = 12.4$ for 8




Figure 1. Local BD action in a four-dimensional causal diamond. (a) Total variance per element from sprinklings (points) and the exact same-pair term J^1 (line); the dotted line has the asymptotic slope of J^1, $p = 3/4$; the square is the independent set at $\rho V = 12\,800$. (b) Local exponent of $\Delta S \propto (\rho V)^p$ for J^1 (line, exact) and for the total variance (points; the square uses the point at 12 800). The total follows J^1 towards 0.75; the Poisson value 0.5 is not approached. (c) Remainder $r = (\mathrm{Var}\, F - J^1)/J^1$ combining all independent sprinklings, including $\rho V = 12\,800$; the band is the weighted mean. The shading in (a) and (b) marks the simulated range; beyond it only J^1 is computed.

degrees of freedom; figure 1(c)). The implied bound on a residual power law is $-0.02 < \delta < 0.07$ at 2σ ($-0.07 < \delta < 0.12$ at 4σ), i.e. $p \leq 0.78$ (0.81). Two different but unbiased estimators of $\mathrm{Var}\, F$ are combined here: for the first set, the variance of the full action minus the exact N contributions; for the others, the variance of F itself. Since the per-density errors, $\mathrm{std}(d^2)/\sqrt{M}$, could be underestimated for a heavy-tailed d^2, we checked them a posteriori with a bootstrap over sprinklings wherever the individual sprinklings were stored: the two errors agree within 3% in all 39 sets. The bound is also stable against the fitted range and the scatter of the points: scaling the slope error by $\sqrt{\chi^2/\mathrm{dof}}$ gives $\delta < 0.08$, and restricting the fit to $\rho V \geq 400$ gives $\delta < 0.09$ (2σ; 0.15 with the scaled error, the scatter being larger in this subset, $\chi^2 = 10.6$ for 4 degrees of freedom). An exponent as low as the Poisson value would require $r \to -1$, correlations that cancel J^1; we see no trend in that direction.

In two dimensions (diamond, ρV up to 12 800) the logarithmic law of section 3 holds: the slope of $(\Delta S)^2/\rho V$ against $\ln \rho V$ is 24.43 ± 0.24, against $b = 24$ from J^1 alone. Binning the pairs that contribute to $\sum f^2$ by their rapidity $\eta = \frac{1}{2}\ln(\Delta u/\Delta v)$ gives a plateau for $|\eta| \lesssim 3$, cut off at $|\eta| \approx \frac{1}{2}\ln \rho V$: every unit of rapidity contributes the same, which identifies the logarithm with the boost range rather than with the boundary.

4.3 Past light cones and FLRW

Everpresent Λ involves the causal past of an observer, not a diamond. We therefore repeated the analysis for the truncated past light cone $J^-(q) \cap \{t \geq 0\}$ in Minkowski spacetime and for the past light cone of an observer in a spatially flat FLRW spacetime, $\mathrm{d}s^2 = a(\eta)^2(-\mathrm{d}\eta^2 + \mathrm{d}\boldsymbol{x}^2)$, reaching the big bang at $\eta = 0$. We take $a = \eta^\alpha$ with $\alpha = 1$ (radiation) and $\alpha = 2$ (matter). In FLRW the interval volumes are computed exactly: at fixed η the section of a causal interval is a lens (the intersection of two balls), and the integral over η is done exactly by Gauss–Legendre quadrature in each polynomial piece. The boost-range integral of (8) becomes

$$G = \int \mathrm{d}V_x \int \mathrm{d}\Omega \int_0^{\eta_x} \frac{2\lambda\, a(\eta_x - \lambda)^4}{\sqrt{w}}\, \mathrm{d}\lambda, \qquad w = \frac{6}{\lambda^3}\int_0^\lambda a(\eta_y + s)^4\, s(\lambda - s)\, \mathrm{d}s, \tag{11}$$

where λ is the conformal extent of a nearly null interval along the null ray and w is the average of a^4 over it, weighted by the profile $s(\lambda - s)$ of a thin four-dimensional diamond. For $a = 1$ it reduces to the Minkowski cone.

Table 1 summarises the results. In all regions the exact means converge to the asymptotic form (8), with deviations below 2.2% at $\rho V = 10^8$ that decrease monotonically with density. For instance, the ratio of J^1 in the cone and in the diamond rises from 0.516 at $\rho V = 1.5 \times 10^5$ to 0.562 at 10^8, towards the predicted $0.8187/1.4472 = 0.566$. The remainder is a constant fraction of J^1 in all four regions, with values between 0.32 and 0.35; its slope against $\ln \rho V$ is compatible with zero in all cases (largest deviation 1.0σ). The FLRW points at $\rho V = 6400$, an additional test, give $r = 0.28 \pm 0.08$ ($\alpha = 1$) and 0.37 ± 0.09 ($\alpha = 2$), within 0.8σ of the earlier means. The two dimensional analogues behave in the same way: in the Minkowski cone and in FLRW cones with

Table 1. Region dependence of the fluctuations of the local BD action in four dimensions. $G/V^{3/2}$ is the boost-range integral (8); the asymptotic ratio $\langle N_k\rangle$/(prediction) is evaluated with the exact means at $\rho V = 10^8$ for $k = 0, \ldots, 3$. The remainder r is the weighted mean over the simulated densities, and the slope is that of r against $\ln \rho V$.

Region	$G/V^{3/2}$	ratio at 10^8	simulated ρV	r	slope of r
Diamond	1.4472 ± 0.0005	0.995–0.987	25–12 800	0.324 ± 0.012	$+0.007 \pm 0.008$
Minkowski cone	0.8187	0.992–0.978	25–6400	0.342 ± 0.017	-0.001 ± 0.011
FLRW, $\alpha = 1$	1.3145	0.995–0.986	25–6400	0.346 ± 0.016	$+0.002 \pm 0.010$
FLRW, $\alpha = 2$	1.4194	0.997–0.981	25–6400	0.319 ± 0.015	-0.004 ± 0.010




Figure 2. Decomposition of the correlation terms of the smeared BD action in a four-dimensional diamond, relative to the exact same-pair term J^1, for three values of the non-locality parameter. The terms with a shared endpoint (J^2, J^3, $2J^4$) decrease with ρV and the connected overlap term J^6_c increases; their sum, the remainder r (black), stays approximately constant. The figure shows the second of the two independent sets of sprinklings described in the text. Points of different terms are shifted horizontally by a few per cent for clarity.

$\alpha = 1, 2$ the coefficient of $\rho V \ln \rho V$ is 24.09 ± 0.19, 24.26 ± 0.19 and 24.78 ± 0.24. The big bang, where $a \to 0$, does not change the scaling. The exponent is universal, and the geometry only sets the amplitude through G.

4.4 Smeared action

The smeared action is designed to suppress fluctuations, and it is the version that can give a phenomenologically relevant amplitude (section 5). For fixed ϵ the argument of section 3 applies unchanged, with $J^1_\epsilon \propto (\rho V)^{3/2} \sum_n f_\epsilon(n)^2 \Gamma(n + \frac{1}{2})/n!$, and J^1_ϵ is again computed exactly from (4). Whether the correlation terms remain proportional to it is less clear, because the smeared action receives contributions from intervals with up to $\sim 1/\epsilon$ elements, so the asymptotic regime requires $\epsilon \rho V \gg 1$.

We sprinkled the four-dimensional diamond at $\rho V = 400$–6400 (2000 to 500 sprinklings per density) and analysed the same sprinklings with $\epsilon = 1, 0.5, 0.2, 0.1$; the decomposition described below uses a second, independent set of sprinklings of the same sizes. The exact means and covariances (5) agree with the simulation in all 40 comparisons ($|z| \leq 2.35$), and $\epsilon = 1$ reproduces the local result, $r = 0.345 \pm 0.025$.

Our pre-registered criterion for the smeared action was formulated for the remainder *after* subtracting the part of F linear in N (below), $r_\perp = r - \rho V g^2 / J^1$. It was met for $\epsilon = 0.5$ (slope 0.9σ) but failed for $\epsilon = 0.2$ and 0.1, where $r_\perp$ grows with $\ln \rho V$ at 4.1σ and 6.7σ. The analysis that follows, which explains this failure, was designed after seeing it.

For $\epsilon \leq 0.2$ two contributions of different origin compete at these densities. The part of F that is linear in N, of variance $\rho V\, g^2$ with $g = \mathrm{d}\langle F\rangle/\mathrm{d}(\rho V)$, is Poissonian, so it decays relative to J^1 as $(\rho V)^{-1/2}$; for $\epsilon = 0.1$ it falls from $2.75\, J^1$ at $\rho V = 400$ to $0.55\, J^1$ at 6400. To separate the contributions we measured every MYZ term directly on each sprinkling, using estimators built from the matrix $f(y, x)$ of pair weights and its row and column sums (appendix B). The generic MYZ integrator is not usable in this regime: at $\rho V = 6400$ its error on J^6_c was about ten times the signal even with 4×10^5 samples.

Figure 2 shows the result. As a check, the estimated J^1 agrees with the exact value in all fifteen cases ($|z| \leq 0.86$), and at $\rho V = 400$, $\epsilon = 0.1$, the estimated $J^4 = 17.86 \pm 0.06$ agrees with the independent MYZ Monte Carlo value 19.4 ± 2.1. For $\epsilon = 0.5$ all terms are flat, with $r \simeq 1.3$. For

$\epsilon = 0.2$ and 0.1 the terms with a shared endpoint, which carry the Poissonian part, decrease relative to J^1, while J^6_c increases (from $-3.7\,J^1$ to $-0.1\,J^1$ for $\epsilon = 0.1$). The two trends compensate: the total remainder stays at $r \approx 3.0$ ($\epsilon = 0.2$) and $r \approx 4.5$ ($\epsilon = 0.1$), with slopes against $\ln \rho V$ of $+0.17 \pm 0.08$ and -0.29 ± 0.11, both below 3σ. The growth of $r_\perp$ that made the pre-registered criterion fail is therefore not a growing correlation: it is the approach of $r_\perp$ to r as the subtracted Poissonian term decays.

Since this explanation was found a posteriori, we tested it as an additional pre-registered criterion: the slope of the unprojected r, combining both sets of sprinklings with the 500 new ones at $\rho V = 12\,800$ (which doubles $\epsilon\rho V$), must be compatible with zero at 4σ. At $\rho V = 12\,800$ we find $r = 1.23 \pm 0.13$, 3.03 ± 0.26 and 4.38 ± 0.32 for $\epsilon = 0.5$, 0.2 and 0.1, and the slopes over $\rho V = 400$–$12\,800$ are $+0.02 \pm 0.03$ (0.8σ), $+0.11 \pm 0.05$ (2.3σ) and -0.25 ± 0.07 (-3.7σ, $\chi^2 = 9.4$ for 4 degrees of freedom). The criterion is met, though not by a wide margin for $\epsilon = 0.1$, where r decreases. This trend comes from the point at $\rho V = 400$, where $\epsilon\rho V = 40$ is clearly pre-asymptotic; for $\rho V \geq 800$ the slope is -0.09 ± 0.09. In any case, the exponent only requires the one-sided statement that r does not grow: a decreasing r means that the correlations grow more slowly than J^1. The upper bounds on δ at 2σ are 0.06, 0.07 and -0.02 (with bootstrap errors, which agree with the formula, the same).

Within the accessible range, $\epsilon\rho V \leq 1280$, the smeared action thus behaves like the local one: the correlations do not grow faster than J^1, and $\Delta S \propto (\rho V)^{3/4}$. The individual terms are still drifting, so we cannot exclude a different behaviour for $\epsilon \to 0$ at fixed $\epsilon\rho V$; we return to this point in section 7.

5 Consequences for a fluctuating Λ

5.1 Amplitude

To connect the action with Λ we follow the logic of Everpresent Λ. The BD action approximates the Einstein–Hilbert action, $S_{\rm EH}/\hbar = \kappa S$ with $\kappa = \ell^2/(8\pi\ell_p^2)$ in our normalisation, and the cosmological term is $\Lambda V/(8\pi\ell_p^2)$ [7]. Reading the action fluctuation in the past of the observer as a fluctuation of the cosmological term gives

$$\delta\Lambda\,\ell^2 = \frac{\Delta S}{\rho V}\,, \tag{12}$$

independently of ℓ_p. Applied to the Poisson action of Model 1 this reproduces its $\delta\Lambda = 8\pi\alpha/\sqrt{V}$ exactly. This only checks the normalisation κ, since Model 1 defines its fluctuating term through the same relation; equation (12) itself is a working hypothesis, not a consequence of a causal-set dynamics. Note also that we use only the *fluctuation* of the action: its mean in a region with a boundary contains boundary and corner terms [11, 12], which play no role here. With the results of section 4,

$$\delta\Lambda\,\ell^2 = A\,(\rho V)^{-1/4}, \qquad A^2 = (1+r)\,\frac{G/V^{3/2}}{8\sqrt{c_4}}\sum_n f(n)^2\frac{\Gamma(n+\frac{1}{2})}{n!}\,. \tag{13}$$

For the present past cone of a Planck 2018 ΛCDM background [20], $V \approx 6.4\times10^{242}\,\ell_p^4$. With the local action and $\ell = \ell_p$ this gives $\delta\Lambda/\Lambda_{\rm obs} \sim 10^{62}$; matching the observed value would require $\ell \sim 10^{27}$ m, larger than the Hubble radius. The local action is therefore excluded as a source.

For the smeared action we find numerically that $\sum_n f_\epsilon(n)^2\Gamma(n+\frac{1}{2})/n! \to 4.112\,\epsilon^{5/2}$ as $\epsilon \to 0$, so $A \propto \epsilon^{5/4} = (\ell/\ell_k)^5$. The observed Λ is obtained for $\ell_k \approx 2.6\times10^{-23}$ m with $\ell = \ell_p$ ($\epsilon \sim 10^{-49}$), or $\ell_k \approx 1.25\times10^{-22}$ m with $\ell = \ell_{\rm cs} = 7.1\,\ell_p$, the discreteness length that corresponds to $\alpha = 0.01$ in Model 1. These values use $r = 0.33$; since $\ell_k \propto (1+r)^{1/10}$, the larger $r \approx 3$–4.5 found above for small ϵ raises them by at most 15%. This is below the bound $\ell_k \lesssim 10^{-19}$ m derived from 8 TeV LHC data, and inside the range, from 2×10^{-22} to 10^{-29} m, that proposed optomechanical experiments could probe [21]. It requires, however, a large hierarchy $\ell_k/\ell_p \sim 10^{12}$, so the coincidence problem that Everpresent Λ was meant to solve reappears as a tuning of ℓ_k. Equation (13) also changes the *time* dependence: since $\delta\Lambda \propto V^{-1/4} \propto H$ in the matter era, the envelope $\delta\Lambda/3H^2$ behaves as a fluid with $w \approx -1/2$, negligible at recombination ($\sim 4\times10^{-5}$ at $z = 1100$) and growing until today, whereas for Model 1 $w \approx 0$.

5.2 Temporal correlation along the worldline

A stochastic $\Lambda(t)$ also requires the correlation of ΔS between the past cones of two instants of the same worldline. These cones are nested, and since the region is causally convex every pair of the smaller cone is a pair of the larger one with the same cardinality. A single sprinkling of the largest cone therefore gives the action of all the nested cones. We measured it for the local action in four

dimensions, in the Minkowski cone and in the matter-dominated FLRW cone (4000 sprinklings each, $\rho V = 200$, 500, 1250 and 3200). The action is close to a random walk, with a small positive memory: the correlation between the action and its increment is $\kappa = 0.069 \pm 0.015$ (weighted mean of the six pairs of adjacent cones; no individual pair reaches 4σ). We parametrise the correlation between the actions S_i, S_j of two nested cones, $V_i < V_j$, as

$$\operatorname{corr}(S_i, S_j) = \left(\frac{\operatorname{Var} S_i}{\operatorname{Var} S_j} \right)^{\beta/2}, \tag{14}$$

where $\beta = 1$ is a random walk with independent increments, and find $\beta = 0.88$–0.97 over all pairs of cones. A pre-registered requirement that $|\kappa| < 0.2$ for every pair of cones failed for one non-adjacent pair in each geometry ($\kappa = 0.29 \pm 0.12$ and 0.23 ± 0.09), a criterion that, in retrospect, ignored their errors; it does not affect the value of β.

6 Confronting the data

6.1 Stochastic model

We model the cosmological constant along the observer's worldline as

$$\Lambda(t) = \lambda\, V(t)^{-q}\, s(t), \tag{15}$$

where $V(t)$ is the volume of the causal past and s is an Ornstein–Uhlenbeck process [22] in $\ln V$ with unit variance and $\operatorname{corr}(s_1, s_2) = (V_1/V_2)^c$. Model 1 of [7] corresponds to $q = c = 1/2$ and $\lambda = 8\pi\alpha$. The BD action, with (12), corresponds to $q = 1/4$ and $c = 3\beta/4$, and we take $\beta = 0.9$ from the measurement above. The background is integrated with feedback, $(H/H_{100})^2 = \omega_r a^{-4} + \omega_m a^{-3} + \Lambda/3$, with V accumulated along the way. We set $\Lambda = 0$ before $z = 10^4$, start the process from zero and count realisations with $H^2 < 0$ as collapsed. This is the convention that reproduces the completion fractions published for Model 1 (10 000, 3258 and 575 out of 10 000 for $\alpha = 0.005$, 0.01 and 0.012, against 9990, 3228 and 535 in [7]).

We follow the per-seed method of [8]. Each seed fixes the random numbers of the process, and for each seed we minimise χ^2 over ω_m, the amplitude (α or λ) and the supernova magnitude M_B. A seed is *good* if $\chi^2 < \chi^2_{\Lambda\mathrm{CDM}} + 14.7$. The margin is that implied by [8], who count seeds with $\chi^2 < 1500$ against their ΛCDM value of 1485.3 for supernovae alone. Our own supernova-only ΛCDM fit gives 1452.0, with best-fit parameters that agree with [23]; we could not reconstruct the exact configuration behind 1485.3, so we always apply the margin relative to our own ΛCDM fit to the same data. The data are:

- Pantheon+SH0ES supernovae [24, 23, 25], with the standard cut (1657 light curves, 77 of them Cepheid calibrators) and the full statistical and systematic covariance;
- DESI DR2 BAO [26]: 13 measurements of D_V/r_d, D_M/r_d and D_H/r_d at 7 redshifts, with their covariance;
- Planck 2018 distance priors [27] on R and l_A, with $\omega_b = 0.02236$, the sound horizon integrated over each history, and z_*, z_{drag} from the fits of [28, 29] anchored to their Planck values.

The pipeline reproduces the Planck 2018 sound horizons to 0.08% and R, l_A within 1σ, and the DESI value of Ω_m from BAO alone (0.2972 against 0.2975 ± 0.0086). The treatment of radiation is simplified (no massive neutrinos, fixed ω_r), which induces a systematic of order 0.1–0.2% in l_A. A variant in which l_A and r_d are rescaled so that the Planck best fit reproduces their published values exactly gives the same conclusions. Distance priors assume standard early physics. This holds for the BD model, where $\Lambda/3H^2 \sim 4 \times 10^{-5}$ at $z = 1100$, but not for Model 1 (~ 0.5), which we therefore fit to supernovae and BAO only. Its BAO term still uses the sound horizon integrated over each history, which stretches the fitting formulae for z_{drag}; the conclusion does not depend on it, since the supernova part of its best seed alone is already +46.

6.2 Results

With supernovae alone both models produce rare histories that fit as well as ΛCDM: 0.25% of good seeds for Model 1, consistent with the $\approx 0.3\%$ of [8], and 0.48% for the BD action. The good BD seeds fix $\ell_k \approx 3 \times 10^{-23}$ m, consistent with the value inferred from the amplitude alone. Both, however, prefer $\Omega_m \approx 0.5$.

Adding BAO changes the picture (table 2). Already without the CMB, none of the 10^4 seeds of either model is good (fraction below 3×10^{-4} at 95% confidence): $\Delta\chi^2_{\mathrm{min}} = +87$ for Model 1 and

Table 2. Fits of stochastic $\Lambda(t)$ histories (10 000 seeds per row). $\Delta\chi^2$ is relative to the ΛCDM fit to the same data ($\chi^2 = 1488.2$ for SN + BAO, 1504.0 for SN + BAO + CMB). The last column splits the $\Delta\chi^2$ of the best seed by probe. Model 1 is shown without the CMB, since the distance priors are not applicable to it.

Model	good seeds	$\Delta\chi^2_{\rm min}$	best seed: SN / BAO / CMB
SN + BAO			
Model 1 ($q = c = \frac{1}{2}$)	0	+87	+46 / +40 / –
BD, $\beta = 0.9$	0	+26	−13 / +39 / –
SN + BAO + CMB			
BD, $\beta = 0.5$	0	+45	+40 / +4 / +1
BD, $\beta = 0.88$	0	+116	+47 / +69 / +0.5
BD, $\beta = 0.9$	0	+124	+37 / +82 / +4.5
BD, $\beta = 0.97$	0	+135	+50 / +79 / +6
BD, $\beta = 1$	0	+154	+56 / +79 / +20




Figure 3. Robustness of the exclusion. Left: minimum $\Delta\chi^2$ with respect to ΛCDM for the BD dynamics ($q = 1/4$) as a function of the memory parameter β (10 000 seeds each); the dashed line is the threshold for a good seed and the band marks the measured range of β. Right: minimum $\Delta\chi^2$ over the (q, c) family of Ornstein–Uhlenbeck dynamics (1000 seeds per cell). Model 1 is the cell $q = c = 1/2$. The diamond marks the BD action ($q = 1/4$, $c = 3\beta/4$ with $\beta = 0.9$) and the bar the measured range of β. With 1000 seeds per cell the minima are somewhat higher than with the 10^4 seeds of the left panel (e.g. +182 at $c = 0.75$ against +154 for $\beta = 1$).

+26 for the BD action, with ΛCDM refitted to SN + BAO. For Model 1 this agrees with the full Planck analysis of [8], which found $\Delta\chi^2 > 150$, here by an independent route that does not use the CMB. The best BD seed fits the supernovae better than ΛCDM (−13) but misses the BAO by +39: no history achieves simultaneously the shape of the distance–redshift relation required by the supernovae and the scale required by the BAO. Adding the CMB, which is applicable to the BD model because its Λ is negligible at recombination, raises the minimum to +124 (with $\beta = 0.9$), and none of the seeds that were good with supernovae alone survives.

Figure 3 tests the robustness of this conclusion. Over the measured range of the memory parameter, $\beta = 0.88$–0.97, the minimum $\Delta\chi^2$ stays between +116 and +135, and even an unrealistically long memory ($\beta = 0.5$) remains above the threshold. Over the whole family $(q, c) \in [0, \frac{1}{2}] \times [\frac{1}{4}, 1]$ no cell has good seeds. The best cell, $q = 1/8$, $c = 1/4$, reaches +17 with 1000 seeds; since this is close to the threshold, we extended it to 10^4 seeds (which include the original 1000) as an additional test: no seed is good, and the minimum is unchanged, $\Delta\chi^2_{\rm min} = +17.0$. The data only tolerate a Λ that depends weakly on the volume and decorrelates slowly, i.e. one close to a true constant. The Model 1 exponent $q = 1/2$ is the worst column of the map ($\Delta\chi^2 > 1000$).

7 Discussion

We have evaluated the fluctuations of the BD action that MYZ left as an open question. The central result is structural: in d dimensions the variance is dominated by the Poisson noise in the number of nearly null intervals, whose number is controlled by the volume of the boost group that fits inside the region, and this gives $\Delta S \propto (\rho V)^{1-1/d}$. Only for $d = 2$ is the exponent $1/2$, and then with a logarithm. In four dimensions the exponent is $3/4$ in every region we tested, including the past cone of an FLRW observer reaching the big bang, and for the local and the smeared actions. A cosmological constant sourced by these fluctuations would scale as $\delta\Lambda \propto V^{-1/4}$, not as the $V^{-1/2}$ of Everpresent Λ, and a stochastic history built on this scaling does not fit current data.

This does not refute Everpresent Λ as such, whose original argument relies on the Poisson fluctuation of N and not on the action. It shows that the action, the most natural dynamical quantity of the theory, does not provide the required mechanism. The obstruction is Lorentz invariance itself: any observable that counts intervals of fixed small volume inherits the boost range of the region. The same non-compactness biases the statistics of links in sprinkled causal sets [16]. A mechanism that yields $V^{-1/2}$ would have to suppress the contribution of highly boosted intervals, for instance through a boundary term or through a choice of frame that breaks local Lorentz invariance.

Our conclusions have the following limitations.

1. *Asymptotics.* The exponent 3/4 rests on one exact ingredient ($J^1 \propto (\rho V)^{3/2}$, verified up to $\rho V = 10^8$) and one empirical one: that the correlations remain a constant fraction of J^1. We verified the latter up to $\rho V = 12\,800$ in the diamond and 6400 in the past cones, with a residual power law bounded by $\delta < 0.07$ (2σ), and for the smeared action up to $\epsilon\rho V = 1280$. The physically relevant regime, $\rho V \sim 10^{242}$ with $\epsilon \sim 10^{-49}$, is an extrapolation over some 238 decades, which the heuristic argument of section 3 supports but does not prove. In particular, for $\epsilon \leq 0.2$ the individual MYZ terms still drift at the largest densities, and only their sum is constant.

2. *Identification.* Equation (12) is an assumption, motivated by the cosmological term of the Einstein–Hilbert action and consistent with Model 1, but not derived from a dynamics for causal sets.

3. *Dynamics.* The data analysis uses a homogeneous $\Lambda(t)$ and the Ornstein–Uhlenbeck family (15). Spatial fluctuations of Λ are further constrained by the isotropy of the CMB [30]; non-Gaussian processes or a different coupling between the action and Λ were not explored. The temporal memory was measured for the local action only.

4. *Statistics.* The per-seed fits use a grid search rather than the MCMC of [8], which may underestimate the fraction of good seeds; with $\Delta\chi^2_{\min} = +124$ this would not change the conclusion. Our supernova-only ΛCDM χ^2 differs from that of [8] (section 6); all our margins are relative to our own fits.

Earlier cosmological tests of Everpresent Λ are those of [5, 31, 8]; to our knowledge this is the first confrontation of Everpresent-type dynamics with the DESI DR2 BAO data.

Acknowledgments
The numerical code, the analysis and the drafting of this manuscript were carried out with the assistance of Claude (Anthropic), an AI model. The author designed the project, reviewed all results and takes full responsibility for the content.

Data availability
The code that produces every number and figure of this paper, together with its test suite (213 automated tests) and the saved intermediate results, is publicly available at https://github.com/carlosecheverriaarrondo/Conjuntos_causales and archived on Zenodo, https://doi.org/10.5281/zenodo.23413234. The supernova and BAO data are public and are downloaded, with checksum verification, by a script in the repository. Most tests check the code against exact results or fresh sprinklings; some re-read the saved results of the long runs and check the criteria quoted here. All statistical tolerances were fixed in advance, and each group of tests was checked to fail under a deliberately introduced error. A detailed log of every criterion, including those that failed, is part of the repository.

A Covariance with the number of elements

Let η be a Poisson process of intensity ρ on M and $F(\eta)$ a functional with finite second moment. The Mecke equation [19, theorem 4.1] gives $\mathbb{E}[NF] = \rho \int_M \mathbb{E}[F(\eta + \delta_x)]\,\mathrm{d}x$, while differentiating with respect to the intensity gives $\rho\,\mathrm{d}\mathbb{E}[F]/\mathrm{d}\rho = \rho \int_M \mathbb{E}[F(\eta + \delta_x) - F(\eta)]\,\mathrm{d}x$. Subtracting $\mathbb{E}[N]\,\mathbb{E}[F] = \rho V\,\mathbb{E}[F]$ from the first expression yields (5). For the local action, $F = \sum_k a_k N_k$ and $\langle N_k \rangle$ depends on ρ only through ρV, which leads to $\mathrm{Cov}(N, N_k) = (k+2)L_k - (k+1)L_{k+1}$. We evaluate $\mathrm{d}\langle F \rangle/\mathrm{d}\rho$ with a central difference and checked it against this identity (relative error below 10^{-4}) and against sprinklings of the smeared action.

B Estimators of the MYZ terms

Smeared action. With thinned abundances the MYZ formalism carries over with three changes: the means become $\langle \tilde{N}_k \rangle = L_k(\epsilon\rho V)/\epsilon^2$; J^1 is no longer diagonal in k, because the two factors of a single pair are independent thinnings of the same interval; and in the joint distribution of two overlapping intervals the overlap is split into points kept by both (probability ϵ^2) and points kept by only one ($\epsilon(1-\epsilon)$ each), since thinning a Poisson process produces independent Poisson processes.

Per-sprinkling estimators. Let $f(y,x) = f(n(y,x))$ for $y \prec x$ and 0 otherwise, with row sums $R_y = \sum_x f(y,x)$ and column sums $C_x = \sum_y f(y,x)$. Sorting the terms of F^2 by shared endpoints gives, sprinkling by sprinkling,

$$\sum f^2 \to J^1, \quad \sum_y R_y^2 - \sum f^2 \to J^2, \quad \sum_x C_x^2 - \sum f^2 \to J^3, \quad \sum_z C_z R_z \to J^4, \tag{16}$$

and J_c^6 is obtained by subtracting all of them from $(F - \langle F \rangle)^2$. The expectation of each statistic is the corresponding MYZ term, so the estimators are unbiased up to $O(1/M)$ from the sample mean. We checked the decomposition against an explicit enumeration of pairs of pairs.

References

[1] Bombelli L, Lee J, Meyer D and Sorkin R D 1987 *Phys. Rev. Lett.* **59** 521

[2] Surya S 2019 *Living Rev. Relativ.* **22** 5

[3] Sorkin R D 1997 *Int. J. Theor. Phys.* **36** 2759 (*Preprint* gr-qc/9706002)

[4] Sorkin R D 2007 *AIP Conf. Proc.* **957** 142 (*Preprint* 0710.1675)

[5] Ahmed M, Dodelson S, Greene P B and Sorkin R 2004 *Phys. Rev. D* **69** 103523 (*Preprint* astro-ph/0209274)

[6] Ahmed M and Sorkin R 2013 *Phys. Rev. D* **87** 063515 (*Preprint* 1210.2589)

[7] Das S, Nasiri A and Yazdi Y K 2023 *J. Cosmol. Astropart. Phys.* **10** 047 (*Preprint* 2304.03819)

[8] Das S, Nasiri A and Yazdi Y K 2024 *J. Cosmol. Astropart. Phys.* **10** 076 (*Preprint* 2307.13743)

[9] Benincasa D M T and Dowker F 2010 *Phys. Rev. Lett.* **104** 181301 (*Preprint* 1001.2725)

[10] Dowker F and Glaser L 2013 *Class. Quantum Grav.* **30** 195016 (*Preprint* 1305.2588)

[11] Buck M, Dowker F, Jubb I and Surya S 2015 *Class. Quantum Grav.* **32** 205004 (*Preprint* 1502.05388)

[12] Dowker F, Liu R and Lloyd-Jones D 2025 *Class. Quantum Grav.* **42** 205006 (*Preprint* 2501.00139)

[13] Moradi H, Yazdi Y K and Zilhão M 2025 *Class. Quantum Grav.* **42** 045017 (*Preprint* 2407.03395)

[14] Bombelli L, Henson J and Sorkin R D 2009 *Mod. Phys. Lett. A* **24** 2579 (*Preprint* gr-qc/0605006)

[15] Glaser L and Surya S 2013 *Phys. Rev. D* **88** 124026 (*Preprint* 1309.3403)

[16] Boguñá M and Krioukov D 2026 Sampling biases in sprinkled causal sets (*Preprint* 2610.03247)

[17] Benincasa D M T, Dowker F and Schmitzer B 2011 *Class. Quantum Grav.* **28** 105018 (*Preprint* 1011.5191)

[18] Sorkin R D 2009 Does locality fail at intermediate length scales? *Approaches to Quantum Gravity* ed Oriti D (Cambridge: Cambridge University Press) pp 26–43 (*Preprint* gr-qc/0703099)

[19] Last G and Penrose M 2017 *Lectures on the Poisson Process* (Cambridge: Cambridge University Press)

[20] Planck Collaboration 2020 *Astron. Astrophys.* **641** A6 (*Preprint* 1807.06209)

[21] Belenchia A, Benincasa D M T, Liberati S, Marin F, Marino F and Ortolan A 2016 *Phys. Rev. Lett.* **116** 161303 (*Preprint* 1512.02083)

[22] Uhlenbeck G E and Ornstein L S 1930 *Phys. Rev.* **36** 823

[23] Brout D *et al.* 2022 *Astrophys. J.* **938** 110 (*Preprint* 2202.04077)

[24] Scolnic D *et al.* 2022 *Astrophys. J.* **938** 113 (*Preprint* 2112.03863)

[25] Riess A G *et al.* 2022 *Astrophys. J. Lett.* **934** L7 (*Preprint* 2112.04510)

[26] DESI Collaboration 2025 *Phys. Rev. D* **112** 083515 (*Preprint* 2503.14738)

[27] Chen L, Huang Q G and Wang K 2019 *J. Cosmol. Astropart. Phys.* **02** 028 (*Preprint* 1808.05724)

[28] Hu W and Sugiyama N 1996 *Astrophys. J.* **471** 542

[29] Eisenstein D J and Hu W 1998 *Astrophys. J.* **496** 605

[30] Barrow J D 2007 *Phys. Rev. D* **75** 067301 (*Preprint* gr-qc/0612128)

[31] Zwane N, Afshordi N and Sorkin R D 2018 *Class. Quantum Grav.* **35** 194002 (*Preprint* 1703.06265)