

The twelve-block colouring is optimal: $\delta_3 = 117/2192$
a computer-assisted proof, its certificates re-decided by two programs that share no code

Carlos Toledo*

October 2026 — draft v0.2, rebuilt in cert-machine; every number interpolated from the
record; not refereed, not formalised

Abstract

Let $V(n)$ be the least number of monochromatic three-term arithmetic progressions over all two-colourings of $\{1, \dots, n\}$. Graham offered \$100 for the value of $\delta_3 = \lim V(n)/n^2$, the case $k = 3$ of Erdős problem #1186. Parrilo, Robertson and Saracino (2008) proved $1675/32768 \leq \delta_3 \leq 117/2192$, with the upper bound attained by a colouring in twelve blocks, and conjectured the upper bound correct. We prove it: $\delta_3 = 117/2192$. By their counting lemma the statement reduces to a quadratic-form inequality, $Q(\varphi) = \iint_{a/2 \leq b \leq (1+a)/2} \varphi(a)\varphi(b) \geq -5/137$ for every measurable $\varphi : [0, 1] \rightarrow [-1, 1]$, and we show more: the twelve-block function and its negative are the only minimisers. The proof expands Q around the conjectured optimum, where the first-order term is an explicit nonnegative piecewise-linear function h ; a discretisation built on h loses nothing to first order; the space is cut into five regions by the L^1 distance to the optimum, and each region is closed by an exact rational certificate — zero gap near the optimum, a margin of at least 2.747×10^{-4} away from it. The certificates were generated and checked in one session by a program and a verifier written together; they are re-decided here by a second verifier, written in a second language from the mathematics and the file format alone, which shares no code with either. Both accept the five certificates; both refuse every planted forgery. Nothing here has been refereed or formalised, and we say exactly what that leaves open.

1 Introduction

Colour $1, \dots, n$ red and blue. How few monochromatic progressions $\{a, a + d, a + 2d\}$, $d \geq 1$, can there be? A random colouring gives about $n^2/16$. At the Erdős conference in Budapest in 1999 Graham proposed, as a \$100 problem, to determine β in $V(n) = \beta n^2(1 + o(1))$ [1]. Parrilo, Robertson and Saracino [1] disproved the folklore guess $\beta = 1/16$ with a colouring in twelve blocks of relative lengths

$$28, 6, 28, 37, 59, 116, 116, 59, 37, 28, 6, 28 \quad (\text{out of } 548),$$

alternating in colour, which gives $V(n) \leq \frac{117}{2192}n^2(1 + o(1))$; they proved $V(n) \geq \frac{1675}{32768}n^2(1 + o(1))$ by a semidefinite relaxation and conjectured the upper bound to be the truth. Butler, Costello

*Independent researcher. Contact: carlos@carlostoledo.co. This work was produced with substantial AI assistance in a certified-verification setting: constructions and exposition were proposed with the help of large language models, and every quantitative claim is decided by outward-rounded interval arithmetic or exact rational arithmetic, with adversarial controls that must fire for the certificate to be accepted. The certificates, not the models, are the evidence, and they are public and re-runnable. Machine-derived and not peer-reviewed.

and Graham [2] recover the twelve blocks by experiment, crediting [1], and give block colourings for longer progressions. Greenwood, Kariv and Williams [3] prove the twelve-block colouring optimal among antisymmetric colourings with at most twelve blocks. On the problem's page [5] the question is open, as is the general δ_k .

Theorem 1. $\delta_3 = \lim_{n \rightarrow \infty} \frac{V(n)}{n^2} = \frac{117}{2192} = 0.0533759124\dots$

Scope. Erdős problem #1186 asks for bounds or an asymptotic formula for δ_k for every k . Theorem 1 settles $k = 3$, Graham's question and the conjecture of [1]. It says nothing about $k \geq 4$.

How the proof is built, and how it is checked. Sections 2–4 are mathematics a reader can check by hand in an hour: the reduction of [1], an exact expansion around the twelve blocks, and a discretisation lemma. Section 5 is five finite exact certificates. Section 6 is how they were checked: by the verifier written with them, and by a second verifier written afterwards under a clean-room rule. Section 8 lists what is not established.

2 The reduction

We re-derive the counting lemma of [1]. Fix a colouring χ of $[n]$. A progression $\{a < c < b\}$, $a+b = 2c$, that is not monochromatic has exactly two bichromatic pairs among (a, c) , (c, b) , (a, b) . Count incidences of (non-monochromatic progression, bichromatic pair):

- the *outer* pair (a, b) determines the progression and has $a \equiv b \pmod{2}$; so outer incidences number $T = \#\{a < b : a \equiv b \pmod{2}, \chi(a) \neq \chi(b)\}$;
- an ordered bichromatic pair (u, v) is *inner* to the progression $\{u, v, 2v - u\}$ with v the middle term, whenever $2v - u \in [n]$; every progression has exactly two inner ordered pairs, (a, c) and (b, c) ; so inner incidences number $N^+ = \#\{(u, v) : \chi(u) \neq \chi(v), 2v - u \in [n]\}$.

Hence the number of non-monochromatic progressions is $(T + N^+)/2$, and with $n^2/4 + O(n)$ progressions in all,

$$2V = \frac{n^2}{2} - N^+ - T + O(n).$$

Writing r_o, b_o and r_e, b_e for the colour counts in the odd and even classes, $T = r_o b_o + r_e b_e \leq 2 \cdot (n/4)^2 + O(n) = n^2/8 + O(n)$.

Let $\varphi : [0, 1] \rightarrow \{\pm 1\}$ be the step function of χ , constant on each $((u-1)/n, u/n]$. With $u = na$, $v = nb$ the condition $1 \leq 2v - u \leq n$ reads $a/2 < b \leq (1+a)/2$ up to the cells that meet the boundary of

$$R = \{(a, b) \in [0, 1]^2 : a/2 \leq b \leq (1+a)/2\}, \quad |R| = \frac{1}{2},$$

and there are $O(n)$ such cells, whatever the colouring. So, uniformly in χ ,

$$N^+ = n^2 \iint_R \frac{1}{2} (1 - \varphi(a)\varphi(b)) da db + O(n) = \frac{n^2}{2} \left(\frac{1}{2} - Q(\varphi) \right) + O(n), \quad Q(\varphi) = \iint_R \varphi(a)\varphi(b).$$

Combining,

$$\frac{V(n)}{n^2} \geq \frac{1}{16} + \frac{Q^*}{4} - O\left(\frac{1}{n}\right), \quad Q^* = \inf \{Q(\varphi) : \varphi : [0, 1] \rightarrow [-1, 1] \text{ measurable}\}. \quad (1)$$

Let φ^* be the twelve-block function, +1 on the first block. Then $Q(\varphi^*) = -5/137$ (Lemma 1 recomputes it), and $\frac{1}{16} - \frac{5}{548} = \frac{117}{2192}$. The twelve-block colouring is antisymmetric, so its parity classes are balanced, $T = n^2/8 + O(n)$, and (1) is an equality for it: that is the upper bound of [1], and §6 counts it exactly for n up to 35,072. So Theorem 1 follows from:

Theorem 2. $Q(\varphi) \geq -5/137$ for every measurable $\varphi : [0, 1] \rightarrow [-1, 1]$, with equality only for $\varphi = \pm\varphi^*$ almost everywhere.

3 Expanding around the twelve blocks

Let K be the symmetrised kernel of R , so that $Q(\varphi) = \langle \varphi, K\varphi \rangle$, and put $g^* = K\varphi^*$,

$$g^*(a) = \frac{1}{2} \left(\int_{a/2}^{(1+a)/2} \varphi^* + \int_{[2a-1, 2a] \cap [0, 1]} \varphi^* \right), \quad h = -\varphi^* g^*.$$

Every φ with values in $[-1, 1]$ is $\varphi = \varphi^*(1 - 2\sigma)$ with $\sigma = (1 - \varphi\varphi^*)/2 \in [0, 1]$, the part of φ flipped against φ^* ; $m := \int \sigma = \frac{1}{2} \|\varphi - \varphi^*\|_1$. Expanding the quadratic form,

$$Q(\varphi) = Q(\varphi^*) - 4\langle \varphi^* \sigma, K\varphi^* \rangle + 4Q(\varphi^* \sigma),$$

which is the exact, global identity

$$E(\sigma) := \frac{Q(\varphi) - Q(\varphi^*)}{4} = \int_0^1 h \sigma + Q(\varphi^* \sigma). \quad (2)$$

Lemma 1. h is continuous and piecewise linear with nodes at the multiples of $1/1096$; $h \geq 0$; h vanishes exactly at the eleven breakpoints of φ^* , where its one-sided slopes are 1 or $3/2$; and $\int_0^1 h = 5/137 = -Q(\varphi^*)$.

Proof. The kinks of g^* occur only where $a/2$, $(1+a)/2$, $2a$ or $2a-1$ meets an edge of φ^* , or at $a = \frac{1}{2}$. Every edge is a multiple of $1/548 = 2/1096$, so every kink is a multiple of $1/1096$, and φ^* is constant on every interval $[k/1096, (k+1)/1096]$. The node values are exact rationals, computed; at every edge $g^* = 0$, so h is continuous there. The remaining claims are finite checks on the 1097 node values. $\square$

Since $\varphi \mapsto -\varphi$ sends σ to $1 - \sigma$ and does not change Q , it suffices to prove $E(\sigma) \geq 0$ when $m \leq \frac{1}{2}$.

4 A discretisation without first-order loss

Partition $[0, 1]$ into cells I_i whose endpoints include every edge of φ^* . On I_i the sign of φ^* is a constant s_i ; write $w_i = |I_i|$ and $y_i = w_i^{-1} \int_{I_i} \sigma \in [0, 1]$, so that $m = \sum_i w_i y_i$.

Lemma 2 (bathtub). *Let $a_i = w_i \min_{I_i} h$ and $m_i(v) = |\{a \in I_i : h(a) < v\}|$; since h is piecewise linear, $m'_i(v)$ is the sum of $1/|\text{slope}|$ over every linear piece of h on I_i whose range contains v . Put $b_i = \frac{w_i^2}{2} / \sup_v m'_i(v)$, or $b_i = 0$ if h is constant on some piece of I_i . Then $\int_{I_i} h \sigma \geq a_i y_i + b_i y_i^2$ whenever $0 \leq \sigma \leq 1$.*

Proof. For a fixed average y , the integral is least when σ fills the lowest levels of h , so it is at least $\beta(y) = \int_0^{w_i y} h^\uparrow, h^\uparrow$ the increasing rearrangement of h on I_i . Then $\beta(0) = 0$, $\beta'(0) = w_i \min h = a_i$, and $\beta''(y) = w_i^2/m'_i(h^\uparrow(w_i y)) \geq 2b_i$; Taylor's formula gives the claim. $\square$

Pairs of cells are ordered: (i, j) means $a \in I_i, b \in I_j$. A pair is *full* if $I_i \times I_j \subset R$ and *cut* if ∂R crosses it, with A_{ij}^{in} and A_{ij}^{out} the areas inside and outside R .

Lemma 3. $E(\sigma) \geq L(y)$, where

$$L(y) = \sum_i (a_i y_i + b_i y_i^2) + \sum_{\text{full}} s_i s_j w_i w_j y_i y_j \\ + \sum_{\text{cut}, s_i s_j = 1} \max(0, w_i w_j y_i y_j - A_{ij}^{\text{out}}) + \sum_{\text{cut}, s_i s_j = -1} \max(-A_{ij}^{\text{in}}, -w_i w_j y_i y_j).$$

Proof. Lemma 2 bounds $\int h \sigma$ cell by cell. In $Q(\varphi^* \sigma) = \sum_{(i,j)} s_i s_j \iint_{R \cap (I_i \times I_j)} \sigma(a) \sigma(b)$ a full pair factorises exactly. On a cut pair $0 \leq \iint_{\text{in}} \sigma \sigma \leq \min(A^{\text{in}}, w_i w_j y_i y_j)$ because $0 \leq \sigma \leq 1$, and $\iint_{\text{in}} \sigma \sigma = \iint_{I_i \times I_j} \sigma \sigma - \iint_{\text{out}} \sigma \sigma \geq w_i w_j y_i y_j - A^{\text{out}}$. $\square$

$L(0) = 0$, and the first-order part of L at $y = 0$ is exact up to the bathtub. That is the difference from discretising in φ itself, as in [1]: there every cut pair loses a first-order amount, and §7 measures what that costs.

5 Five certificates

We cover $m \in [0, \frac{1}{2}]$ by five regions, each with its own grid and an exact certificate (Table 1). The grids are in units of $1/1096$: one of 130 cells (multiples of $24/1096$, steps of $4/1096$ within $8/1096$ of every edge, and mirror images), one of 146 cells (multiples of $8/1096$ and the edges).

Slabs $\text{lo} \leq m \leq \text{hi}$. An exact polynomial identity in $v = (1, y)$,

$$L_\theta(y) - B = v^\top S v + \sum_f \mu_f f(y), \quad \mu \geq 0, \theta \in [0, 1], S + \varepsilon I \succeq 0.$$

$L_\theta \leq L$ replaces each maximum in Lemma 3 by the convex combination with weight θ . Each form f is a product of two factors among $y_i, 1 - y_i, m - \text{lo}, \text{hi} - m$, nonnegative on the region, or a triangle inequality $1 + s_1 z_i z_j + s_2 z_j z_k + s_3 z_i z_k$ with $s_1 s_2 s_3 = 1, z = 1 - 2y$ and distinct i, j, k , nonnegative on the cube because it is multilinear and nonnegative at the vertices. Since $v^\top v \leq 1 + n, L \geq B - \varepsilon(1 + n)$ on the region. The products with $m - \text{lo}$ and $\text{hi} - m$ matter: they stop the relaxation from averaging the two optima $\pm \varphi^*$, and from averaging points of different regions.

The region $m \leq 1/5$. An identity in which every term vanishes at $y = 0$,

$$L_0(y) = y^\top P y + \sum_{i,j} \lambda_{ij} y_i (1 - y_j) + \sum_j \nu_j y_j (1/5 - m) + \kappa m (1/5 - m) + y^\top N y + \sum_i c_i y_i,$$

with $P \succ 0$, every multiplier nonnegative and N entrywise nonnegative (enough, since $y \geq 0$; the κ term uses $m \geq 0$), where $L_0 \leq L$ takes the cut bounds 0 and $-w_i w_j y_i y_j$. The forms

region $m = \int \sigma$	cells	certified bound on E	forms	verdict here
$[0, 1/5]$	130	0 (exact at $\sigma = 0$)	22,808	verified
$[1/5, 3/10]$	130	2.784897×10^{-4}	13,324	verified
$[3/10, 2/5]$	130	3.035513×10^{-4}	9,849 (2,204 triangles)	verified
$[2/5, 9/20]$	146	2.747055×10^{-4}	1,998	verified
$[9/20, 1/2]$	146	4.638341×10^{-4}	2,332 (470 triangles)	verified

Table 1: The cover of $m \in [0, \frac{1}{2}]$. Bounds are the exact certified values recomputed by the second verifier, printed truncated; the inner region’s bound is 0, attained only at $\sigma = 0$.

$y_j(1 - y_i)$ turn the linear budget $a_i y_i$ into exactly the bathtub’s quadratic cost, and the positive-definite Hessian of Q in the breakpoint shifts of φ^* (smallest eigenvalue about 0.1266, a float measurement of the claimant’s, not used) supplies the margin.

Proof of Theorem 2. The five regions cover $m \in [0, \frac{1}{2}]$, so $E \geq 0$ whenever $m \leq \frac{1}{2}$, and by symmetry everywhere. If $m \leq 1/5$ and $y \neq 0$, then $E \geq L_0(y) \geq y^\top P y > 0$, because P is proved positive definite (§6); if $y = 0$ then $\sigma = 0$ almost everywhere. Every slab has $E > 0$. So $E = 0$ only at $\sigma = 0$, that is $\varphi = \varphi^*$; the case $m \geq \frac{1}{2}$ gives $-\varphi^*$. $\square$

Corollary 1. φ^* minimises Q on the ball $\|\varphi - \varphi^*\|_1 \leq 2/5$, fractional perturbations included, and every φ with $\frac{1}{2}\|\varphi \mp \varphi^*\|_1 \geq 1/5$ for both signs has $Q(\varphi) \geq -\frac{5}{137} + 4 \cdot 2.747 \times 10^{-4}$.

6 Verification

The claimant’s pipeline. The multipliers come from floating-point semidefinite programs, rounded to rationals; S or P is then recomputed exactly and its semidefiniteness proved exactly. A verifier written in the same session, separately from the generators, re-derives φ^* , h , the bathtub minorants and the cell areas and re-checks the five files (87 s on this machine, re-run from the pinned bytes). Both were written by one AI agent in one session, so a misreading common to both would not be caught — the gap this section closes as far as a program can.

The second verifier, under a clean-room rule. It was written in JavaScript with BigInt rationals, in this repository, by an agent that was given only the mathematics (the statement and the proofs of the lemmas above, as the claimant wrote them) and a field-by-field description of the certificate files written from the files’ serialisers. It did not read the generators, the claimant’s verifier or its logs, and the claimant’s code entered the repository only after it was finished. It derives φ^* , g^* , h , the bathtub constants and the residual matrices from the mathematics, and every cell area by two exact routes that must agree pair by pair (the trapezoid rule on the piecewise-linear section length, and polygon clipping with the shoelace formula). It refuses any form it cannot show nonnegative on its region and any multiplier of the wrong sign. It proves $S + \varepsilon I \succeq 0$ and $P \succ 0$ exactly: after an exact rescaling by powers of two, a floating-point Cholesky factor of $A - 2^{-k}I$ is rounded to a dyadic matrix L , and the remainder $A - LL^\top$ is shown, exactly, to be diagonally dominant; a matrix it cannot so prove is refused. It accepts the five certificates in 0.8 s and recomputes every bound in Table 1, each equal to the bound the file claims. $Q(\varphi^*) = -5/137$ and $\int h = 5/137$ are recomputed independently, the first from block-pair areas alone.

What the second reading found. Nothing that changes a verdict, and four points the text above now states. (i) The bathtub constant needs the *summed* m'_i of Lemma 2: read with the largest single $1/|\text{slope}|$ instead, b_i is too large and the inequality fails — 1,358 exact violations in 2,432 tests — on the cells where h is not monotone. The certificates were built with the summed reading. (ii) ε is load-bearing: no slab’s S is semidefinite by itself (exact negative witnesses), and the bound $B - \varepsilon(1 + n)$ pays for it. (iii) The triangle forms need three distinct indices. (iv) The kink list of Lemma 1 includes $a = \frac{1}{2}$ and the edges themselves, all on the lattice.

Forgeries and mutations. The second verifier’s battery runs 76 checks; 40 of them are planted forgeries it must refuse, each by the rule it breaks: bounds raised by 10^{-3} and by 10^{-6} (the certificates are that tight), dropped triangle forms, every θ set to 1 or to 0, widened slabs, ε set to 0 or inflated together with B , a grid missing an edge of φ^* , a doubled inner radius, an overspent linear budget, dropped inner multipliers, missing regions, and single entries off by 2^{-100} — a negative multiplier, a sign-violating or repeated-index triangle, θ outside $[0, 1]$, an unknown form. Every one is refused at this build. The verifier itself was then mutated by one-line edits (a flipped h , a mirrored R , a dropped ε , an optimistic cut bound, a skipped sign check, a trusted floating-point Cholesky, an accepted zero pivot, ...): 23 of 24 are caught by the genuine set or a forgery, and the one survivor removes a rule that is enforced twice.

The two lemmas, tested. Identity (2) and Lemma 3 are proved above; the second verifier also tests them exactly, in rationals, on step functions finer than the certificate grids: 1,400 cases of (2), all equalities, and 4,200 cases of Lemma 3 from seven generators, adversarial ones included, all satisfied, with $E - L$ exactly 0 on breakpoint flips and single cells. These are tests, not proofs.

The upper bound, counted. The exact number of monochromatic progressions of the twelve-block colouring of $[n]$, $n = 548k$; for n a multiple of 1096 it is exactly $\frac{117}{2192}n^2 - \frac{n}{2}$:

n	$V_{12\text{-block}}(n)$	V/n^2	$n(V/n^2 - \frac{117}{2192})$
548	15,756	0.0524668336	−0.498175
1,096	63,568	0.0529197080	−0.500000
2,192	255,368	0.0531478102	−0.500000
4,384	1,023,664	0.0532618613	−0.500000
8,768	4,099,040	0.0533188869	−0.500000
17,536	16,404,928	0.0533473996	−0.500000
35,072	65,637,248	0.0533616560	−0.500000

approaching $117/2192 = 0.0533759124\dots$ from below at rate $O(1/n)$, as the reduction predicts.

7 Why the obvious discretisation stops short

Before the expansion around φ^* , the same claimant certified lower bounds by discretising φ itself on L equal cells, with the cut-pair bound $\max(-A^{\text{in}}, x_i x_j / L^2 - A^{\text{out}})$ and a Shor relaxation with triangle cuts. Its own verifier accepted (these were not re-decided here):

L	certified $\delta_3 \geq$	local-search minimum of the L -cell model (float)
32	0.05020594	0.052000
64	0.05219317	0.053031
128	0.05296151	0.053267
192	0.05317533	0.053331
256	0.05324921	0.053347

Every cut pair gives away a first-order amount in this model, so even its minimum — the right column, measured by local search in floating point — sits strictly below 117/2192 at every L : the method cannot reach the answer, however fine the grid. In the coordinates σ the first-order term is the exact $\int h\sigma$, bounded cell by cell without loss at $\sigma = 0$, and the discretisation error is second order. That measurement is what led to the proof.

8 What is established, and what is not

- **Established, by machine:** five exact certificates, accepted by two verifiers written independently in two languages, each refusing the planted forgeries; the reduction, the expansion and the two lemmas proved in §§2–4.
- **Read, not refereed:** the author read the proof line by line on 5 October 2026, after both verifiers had accepted the certificates; no referee has read it.
- **Not established:** nothing is formalised; both verifiers and the generators were written by AI agents, and the clean-room rule removes shared code, not a shared misreading of the mathematics the agents were both given — §§2–4 are where a reader should look. Generation is not reproducible byte for byte (the solver is nondeterministic); the certificate files are the record, and their force does not depend on regeneration.
- **Not claimed:** anything about δ_k for $k \geq 4$, or about the $\mathbb{F}_p$ analogue. The method does not carry over as it stands: for $k = 4$ the counting functional has a quartic term that no reduction of the kind in §2 removes, and the best known colouring is arithmetic, not a block colouring (1/72, Lu and Peng [4], against 0.01722 for the best block colouring [2]); both values were re-decided in exact arithmetic in a companion scout.

Reproduction

Every constant in this paper is interpolated from a certificate record in the `cert-machine` repository, so the document cannot quote a number the machine did not certify. The five certificates are `certs/delta3/*.json` (sha256 in `certs/delta3-ledger.json`); `node instruments/delta3/verify.js` re-decides them, `node instruments/delta3/battery.js` runs the forgeries, `node instruments/delta3/lemmas` the lemma tests and the upper-bound counts. The claimant’s generators and verifier are kept, pinned by sha256, in `instruments/delta3/claimant/`. Repository commit `ce470dd1d090`.

References

- [1] P. A. Parrilo, A. Robertson, D. Saracino, On the asymptotic minimum number of monochromatic 3-term arithmetic progressions, *J. Combin. Theory Ser. A* 115 (2008) 185–192; arXiv:math/0609532.
- [2] S. Butler, K. P. Costello, R. Graham, Finding patterns avoiding many monochromatic constellations, *Experiment. Math.* 19 (2010) 399–411.
- [3] T. Greenwood, J. Kariv, N. Williams, From discrete to continuous: monochromatic 3-term arithmetic progressions, arXiv:2301.00336.
- [4] L. Lu, X. Peng, Monochromatic 4-term arithmetic progressions in 2-colorings of $\mathbb{Z}_n$, arXiv:1107.2888.
- [5] T. F. Bloom, Erdős problem #1186, <https://www.erdosproblems.com/1186>, snapshot of 5 October 2026.