

Central Limit Theorem

David Carlson

2021

CLT Overview

- ▶ Incredibly important, fundamental theorem in probability theory

CLT Overview

- ▶ Incredibly important, fundamental theorem in probability theory
- ▶ Used in many cases, perhaps most importantly in regression

CLT Overview

- ▶ Incredibly important, fundamental theorem in probability theory
- ▶ Used in many cases, perhaps most importantly in regression
- ▶ Establishes that when independent random variables are summed, their properly normalized sum tends towards a normal distribution *irrelevant* of the variables' distributions

CLT Overview

- ▶ Incredibly important, fundamental theorem in probability theory
- ▶ Used in many cases, perhaps most importantly in regression
- ▶ Establishes that when independent random variables are summed, their properly normalized sum tends towards a normal distribution *irrelevant* of the variables' distributions
- ▶ Implies that probabilistic and statistical methods that work for normal distributions can be applicable to many problems involving other types of distributions

More Formally

- ▶ Let X_1, X_2, \dots, X_n are n random samples drawn from a population with overall mean μ and *finite* variance σ^2

More Formally

- ▶ Let X_1, X_2, \dots, X_n are n random samples drawn from a population with overall mean μ and *finite* variance σ^2
- ▶ Let \bar{X}_n be the sample mean

More Formally

- ▶ Let X_1, X_2, \dots, X_n are n random samples drawn from a population with overall mean μ and *finite* variance σ^2
- ▶ Let \bar{X}_n be the sample mean
- ▶ Let $Z = \lim_{n \rightarrow \infty} \sqrt{n} \left(\frac{\bar{X}_n - \mu}{\sigma} \right)$

More Formally

- ▶ Let X_1, X_2, \dots, X_n are n random samples drawn from a population with overall mean μ and *finite* variance σ^2
- ▶ Let \bar{X}_n be the sample mean
- ▶ Let $Z = \lim_{n \rightarrow \infty} \sqrt{n} \left(\frac{\bar{X}_n - \mu}{\sigma} \right)$
- ▶ Z is a *standard* normal distribution

Example

- ▶ Suppose that a sample is obtained containing many observations, each observation being randomly generated in a way that does not depend on the values of the other observations, and that the arithmetic mean of the observed values is computed

Example

- ▶ Suppose that a sample is obtained containing many observations, each observation being randomly generated in a way that does not depend on the values of the other observations, and that the arithmetic mean of the observed values is computed
- ▶ If this procedure is performed many times, the central limit theorem says that the probability distribution of the average will closely approximate a normal distribution

Example

- ▶ Suppose that a sample is obtained containing many observations, each observation being randomly generated in a way that does not depend on the values of the other observations, and that the arithmetic mean of the observed values is computed
- ▶ If this procedure is performed many times, the central limit theorem says that the probability distribution of the average will closely approximate a normal distribution
- ▶ A simple example of this is that if one flips a coin many times, the probability of getting a given number of heads will approach a normal distribution, with the mean equal to half the total number of flips

Example

- ▶ Suppose that a sample is obtained containing many observations, each observation being randomly generated in a way that does not depend on the values of the other observations, and that the arithmetic mean of the observed values is computed
- ▶ If this procedure is performed many times, the central limit theorem says that the probability distribution of the average will closely approximate a normal distribution
- ▶ A simple example of this is that if one flips a coin many times, the probability of getting a given number of heads will approach a normal distribution, with the mean equal to half the total number of flips
- ▶ At the limit of an infinite number of flips, it will equal a normal distribution

Regression and CLT

- ▶ Regression analysis and in particular ordinary least squares specifies that a dependent variable depends according to some function upon one or more independent variables, with an additive error term

Regression and CLT

- ▶ Regression analysis and in particular ordinary least squares specifies that a dependent variable depends according to some function upon one or more independent variables, with an additive error term
- ▶ Various types of statistical inference on the regression assume that the error term is normally distributed

Regression and CLT

- ▶ Regression analysis and in particular ordinary least squares specifies that a dependent variable depends according to some function upon one or more independent variables, with an additive error term
- ▶ Various types of statistical inference on the regression assume that the error term is normally distributed
- ▶ This assumption can be justified by assuming that the error term is actually the sum of many independent error terms; even if the individual error terms are not normally distributed, by the central limit theorem their sum can be well approximated by a normal distribution

Illustration

```
http://195.134.76.37/applets/  
AppletCentralLimit/App1_CentralLimit2.  
html
```