


Sets and functions

David Carlson

2021

Sets and Functions



Coordinate Geometry

- Pairs of numbers represent points:
 $(x_1, y_1), (x_2, y_2), (x_3, y_3), \dots, (x_n, y_n)$

Coordinate Geometry

- ▶ Pairs of numbers represent points:
 $(x_1, y_1), (x_2, y_2), (x_3, y_3), \dots, (x_n, y_n)$
- ▶ The plot has two axes:

Coordinate Geometry

- ▶ Pairs of numbers represent points:
 $(x_1, y_1), (x_2, y_2), (x_3, y_3), \dots, (x_n, y_n)$
- ▶ The plot has two axes:
 - ▶ x-axis is the horizontal axis at the bottom of the plot

Coordinate Geometry


- ▶ Pairs of numbers represent points:
 $(x_1, y_1), (x_2, y_2), (x_3, y_3), \dots, (x_n, y_n)$
- ▶ The plot has two axes:
 - ▶ x -axis is the horizontal axis at the bottom of the plot
 - ▶ y -axis is the vertical line on the left hand of the plot

Coordinate Geometry

- ▶ Pairs of numbers represent points:
 $(x_1, y_1), (x_2, y_2), (x_3, y_3), \dots, (x_n, y_n)$
- ▶ The plot has two axes:
 - ▶ x-axis is the horizontal axis at the bottom of the plot
 - ▶ y-axis is the vertical line on the left hand of the plot
- ▶ To plot some point (x_i, y_i) :
 - ▶ Move along the x-axis to the point x_i from zero
 - ▶ Move up or down to the position of y_i

Example

```
plot(x=2, y=3, xlim=c(0, 4), ylim=c(0,4), pch=19, cex=3  
      , xlab="X-axis", ylab="Y-axis", col="darkred")  
abline(h=c(0,1,2,3,4),v=c(0,1,2,3,4), lty=2, col="gray40")  
text(x=2.2, y=3.2, labels="(2, 3)", cex=1)
```



Lines

- ▶ The equation for a line will most often be written as:
 - ▶ $y = mx + b$
 - ▶ $y = a + bx$
 - ▶ $y = \beta_0 + \beta_1 x$

Lines

- ▶ The equation for a line will most often be written as:
 - ▶ $y = mx + b$
 - ▶ $y = a + bx$
 - ▶ $y = \beta_0 + \beta_1 x$
- ▶ Hardly a day will go by for the next two years when you do not encounter (implicitly or explicitly) an equation that looks like this.

Lines

- ▶ The equation for a line will most often be written as:
 - ▶ $y = mx + b$
 - ▶ $y = a + bx$
 - ▶ $y = \beta_0 + \beta_1x$
- ▶ Hardly a day will go by for the next two years when you do not encounter (implicitly or explicitly) an equation that looks like this.
- ▶ From any two points we can calculate the “slope” (e.g., m) and the “intercept.”

Lines

- ▶ The equation for a line will most often be written as:
 - ▶ $y = mx + b$
 - ▶ $y = a + bx$
 - ▶ $y = \beta_0 + \beta_1x$
- ▶ Hardly a day will go by for the next two years when you do not encounter (implicitly or explicitly) an equation that looks like this.
- ▶ From any two points we can calculate the “slope” (e.g., m) and the “intercept.”
- ▶ The slope is the ratio of the differences in the two y pairs and the two x pairs (i.e., rise over run)

Lines

- ▶ The equation for a line will most often be written as:
 - ▶ $y = mx + b$
 - ▶ $y = a + bx$
 - ▶ $y = \beta_0 + \beta_1 x$
- ▶ Hardly a day will go by for the next two years when you do not encounter (implicitly or explicitly) an equation that looks like this.
- ▶ From any two points we can calculate the “slope” (e.g., m) and the “intercept.”
- ▶ The slope is the ratio of the differences in the two y pairs and the two x pairs (i.e., rise over run)
- ▶ The intercept is the value of the function where $x = 0$. In other words, it is the point where the line hits the y -axis.

Example

$[2, 1], [3, 5]$

Example

$$[2, 1], [3, 5]$$

$$m = \frac{5-1}{3-2} = 4$$


Example

$[2, 1], [3, 5]$

$$m = \frac{5-1}{3-2} = 4$$

$$1 = 4(2) + b \iff b = 1 - 8 = -7$$

```
plot(x=c(2,3), y=c(1, 5), xlim=c(2, 3), ylim=c(1,5), pch=19  
      , xlab="X-axis", ylab="Y-axis", col="darkred")  
abline(a=-7, b=4)
```



Drill



1. Find the line that goes through the points $[-3, 2], [4, 20]$
2. Solve for x : $3x - 4 = \frac{2x}{2}$
3. Solve for x : $(1 - x^2) = -6$
4. What are the coordinates for the point on the line $y = 3.4x - 8$ where $x = 14$?
5. Solve for x : $5x - 7 = 0$
6. Solve for x : $-6x - 17 = 0$
7. Solve for x : $-6x - 17 < 0$

Sets

- ▶ **Set:** A set is any well defined collection of elements. If x is an element of S , $x \in S$.

Sets

- ▶ **Set:** A set is any well defined collection of elements. If x is an element of S , $x \in S$.
- ▶ Types of sets:
 - ▶ Countably finite: a set with a finite number of elements, which can be mapped onto positive integers. $S = \{1, 2, 3, 4, 5, 6\}$

Sets

- ▶ **Set:** A set is any well defined collection of elements. If x is an element of S , $x \in S$.
- ▶ Types of sets:
 - ▶ Countably finite: a set with a finite number of elements, which can be mapped onto positive integers. $S = \{1, 2, 3, 4, 5, 6\}$
 - ▶ Countably infinite: a set with an infinite number of elements, which can still be mapped onto positive integers.
 $S = \{1, \frac{1}{2}, \frac{1}{3}, \dots\}$


Sets

- ▶ **Set:** A set is any well defined collection of elements. If x is an element of S , $x \in S$.
- ▶ Types of sets:
 - ▶ Countably finite: a set with a finite number of elements, which can be mapped onto positive integers. $S = \{1, 2, 3, 4, 5, 6\}$
 - ▶ Countably infinite: a set with an infinite number of elements, which can still be mapped onto positive integers.
 $S = \{1, \frac{1}{2}, \frac{1}{3}, \dots\}$
 - ▶ Uncountably infinite: a set with an infinite number of elements, which cannot be mapped onto positive integers.
 $S = \{x : x \in [0, 1]\}$

Sets


- ▶ **Set:** A set is any well defined collection of elements. If x is an element of S , $x \in S$.
- ▶ Types of sets:
 - ▶ Countably finite: a set with a finite number of elements, which can be mapped onto positive integers. $S = \{1, 2, 3, 4, 5, 6\}$
 - ▶ Countably infinite: a set with an infinite number of elements, which can still be mapped onto positive integers.
 $S = \{1, \frac{1}{2}, \frac{1}{3}, \dots\}$
 - ▶ Uncountably infinite: a set with an infinite number of elements, which cannot be mapped onto positive integers.
 $S = \{x : x \in [0, 1]\}$
 - ▶ Empty: a set with no elements. $S = \{\}$ or $S = \emptyset$

Set operations




- **Union:** The union of two sets A and B , $A \cup B$, is the set containing all of the elements in A or B .

Set operations



- ▶ **Union:** The union of two sets A and B , $A \cup B$, is the set containing all of the elements in A or B .
- ▶ **Intersection:** The intersection of sets A and B , $A \cap B$, is the set containing all of the elements in both A and B .

Set operations



- ▶ **Union:** The union of two sets A and B , $A \cup B$, is the set containing all of the elements in A or B .
- ▶ **Intersection:** The intersection of sets A and B , $A \cap B$, is the set containing all of the elements in both A and B .
- ▶ **Complement:** If set A is a subset of S , then the complement of A , denoted A^C , is the set containing all of the elements in S that are not in A . Sometimes notated as \tilde{A} .

Drill!



Draw the following:

1. $A \cap B$
2. $\tilde{A} \cap C$
3. $(A \cap B) \cup (\tilde{A} \cap C)$
4. $(A \cap \tilde{B}) \cup (\tilde{A} \cap C)$
5. $(A \cap \tilde{B}) \cap (\tilde{A} \cup C)$

Note: There are many ways to denote the complement. Another common one would be A' .

6. Give examples of $A \cap \tilde{B}$ where A is the set of all vegetables and B is the set of green foods
7. Given $U = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$, $A = \{2, 4, 6, 8\}$, $B = \{1, 3, 4, 5, 7\}$, $C = \{7, 8\}$, find $A \cap B$.
8. \tilde{B} (in U)
9. $\tilde{A} \cup \tilde{B}$ (in U)
10. $(A \cap C) \cup (A \cap B)$

The Reals

- ▶ \mathbf{R}^1 is the set of all *real* numbers extending from $-\infty$ to $+\infty$ — i.e., the real number line.

The Reals

- ▶ \mathbf{R}^1 is the set of all *real* numbers extending from $-\infty$ to $+\infty$ — i.e., the real number line.
- ▶ \mathbf{R}^n is an n -dimensional space (often referred to as Euclidean space), where each of the n axes extends from $-\infty$ to $+\infty$.

The Reals

- ▶ \mathbf{R}^1 is the set of all *real* numbers extending from $-\infty$ to $+\infty$ — i.e., the real number line.
- ▶ \mathbf{R}^n is an n -dimensional space (often referred to as Euclidean space), where each of the n axes extends from $-\infty$ to $+\infty$.
- ▶ Examples:
- ▶ \mathbf{R}^1 is a line.
- ▶ \mathbf{R}^2 is a plane.

The Reals

- ▶ \mathbf{R}^1 is the set of all *real* numbers extending from $-\infty$ to $+\infty$ — i.e., the real number line.
- ▶ \mathbf{R}^n is an n -dimensional space (often referred to as Euclidean space), where each of the n axes extends from $-\infty$ to $+\infty$.
- ▶ Examples:
- ▶ \mathbf{R}^1 is a line.
- ▶ \mathbf{R}^2 is a plane.
- ▶ \mathbf{R}^3 is a 3-D space.
- ▶ \mathbf{R}^4 could be 3-D plus time.

The Reals

- ▶ \mathbf{R}^1 is the set of all *real* numbers extending from $-\infty$ to $+\infty$ — i.e., the real number line.
- ▶ \mathbf{R}^n is an n -dimensional space (often referred to as Euclidean space), where each of the n axes extends from $-\infty$ to $+\infty$.
- ▶ Examples:
- ▶ \mathbf{R}^1 is a line.
- ▶ \mathbf{R}^2 is a plane.
- ▶ \mathbf{R}^3 is a 3-D space.
- ▶ \mathbf{R}^4 could be 3-D plus time.
- ▶ Points in \mathbf{R}^n are ordered n -tuples, where each element of the n -tuple represents the coordinate along that dimension.

Interval notation

- ▶ Open interval:

$$(a, b) \equiv \{x \in \mathbf{R}^1 : a < x < b\}$$

Interval notation

- ▶ Open interval:

$$(a, b) \equiv \{x \in \mathbf{R}^1 : a < x < b\}$$

- ▶ Closed interval:

$$[a, b] \equiv \{x \in \mathbf{R}^1 : a \leq x \leq b\}$$

Interval notation

- ▶ Open interval:

$$(a, b) \equiv \{x \in \mathbf{R}^1 : a < x < b\}$$

- ▶ Closed interval:

$$[a, b] \equiv \{x \in \mathbf{R}^1 : a \leq x \leq b\}$$

- ▶ Half open, half closed:

$$(a, b] \equiv \{x \in \mathbf{R}^1 : a < x \leq b\}$$

Neighborhoods: Intervals, Disks, and Balls

In many areas of math, we need a formal construct for what it means to be “near” a point \mathbf{c} in \mathbf{R}^n .

Neighborhoods: Intervals, Disks, and Balls

In many areas of math, we need a formal construct for what it means to be “near” a point \mathbf{c} in \mathbf{R}^n . This is generally called the **neighborhood** of \mathbf{c} and is represented by an open interval, disk, or ball, depending on whether \mathbf{R}^n is of one, two, or more dimensions, respectively.

Neighborhoods: Intervals, Disks, and Balls

In many areas of math, we need a formal construct for what it means to be “near” a point \mathbf{c} in \mathbf{R}^n . This is generally called the **neighborhood** of \mathbf{c} and is represented by an open interval, disk, or ball, depending on whether \mathbf{R}^n is of one, two, or more dimensions, respectively. Given the point c , these are defined as:

- ▶ ϵ -interval in \mathbf{R}^1 : $\{x : |x - c| < \epsilon\}$
 - ▶ The open interval $(c - \epsilon, c + \epsilon)$.

Neighborhoods: Intervals, Disks, and Balls

In many areas of math, we need a formal construct for what it means to be “near” a point \mathbf{c} in \mathbf{R}^n . This is generally called the **neighborhood** of \mathbf{c} and is represented by an open interval, disk, or ball, depending on whether \mathbf{R}^n is of one, two, or more dimensions, respectively. Given the point c , these are defined as:

- ▶ ϵ -interval in \mathbf{R}^1 : $\{x : |x - c| < \epsilon\}$
 - ▶ The open interval $(c - \epsilon, c + \epsilon)$.
- ▶ ϵ -disk in \mathbf{R}^2 : $\{\mathbf{x} : \|\mathbf{x} - \mathbf{c}\| < \epsilon\}$
 - ▶ The open interior of the circle centered at \mathbf{c} with radius ϵ .

Neighborhoods: Intervals, Disks, and Balls

In many areas of math, we need a formal construct for what it means to be “near” a point \mathbf{c} in \mathbf{R}^n . This is generally called the **neighborhood** of \mathbf{c} and is represented by an open interval, disk, or ball, depending on whether \mathbf{R}^n is of one, two, or more dimensions, respectively. Given the point c , these are defined as:

- ▶ ϵ -interval in \mathbf{R}^1 : $\{x : |x - c| < \epsilon\}$
 - ▶ The open interval $(c - \epsilon, c + \epsilon)$.
- ▶ ϵ -disk in \mathbf{R}^2 : $\{\mathbf{x} : \|\mathbf{x} - \mathbf{c}\| < \epsilon\}$
 - ▶ The open interior of the circle centered at \mathbf{c} with radius ϵ .
- ▶ ϵ -ball in \mathbf{R}^n : $\{\mathbf{x} : \|\mathbf{x} - \mathbf{c}\| < \epsilon\}$
 - ▶ The open interior of the sphere centered at \mathbf{c} with radius ϵ .

Advanced set vocabulary

- ▶ **Interior Point:** The point \mathbf{x} is an interior point of the set S if \mathbf{x} is in S and if there is some ϵ -ball around \mathbf{x} that contains only points in S .

Advanced set vocabulary

- ▶ **Interior Point:** The point \mathbf{x} is an interior point of the set S if \mathbf{x} is in S and if there is some ϵ -ball around \mathbf{x} that contains only points in S . The **interior** of S is the collection of all interior points in S .

Advanced set vocabulary

- **Interior Point:** The point \mathbf{x} is an interior point of the set S if \mathbf{x} is in S and if there is some ϵ -ball around \mathbf{x} that contains only points in S . The **interior** of S is the collection of all interior points in S .

Example: The interior of the set $\{(x, y) : x^2 + y^2 \leq 4\}$ is $\{(x, y) : x^2 + y^2 < 4\}$.

- ▶ **Boundary Point:** The point \mathbf{x} is a boundary point of the set S if every ϵ -ball around \mathbf{x} contains both points that are in S and points that are outside S .

- ▶ **Boundary Point:** The point \mathbf{x} is a boundary point of the set S if every ϵ -ball around \mathbf{x} contains both points that are in S and points that are outside S . The **boundary** is the collection of all boundary points.

- ▶ **Boundary Point:** The point \mathbf{x} is a boundary point of the set S if every ϵ -ball around \mathbf{x} contains both points that are in S and points that are outside S . The **boundary** is the collection of all boundary points.

Example: The boundary of $\{(x, y) : x^2 + y^2 \leq 4\}$ is $\{(x, y) : x^2 + y^2 = 4\}$.

- ▶ **Open:** A set S is open if for each point \mathbf{x} in S , there exists an open ϵ -ball around \mathbf{x} completely contained in S .

Example: $\{(x, y) : x^2 + y^2 < 4\}$

- ▶ **Open:** A set S is open if for each point \mathbf{x} in S , there exists an open ϵ -ball around \mathbf{x} completely contained in S .

Example: $\{(x, y) : x^2 + y^2 < 4\}$

- ▶ **Closed:** A set S is closed if it contains all of its boundary points.

Example: $\{(x, y) : x^2 + y^2 \leq 4\}$

- ▶ **Open:** A set S is open if for each point \mathbf{x} in S , there exists an open ϵ -ball around \mathbf{x} completely contained in S .

Example: $\{(x, y) : x^2 + y^2 < 4\}$

- ▶ **Closed:** A set S is closed if it contains all of its boundary points.

Example: $\{(x, y) : x^2 + y^2 \leq 4\}$

- ▶ Note: a set may be neither open nor closed.

Example: $\{(x, y) : 2 < x^2 + y^2 \leq 4\}$

- **Complement:** The complement of set S is everything outside of S .

Example: The complement of $\{(x, y) : x^2 + y^2 \leq 4\}$ is $\{(x, y) : x^2 + y^2 > 4\}$.

- ▶ **Complement:** The complement of set S is everything outside of S .

Example: The complement of $\{(x, y) : x^2 + y^2 \leq 4\}$ is $\{(x, y) : x^2 + y^2 > 4\}$.

- ▶ **Closure:** The closure of set S is the smallest closed set that contains S .

Example: The closure of $\{(x, y) : x^2 + y^2 < 4\}$ is $\{(x, y) : x^2 + y^2 \leq 4\}$

- ▶ **Complement:** The complement of set S is everything outside of S .

Example: The complement of $\{(x, y) : x^2 + y^2 \leq 4\}$ is $\{(x, y) : x^2 + y^2 > 4\}$.

- ▶ **Closure:** The closure of set S is the smallest closed set that contains S .

Example: The closure of $\{(x, y) : x^2 + y^2 < 4\}$ is $\{(x, y) : x^2 + y^2 \leq 4\}$

- ▶ **Bounded:** A set S is bounded if it can be contained within an ϵ -ball.

- ▶ **Bounded:** A set S is bounded if it can be contained within an ϵ -ball.

Examples:

Bounded: any interval that doesn't have ∞ or $-\infty$ as endpoints;
any disk in a plane with finite radius.

Unbounded: the set of integers in \mathbf{R}^1 .

- ▶ **Bounded:** A set S is bounded if it can be contained within an ϵ -ball.

Examples:

Bounded: any interval that doesn't have ∞ or $-\infty$ as endpoints;
any disk in a plane with finite radius.

Unbounded: the set of integers in \mathbf{R}^1 .

- ▶ **Compact:** A set is compact if and only if it is both closed and bounded.

DRILL!



Create a graph for the following sets:

1. $(2, 6]$

2. $[-2, 3) \cup [5, \infty)$

Express the following in plain English (note that \mathbf{Z} is the set of all integers):

3. $[5, 7)$

4. $(3, \infty)$

5. $\{x \in \mathbf{Z} : x \geq 2\}$

6. $\{y \in \mathbf{R} : y \in \mathbf{Z}\}$

Express the following in math notation:

7. The set of real numbers greater than or equal to -5 and less than 4 .
8. The set of real numbers greater than 12 .
9. The set of real numbers that are divisible by 3 (where when I divide by 3 I will get an integer).
10. The set of numbers that solve the equation $y = 4x + 24$

Introducing functions

- ▶ A **function** (in \mathbf{R}^1) is a rule or relationship or mapping or transformation that assigns one and only one number in \mathbf{R}^1 to each number in \mathbf{R}^1 .

Introducing functions

- ▶ A **function** (in \mathbf{R}^1) is a rule or relationship or mapping or transformation that assigns one and only one number in \mathbf{R}^1 to each number in \mathbf{R}^1 . Example?

Introducing functions

- ▶ A **function** (in \mathbf{R}^1) is a rule or relationship or mapping or transformation that assigns one and only one number in \mathbf{R}^1 to each number in \mathbf{R}^1 . Example?
- ▶ Mapping notation examples
 - ▶ Function of one variable: $f : \mathbf{R}^1 \rightarrow \mathbf{R}^1$
 - ▶ Function of two variables: $f : \mathbf{R}^2 \rightarrow \mathbf{R}^1$

Introducing functions

- ▶ A **function** (in \mathbf{R}^1) is a rule or relationship or mapping or transformation that assigns one and only one number in \mathbf{R}^1 to each number in \mathbf{R}^1 . Example?
- ▶ Mapping notation examples
 - ▶ Function of one variable: $f : \mathbf{R}^1 \rightarrow \mathbf{R}^1$
 - ▶ Function of two variables: $f : \mathbf{R}^2 \rightarrow \mathbf{R}^1$

Introducing functions

- ▶ A **function** (in \mathbf{R}^1) is a rule or relationship or mapping or transformation that assigns one and only one number in \mathbf{R}^1 to each number in \mathbf{R}^1 . Example?
- ▶ Mapping notation examples
 - ▶ Function of one variable: $f : \mathbf{R}^1 \rightarrow \mathbf{R}^1$
 - ▶ Function of two variables: $f : \mathbf{R}^2 \rightarrow \mathbf{R}^1$
- ▶ Examples:



$$f(x) = x + 1$$

For each x in \mathbf{R}^1 , $f(x)$ assigns the number $x + 1$.

Introducing functions

- ▶ A **function** (in \mathbf{R}^1) is a rule or relationship or mapping or transformation that assigns one and only one number in \mathbf{R}^1 to each number in \mathbf{R}^1 . Example?
- ▶ Mapping notation examples

- ▶ Function of one variable: $f : \mathbf{R}^1 \rightarrow \mathbf{R}^1$
- ▶ Function of two variables: $f : \mathbf{R}^2 \rightarrow \mathbf{R}^1$

- ▶ Examples:



$$f(x) = x + 1$$

For each x in \mathbf{R}^1 , $f(x)$ assigns the number $x + 1$.



$$f(x, y) = x^2 + y^2$$

For each ordered pair (x, y) in \mathbf{R}^2 , $f(x, y)$ assigns the number $x^2 + y^2$.

- ▶ Often use one variable x as input and another y as output.

- ▶ Example:

$$y = x + 1$$

- ▶ Input variable also called **independent** variable.

- ▶ Often use one variable x as input and another y as output.

- ▶ Example:

$$y = x + 1$$

- ▶ Input variable also called **independent** variable.
 - ▶ Output variable also called **dependent** variable.

Domain and range

- ▶ Some functions are defined only on proper subsets of \mathbf{R}^n .
- ▶ **Domain:** the set of numbers in X at which $f(x)$ is defined.

Domain and range

- ▶ Some functions are defined only on proper subsets of \mathbf{R}^n .
- ▶ **Domain:** the set of numbers in X at which $f(x)$ is defined.
- ▶ **Range:** elements of Y assigned by $f(x)$ to elements of X , or

$$f(X) = \{y : y = f(x), x \in X\}$$

Domain and range

- ▶ Some functions are defined only on proper subsets of \mathbf{R}^n .
- ▶ **Domain:** the set of numbers in X at which $f(x)$ is defined.
- ▶ **Range:** elements of Y assigned by $f(x)$ to elements of X , or

$$f(X) = \{y : y = f(x), x \in X\}$$

- ▶ Most often used when talking about a function $f : \mathbf{R}^1 \rightarrow \mathbf{R}^1$.

Example 1

$$f(x) = \begin{cases} x + 1, & 1 \leq x \leq 2 \\ 0, & x = 0 \\ 1 - x & -2 \leq x \leq -1 \end{cases}$$

Domain

$$X = [-2, -1] \cup \{0\} \cup [1, 2]$$

Range

$$f(X) = [2, 3] \cup \{0\}$$

Example 2

$$f(x) = 1/x$$

Domain

Example 2

$$f(x) = 1/x$$

Domain

$$X = \mathbf{R}^1 - \{0\}$$

Example 2

$$f(x) = 1/x$$

Domain

$$X = \mathbf{R}^1 - \{0\}$$

Range

Example 2

$$f(x) = 1/x$$

Domain

$$X = \mathbf{R}^1 - \{0\}$$

Range

$$f(X) = \mathbf{R}^1 - \{0\}$$

Example 3

$$f(x, y) = x^2 + y^2$$

Domain

$$X = \mathbf{R}^2$$

Range

$$f(X, Y) = \mathbf{R}_{+}^1$$

DRILL!



Create a graph for the following functions (rough it out):

1. $f(x) = 3 + 2x$

2. $f(x) = 2x^2 + 3x - 4$

3. Consider the following function:

$$h(x) = \frac{\ln(x - 3)}{\sqrt{5 - x}}$$

In set notation express the domain of $h(x)$.

4. For what value of x is $h(x) = 0$?
5. Does the range of $h(x)$ span all of \mathbf{R}^1 ?

Function vocabulary

► **Monomials:** $f(x) = ax^k$; a is the *coefficient*. k is the *degree*.

Examples: $y = x^2$, $y = -\frac{1}{2}x^3$

Function vocabulary

- **Monomials:** $f(x) = ax^k$; a is the *coefficient*. k is the *degree*.

Examples: $y = x^2$, $y = -\frac{1}{2}x^3$

- **Polynomials:** sum of monomials.

Examples: $y = -\frac{1}{2}x^3 + x^2$, $y = 3x + 5$

The *degree* of a polynomial is the highest degree of its monomial terms.

Function vocabulary

- **Monomials:** $f(x) = ax^k$; a is the *coefficient*. k is the *degree*.


Examples: $y = x^2$, $y = -\frac{1}{2}x^3$

- **Polynomials:** sum of monomials.

Examples: $y = -\frac{1}{2}x^3 + x^2$, $y = 3x + 5$


The *degree* of a polynomial is the highest degree of its monomial terms. Also, it's often a good idea to write polynomials with terms in decreasing degree.

```
x<-seq(-4, 4, by=.1); y<-.5*x^3+x^2  
plot(x, y, xlab="x", ylab="f(x)", type="l")
```




► Exponential Functions: Example: $y = 2^x$

```
x<-seq(-4, 4, by=.1); y<-2^x  
plot(x, y, xlab="x", ylab="f(x)", type="l")
```



► Trigonometric Functions: Examples: $y = \cos(x)$, $y = 3 \sin(4x)$

```
x<-seq(-4, 4, by=.1); y<-3*sin(4*x)
plot(x, y, xlab="x", ylab="f(x)", type="l")
```



► **Linear:** polynomial of degree 1.

Example: $y = mx + b$, where m is the slope and b is the y -intercept.

- ▶ **Linear:** polynomial of degree 1.

Example: $y = mx + b$, where m is the slope and b is the y -intercept.

- ▶ **Nonlinear:** anything that isn't constant or polynomial of degree 1.

Examples: $y = x^2 + 2x + 1$, $y = \sin(x)$, $y = \ln(x)$, $y = e^x$

- ▶ **Quadratic:** a second degree polynomial function.

Example: $f(x) = ax^2 + bx + c$

- ▶ **Linear:** polynomial of degree 1.

Example: $y = mx + b$, where m is the slope and b is the y -intercept.

- ▶ **Nonlinear:** anything that isn't constant or polynomial of degree 1.

Examples: $y = x^2 + 2x + 1$, $y = \sin(x)$, $y = \ln(x)$, $y = e^x$

- ▶ **Quadratic:** a second degree polynomial function.

Example: $f(x) = ax^2 + bx + c$

- ▶ Always, always, always, graph your function.

Inverse Functions

- Sometimes we're given a function $y = f(x)$ and we want to find how x varies as a function of y . That is, we are trying to "solve for" x .

Inverse Functions

- ▶ Sometimes we're given a function $y = f(x)$ and we want to find how x varies as a function of y . That is, we are trying to "solve for" x .
- ▶ If f is a one-to-one mapping, then it has an inverse. This will usually be denoted as $f^{-1}(x)$. That means that $f^{-1}(f(x)) = x$.

Inverse Functions

- ▶ Sometimes we're given a function $y = f(x)$ and we want to find how x varies as a function of y . That is, we are trying to "solve for" x .
- ▶ If f is a one-to-one mapping, then it has an inverse. This will usually be denoted as $f^{-1}(x)$. That means that $f^{-1}(f(x)) = x$.
- ▶ A good way to do this is to use algebra to isolate x (your independent variable in $f(x)$) on one side of the equation.

Example 1: Find the inverse functions

$$f(x) = 3x + 2$$

Step 1: Solve for x

$$y = 3x + 2$$

$$y - 2 = 3x$$

Example 1: Find the inverse functions

$$f(x) = 3x + 2$$

Step 1: Solve for x

$$y = 3x + 2$$

$$y - 2 = 3x$$

$$x = \frac{1}{3}(y - 2)$$

Example 1: Find the inverse functions

$$f(x) = 3x + 2$$

Step 1: Solve for x

$$y = 3x + 2$$

$$y - 2 = 3x$$

$$x = \frac{1}{3}(y - 2)$$

Step 2: Identify the inverse function (note the x)

$$f^{-1}(x) = \frac{1}{3}(x - 2)$$

Example 2: Find the inverse of the function

$$f(x) = 3x - 4z + 2$$

Example 2: Find the inverse of the function

$$f(x) = 3x - 4z + 2$$

Step 1: Solve for x

$$y + 4z - 2 = 3x$$

Example 2: Find the inverse of the function

$$f(x) = 3x - 4z + 2$$

Step 1: Solve for x

$$y + 4z - 2 = 3x$$

$$x = \frac{1}{3}(y + 4z - 2)$$

Example 2: Find the inverse of the function

$$f(x) = 3x - 4z + 2$$

Step 1: Solve for x

$$y + 4z - 2 = 3x$$

$$x = \frac{1}{3}(y + 4z - 2)$$

Step 2: Identify the inverse function (note the x)

$$f^{-1}(x) = \frac{1}{3}(x + 4z - 2)$$

For many functions, the inverse will not exist

- ▶ Sometimes it's impossible to figure out what x went into the function just from the output $f(x)$.

For many functions, the inverse will not exist

- ▶ Sometimes it's impossible to figure out what x went into the function just from the output $f(x)$.
- ▶ An undefined inverse is especially likely in non-linear functions.

For many functions, the inverse will not exist

- ▶ Sometimes it's impossible to figure out what x went into the function just from the output $f(x)$.
- ▶ An undefined inverse is especially likely in non-linear functions.
- ▶ Imagine we have the function $f(x) = x^2$ (a parabola).
- ▶ Solving for x , we get $x = \sqrt{y}$ and $x = -\sqrt{y}$ — for each value of y , there are two values of x .

Roots

- ▶ You are going to be spending a lot of time finding **roots** of functions: those values where $f(x) = 0$.
 - ▶ Decision theory/game theory
 - ▶ Dynamic systems
 - ▶ Maximum likelihood

Roots

- ▶ You are going to be spending a lot of time finding **roots** of functions: those values where $f(x) = 0$.
 - ▶ Decision theory/game theory
 - ▶ Dynamic systems
 - ▶ Maximum likelihood
- ▶ Procedure:
 - ▶ Given $y = f(x)$
 - ▶ set $y = 0$.
 - ▶ Solve for x .

Finding the y-intercepts

- ▶ You already found the root of one function when you calculated the **y-intercept** of a line.
- ▶ Where does the line $f(x) = a + bx$ cross the y-axis?

$$a + bx = 0$$

$$a = -bx$$

$$x = -\frac{a}{b}$$

The quadratic equation

- ▶ What is the root of $f(x) = ax^2 + bx + c$

The quadratic equation

- ▶ What is the root of $f(x) = ax^2 + bx + c$
- ▶ The **quadratic equation** is the solution to this question that most of us have been forced to memorize at some point.

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- ▶ These numbers are the two roots of $f(x)$.

Factoring

- ▶ Many times it is possible to **factor** a function into two components multiplied together.
- ▶ This makes finding the root simple

Factoring

- ▶ Many times it is possible to **factor** a function into two components multiplied together.
- ▶ This makes finding the root simple
- ▶ Example

$$f(x) = x^2 + 3x - 4 = 0$$

$$(x - 1)(x + 4) = 0$$

$$x = \{1, -4\}$$

- ▶ **FOIL** (First **O**utside **I**nside **L**ast)
- ▶ The middle term (e.g., $3x$) is the sum of the constants. The final term is the product.

Example

$$x(x - 1) = 6$$

$$x^2 - x - 6 = 0$$

$$(x - 3)(x + 2) = 0$$

Solution are $x = -2$ and $x = 3$.

Completing the square

- ▶ The basic intuition is to solve a quadratic by adding and subtracting the same number to make the factors more recognizable.

Completing the square

- ▶ The basic intuition is to solve a quadratic by adding and subtracting the same number to make the factors more recognizable.

► **Example**

$$x^2 - 6x + 5 = 0$$

► **Example**

$$x^2 - 6x + 5 = 0$$

$$x^2 - 6x + 5 + 4 - 4 = 0$$

► **Example**

$$x^2 - 6x + 5 = 0$$

$$x^2 - 6x + 5 + 4 - 4 = 0$$

$$(x^2 - 6x + 9) - 4 = 0$$

$$(x - 3)^2 - 4 = 0$$

► **Example**

$$x^2 - 6x + 5 = 0$$

$$x^2 - 6x + 5 + 4 - 4 = 0$$

$$(x^2 - 6x + 9) - 4 = 0$$

$$(x - 3)^2 - 4 = 0$$

$$(x - 3)^2 = 4$$

► **Example**

$$x^2 - 6x + 5 = 0$$

$$x^2 - 6x + 5 + 4 - 4 = 0$$

$$(x^2 - 6x + 9) - 4 = 0$$

$$(x - 3)^2 - 4 = 0$$

$$(x - 3)^2 = 4$$

$$x - 3 = \pm 2$$

► **Example**

$$x^2 - 6x + 5 = 0$$

$$x^2 - 6x + 5 + 4 - 4 = 0$$

$$(x^2 - 6x + 9) - 4 = 0$$

$$(x - 3)^2 - 4 = 0$$

$$(x - 3)^2 = 4$$

$$x - 3 = \pm 2$$

$$x = \{5, 1\}$$

► **Example**

$$x^2 - 6x + 5 = 0$$

$$x^2 - 6x + 5 + 4 - 4 = 0$$

$$(x^2 - 6x + 9) - 4 = 0$$

$$(x - 3)^2 - 4 = 0$$

$$(x - 3)^2 = 4$$

$$x - 3 = \pm 2$$

$$x = \{5, 1\}$$

Your turn

Solve this formula by completing the square

$$x^2 + 8x + 6 = 0$$

DRILL! (The pain means it is working)



1. Factor $-7\theta^2 + 21\theta - 14$
2. FOIL: $(2x - 3)(5x + 7)$
3. Factor: $q^2 - 10q + 9$
4. Factor and reduce: $\frac{\beta - \alpha}{\alpha^2 - \beta^2}$
5. Solve: $15\delta + 45 - 5\delta = 36$
6. Solve: $0.30\Omega + 0.05 = 0.25$
7. Solve: $-4x^2 + 64 = 8x - 32$
8. Complete the square and solve: $x^2 + 14x - 14 = 0$
9. Complete the square and solve: $\frac{1}{3}y^2 + \frac{2}{3}y - 16 = 0$
10. Solve using the quadratic formula: $2x^2 + 5x - 7$

Solve the following formulas:

11. $5 + 11x = -3x^2$

12. $\sqrt{4x + 13} = x + 2$

13. $10^{3x^2} 10^x = 100$

14. $6x^2 - 6x - 6 = 0$

15. $5 + 11x = -3x^2$

16. Find the inverse of $f(x) = 5x - 2$
17. Simplify $h(x) = g(f(x))$, where $f(x) = x^2 + 2$ and $g(x) = \sqrt{x - 4}$.
18. Simplify $h(x) = f(g(x))$ with the same f and g . Is it the same as before?
19. Rewrite the following by taking the log of both sides. Is the result a linear function?

$$y = \alpha \times x_1^{\beta_1} \times \beta_2 x_2 \times \beta_3 x_3$$

20. Rewrite the following by taking the log of both sides. Is the result a linear function?

$$y = \alpha \times x_1^{\beta_1} \times \frac{x_2^{\beta_2}}{x_3^{\beta_3}}$$