

Derivatives: Part 1

David Carlson

2021

Derivatives: Part 1

Encounters with basic concepts from calculus will be a daily event in your methods training. In the case of derivatives, you will primarily be using them to:

- ▶ Find the maximum and minimum values of specific functions including utility and likelihood functions;
- ▶ Derive a probability density function given its cumulative density function

This probably doesn't mean much to you at the moment. But what they have in common is that we are trying to understand a certain property of a function – the rate of change.

This probably doesn't mean much to you at the moment. But what they have in common is that we are trying to understand a certain property of a function – the rate of change.

- ▶ Usually we want to find when the rate of change is zero, as that indicates either a maximum or minimum.

This probably doesn't mean much to you at the moment. But what they have in common is that we are trying to understand a certain property of a function – the rate of change.

- ▶ Usually we want to find when the rate of change is zero, as that indicates either a maximum or minimum.
- ▶ These concepts were developed in the 17th century to understand physical motion.
- ▶ If a function describes the location of a baseball flying through the air, the first derivative describes the velocity of the ball, and the second derivative represents the acceleration.

Defining derivatives

- ▶ The derivative of f at x is its rate of change at x — i.e., how much $f(x)$ changes with a change in x .
 - ▶ For a line, the derivative is the slope.
 - ▶ For a curve, the derivative is the tangent line at x .

Defining derivatives

- ▶ The derivative of f at x is its rate of change at x — i.e., how much $f(x)$ changes with a change in x .
 - ▶ For a line, the derivative is the slope.
 - ▶ For a curve, the derivative is the tangent line at x .
- ▶ More formally, let f be a function whose domain includes an open interval containing the point x . The derivative of f at x is given by

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{(x+h) - x} \\&= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}\end{aligned}$$

Discussion

- ▶ Don't succumb to the temptation to take this equation as a given without any understanding. This idea is over 400 years old now, and is not beyond your abilities to understand.

Discussion

- ▶ Don't succumb to the temptation to take this equation as a given without any understanding. This idea is over 400 years old now, and is not beyond your abilities to understand.
- ▶ Imagine that you want to calculate how a function changes between time period 1 and period 2. With a little thought we would calculate this as:

$$m = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

- ▶ This is same “rise over run” calculation we used to calculate the slope of a line from before.

Derivative (Version 1)

- ▶ What we want to calculate is the “instantaneous” rate of change.
- ▶ That is, we want to know the rate of change as x_1 and x_2 get closer and closer together such that time period between when we measure them moves towards zero.

Derivative (Version 1)

- ▶ What we want to calculate is the “instantaneous” rate of change.
- ▶ That is, we want to know the rate of change as x_1 and x_2 get closer and closer together such that time period between when we measure them moves towards zero. This is the conceptual leap that it takes a genius like Newton to discover, but which we can accept with a bit of thought.
- ▶ So one way to think about this is:

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

Derivative (Version 2)

- ▶ We could also think of it this way

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

- ▶ This is just a re-parameterized version of the same thing.

Examples: Solve using version 1

$$f(x) = 3x^2 + 4x - 6$$

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

Examples: Solve using version 2

$$f(x) = 3x^2 + 4x - 6$$

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Note on notation

For historical reasons, there are multiple notations for derivatives.

- ▶ (Prime or Lagrange Notation)
- ▶ (Leibniz's Notation)

Using either version, find the derivatives for the following functions. I need one person to volunteer to draw each function and its derivative.

1. $f(x) = c$

2. $f(x) = 2x$

3. $f(x) = x^2$

4. $f(x) = x^3$

Important concepts for derivatives

- ▶ If $f'(x)$ exists at a point x , then f is said to be **differentiable** at x .

Important concepts for derivatives

- ▶ If $f'(x)$ exists at a point x , then f is said to be **differentiable** at x .
- ▶ Similarly, if $f'(x)$ exists for every point along an interval, then f is differentiable along that interval.

Important concepts for derivatives

- ▶ If $f'(x)$ exists at a point x , then f is said to be **differentiable** at x .
- ▶ Similarly, if $f'(x)$ exists for every point along an interval, then f is differentiable along that interval.
- ▶ For f to be differentiable at x , f must be both continuous and “smooth” at x .

Important concepts for derivatives

- ▶ If $f'(x)$ exists at a point x , then f is said to be **differentiable** at x .
- ▶ Similarly, if $f'(x)$ exists for every point along an interval, then f is differentiable along that interval.
- ▶ For f to be differentiable at x , f must be both continuous and “smooth” at x .
- ▶ The process of calculating $f'(x)$ is called **differentiation**.

Important concepts for derivatives

- ▶ If $f'(x)$ exists at a point x , then f is said to be **differentiable** at x .
- ▶ Similarly, if $f'(x)$ exists for every point along an interval, then f is differentiable along that interval.
- ▶ For f to be differentiable at x , f must be both continuous and “smooth” at x .
- ▶ The process of calculating $f'(x)$ is called **differentiation**.
- ▶ A function is *monotonically increasing* in a domain if it has a positive derivative over that domain. Likewise, it is *monotonically decreasing* if it has a negative derivative.

Existence

- ▶ $f'(x)$ at some particular value of x exists iff $f(x)$ is continuous at that value of x , and there is no point where the right-hand derivative and the left-hand derivative are different.

Existence

- ▶ $f'(x)$ at some particular value of x exists iff $f(x)$ is continuous at that value of x , and there is no point where the right-hand derivative and the left-hand derivative are different.
- ▶ $f'(x)$ exists for all values of x iff $f(x)$ is differentiable at all values of x in the domain of $f(x)$.

Existence

- ▶ $f'(x)$ at some particular value of x exists iff $f(x)$ is continuous at that value of x , and there is no point where the right-hand derivative and the left-hand derivative are different.
- ▶ $f'(x)$ exists for all values of x iff $f(x)$ is differentiable at all values of x in the domain of $f(x)$.
- ▶ The basic test is “can you draw the function without lifting your pencil.” But there are exceptions.

Existence

- ▶ $f'(x)$ at some particular value of x exists iff $f(x)$ is continuous at that value of x , and there is no point where the right-hand derivative and the left-hand derivative are different.
- ▶ $f'(x)$ exists for all values of x iff $f(x)$ is differentiable at all values of x in the domain of $f(x)$.
- ▶ The basic test is “can you draw the function without lifting your pencil.” But there are exceptions.

- ▶ The classic case where this “pencil rule” does not work are:



$$f(x) = |x|$$



$$f(x) = \frac{1}{x-1}$$



$$f(x) = 1, x \in [0, 1]$$

Determine whether or not the following functions are differentiable. If not, at what value of x do they fail?

1.

$$f(x) = |x|$$

2.

$$f(x) = \begin{cases} x^2 & x \leq 0 \\ -x & x < 0 \\ 0 & x = 0 \end{cases}$$

3.

$$f(x) = \sqrt{x}$$

4.

$$f(x) = x^{1/3}$$

5.

$$f(x) = 1/x$$

Properties of derivatives

Suppose that f and g are differentiable at x and that α is a constant. Then the functions $f \pm g$, αf , fg , and f/g (provided $g(x) \neq 0$) are also differentiable at x .

Rules: Making derivatives easier

Doing this calculation for every function is cumbersome, and sometimes difficult. Fortunately, hundreds of mathematical graduate students and professors have developed several “rules” that help us calculate derivatives quickly. All of these rules come with proofs, but for our purposes we usually just memorize them and use them. The first (and easiest) is that the derivative of a constant is zero.

The rules

Suppose that f and g are differentiable at x and that α is a constant.

► **Power rule:** $[x^k]' = kx^{k-1}$

The rules

Suppose that f and g are differentiable at x and that α is a constant.

- ▶ **Power rule:** $[x^k]' = kx^{k-1}$
- ▶ **Sum rule:** $[f(x) \pm g(x)]' = f'(x) \pm g'(x)$

The rules

Suppose that f and g are differentiable at x and that α is a constant.

- ▶ **Power rule:** $[x^k]' = kx^{k-1}$
- ▶ **Sum rule:** $[f(x) \pm g(x)]' = f'(x) \pm g'(x)$
- ▶ **Constant rule:** $[\alpha f(x)]' = \alpha f'(x)$

The rules

Suppose that f and g are differentiable at x and that α is a constant.

- ▶ **Power rule:** $[x^k]' = kx^{k-1}$
- ▶ **Sum rule:** $[f(x) \pm g(x)]' = f'(x) \pm g'(x)$
- ▶ **Constant rule:** $[\alpha f(x)]' = \alpha f'(x)$
- ▶ **Product rule:** $[f(x)g(x)]' = f'(x)g(x) + f(x)g'(x)$

The rules

Suppose that f and g are differentiable at x and that α is a constant.

- ▶ **Power rule:** $[x^k]' = kx^{k-1}$
- ▶ **Sum rule:** $[f(x) \pm g(x)]' = f'(x) \pm g'(x)$
- ▶ **Constant rule:** $[\alpha f(x)]' = \alpha f'(x)$
- ▶ **Product rule:** $[f(x)g(x)]' = f'(x)g(x) + f(x)g'(x)$
- ▶ **Quotient rule:** $[f(x)/g(x)]' = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}, g(x) \neq 0$

Examples

$$f(x) = 3x^2 + 2x^{1/3}$$

Examples

$$f(x) = 3x^2 + 2x^{1/3}$$

$$f'(x) = (3x^2)' + (2x^{1/3})'$$

Examples

$$f(x) = 3x^2 + 2x^{1/3}$$

$$f'(x) = (3x^2)' + (2x^{1/3})'$$

$$f'(x) = 3(2x) + 2(\frac{1}{3}x^{-2/3})$$

Examples

$$f(x) = 3x^2 + 2x^{1/3}$$

$$f'(x) = (3x^2)' + (2x^{1/3})'$$

$$f'(x) = 3(2x) + 2(\frac{1}{3}x^{-2/3})$$

$$= 6x + \frac{2}{3}x^{-2/3}$$

Examples

$$f(x) = 3x^2 + 2x^{1/3}$$

$$f'(x) = (3x^2)' + (2x^{1/3})'$$

$$f'(x) = 3(2x) + 2(\frac{1}{3}x^{-2/3})$$

$$= 6x + \frac{2}{3}x^{-2/3}$$

$$f(x) = (x^3)(2x^4)$$

$$f(x) = (x^3)(2x^4)$$

$$f'(x) = (x^3)'(2x^4) + (x^3)(2x^4)'$$

$$f(x) = (x^3)(2x^4)$$

$$f'(x) = (x^3)'(2x^4) + (x^3)(2x^4)'$$

$$= (3x^2)(2x^4) + (x^3)(8x^3)$$

$$f(x) = (x^3)(2x^4)$$

$$f'(x) = (x^3)'(2x^4) + (x^3)(2x^4)'$$

$$= (3x^2)(2x^4) + (x^3)(8x^3)$$

$$= 6x^6 + 8x^6 = 14x^6$$

$$f(x) = (x^3)(2x^4)$$

$$f'(x) = (x^3)'(2x^4) + (x^3)(2x^4)'$$

$$= (3x^2)(2x^4) + (x^3)(8x^3)$$

$$= 6x^6 + 8x^6 = 14x^6$$

$$f'(x) = (2x^7)' = 14x^6$$

$$f(x) = (x^3)(2x^4)$$

$$f'(x) = (x^3)'(2x^4) + (x^3)(2x^4)'$$

$$= (3x^2)(2x^4) + (x^3)(8x^3)$$

$$= 6x^6 + 8x^6 = 14x^6$$

$$f'(x) = (2x^7)' = 14x^6$$

Derivatives of exp

The derivative of the exp is worth just memorizing. This will make a bit more sense when we cover the chain rule, but for now:

- ▶ $\frac{d}{dx} \alpha e^x = \alpha e^x$
- ▶ $\frac{d^n}{dx^n} \alpha e^x = \alpha e^x$

Derivatives of exponential functions with other bases

$$\frac{d}{dx}(a^x) = (\ln(a)) a^x$$

Derivatives of the natural log

Same thing for the $\ln()$ function

$$\blacktriangleright \frac{d}{dx} \ln x = \frac{1}{x}$$

$$\blacktriangleright \frac{d}{dx} \ln x^k = \frac{d}{dx} k \ln x = \frac{k}{x}$$

Find the derivatives for the following functions:

1. $f(x) = 3x^{1/3}$
2. $f(t) = 14t - 7$
3. $f(y) = y^3 + 3y^2 - 12$
4. $f(x) = \frac{1}{100}x^{25} - \frac{1}{10}x^{0.25}$
5. $f(x) = (x^2 + 1)(x^3 - 1)$
6. $f(y) = (1 - 1/y^2)$
7. $f(y) = (y^3 - 7)(1 - 1/y^2)$
8. $f(x) = \ln(2\pi x^2)$
9. $f(y) = (y - y^{-1})(y - y^{-2})$
10. $f(x) = x^6 + 5x^5 - 2x^2 + 8$

11. $g(y) = 3e^y - \sqrt{y}$

12. $h(z) = \ln(z) + 1/z + 3^z$

13. $f(x) = (x + 3)^7(3x^4 - 2x^2 - 8)$

For these functions, find the derivative at $x = 1$ and $x = 3$

14. $f(x) = 2x^2 + 7$

15. $f(x) = x^3 - x + 1$

Find the derivatives for the following functions:

16. $y = 27x^3 + 5x^2 - x + 13$

17. $y = 81x^2 + 10x - 1$

18. $y = 162x + 10$