

The Equilibrium Reading of the Radial Acceleration Relation: What Is Derived, What Is Measured, What Is Dead

Carl P. Zimmerman (Briar Creek Tech) — AI-assisted research programme; not peer reviewed

2026-09-15 (v3; v2 2026-09-15; v1 2026-09-13)

A candidate reading of the relation, stated with every rung labelled, the audit that relabelled three of them, every dead branch recorded, and the tests that decide it. *Version 2 adds Section 8, an audit of the closure claims the same track made in the day after v1 (a derived temperature, a dark-energy fraction derived from the acceleration scale, a Lorentz-invariant completion, a referee defence declaring the theory complete); none of them is landed, and every label of v1 stands. It also adds, in Section 5, what the weak-lensing relation's shape does and does not test. Version 3 adds the natural-units form of the scale in Section 1 and, in Section 8, the audit of two further same-track lanes that present that form and the Hubble ratio as derivations.*

This is the status report of the glm53 track of an AI-assisted research programme on the de Sitter-locked acceleration scale $a_0 = \frac{1}{2}c\sqrt{G\rho_\Lambda}$. It consolidates that track's computational lanes (G001–G027) and its five Lean 4 certificates (43 theorems, exit 0, zero sorry, axioms $\subseteq$ {propept, Classical.choice, Quot.sound}), together with an independent audit of the track and two companion lanes, L247 and L248, whose certificates live in a separate file of 189 theorems on the same standard. Where the audit relabelled a rung, the relabelling is stated here and the original claim is not repeated. The track's own later lanes include an audit response that accepts three of four defects raised against it, including the missing matter coupling in its action; that correction is incorporated below.

What the Lean certificates certify. Every theorem in the five files is an algebraic identity or inequality of the static problem: the phantom bracket, the sound-speed forms, the lensing-sum cancellation of the bimetric frame algebra $((\Phi + c/2) + (\Psi - c/2) = \Phi + \Psi)$, the identification itself (the isothermal density at $\sigma^2 = GM/2r_M$ equals $\sqrt{GMa_0/4\pi Gr^2}$), the existence and uniqueness of the external-field cap radius. None of them certifies a dynamical statement: that the cold sector reaches the stated temperature, that any equilibrium is stable, or that any branch is selected. The word “certified” below means “the algebra is machine-checked”, never more.

1. The reading in one paragraph

The radial acceleration relation is read not as a force law but as the hydrostatic equilibrium of a cold dark sector in every baryonic potential well: if that sector sits at the virial temperature of its well,

$$\sigma^2 = \frac{GM}{2r_M}, \quad r_M = \sqrt{\frac{GM}{a_0}}, \quad a_0 = \frac{1}{2}c\sqrt{G\rho_\Lambda},$$

(in natural units the scale reads $a_0 = \Lambda_{\text{DE}}^2/(2M_{\text{Pl}})$, with $\rho_{\text{DE}} \equiv \Lambda_{\text{DE}}^4$, $\Lambda_{\text{DE}} = 2.24 \text{ meV}$, and $M_{\text{Pl}} \equiv G^{-1/2}$, not the reduced Planck mass, which would change the coefficient by $\sqrt{8\pi}$; this is Milgrom's a_0 - Λ coincidence with the coefficient fixed at $\frac{1}{2}$, equivalently $a_0 = \frac{1}{2}c\sqrt{G\rho_\Lambda} = cH_0/6.99$, and it is a notation, not a derivation), then an isothermal sphere at that temperature has exactly the deep-MOND phantom density, $\rho_{\text{ph}} = \sqrt{GMa_0}/(4\pi Gr^2)$. That equality is Milgrom's: the deep-MOND phantom of a point mass is a singular isothermal sphere, and the coefficient is one by the definition of σ^2 . What would make it physics is a law that forces the sector to that temperature. The track's own formation calculation (rung 4) does not establish one, and a companion lane (L247) exhibits a constitutive law from which the identification follows but which still does not fix the amplitude. The reading is therefore a *candidate* whose central rung is open, with a two-component architecture (equilibrated inner phantom inside the external-field cap, free cold dust outside) that is proposed, not confirmed, and with one recorded failure in the Milky Way.

2. The derivation chain, every rung labelled

Rung	Statement	Status
0	$s = c\sqrt{G\rho_\Lambda}$, one acceleration from the measured dark energy	MEASURED (cosmology)
1	$a_0 = s/2$: the 2 is the integer $n = 2$ that 155 SPARC curves select with nothing fitted (rms 0.150 dex, the registered L232 value)	SELECTED BY DATA (G002; Lean <code>deep_mond_law</code>); the integer is not derived (rung 9)
2	The interpolating function is the SPARC-selected member $\mu_2(x) = 1 - (1 + x/2)^{-2}$	MEASURED (empirical shape)
3	$r_M = \sqrt{GM/a_0}$ is the dimensionally unique galactic length	DERIVED + LEAN (<code>mond_length_unique</code>)

Rung	Statement	Status
4	The cold sector equilibrates at $\sigma^2 = GM/2r_M$	POSTULATED. The N-body cited for this rung (K001) integrates Newtonian gravity in units where $r_M := 1$; a_0 never enters its equations of motion. Its own initial-condition runs give a relaxed half-mass radius $r_{50} = 1.06, 1.57, 3.17$ for start radii $R_0 = 2, 3, 6$, i.e. $r_{50} = 0.53 R_0$: the confinement radius is the initial radius, not an attractor at r_M , and the settled dispersion is half the target. No formation calculation in the repository establishes this rung. v2: three lanes written after v1 (G046, G056, Q001) claim to derive it; each inserts the r^{-2} profile it claims to derive, and the first fails eight of its own eleven checks (Section 8).
5	The identification: the equilibrated density is the deep-MOND phantom, coefficient one	IDENTITY (Milgrom; Lean <code>equilibrated_is_phantom</code>). It is a consequence of rung 4, not evidence for it. L247 derives it instead from a constitutive law $p = P(a)$ (Section 5).
6	The relation is tight: with a free stellar mass-to-light ratio per galaxy in $[0.2, 1.2]$ the per-galaxy floor is 0.064 dex median, and the outer half is tighter (0.055)	MEASURED WITH A PER-GALAXY NUISANCE (G013). A nuisance minimised per galaxy lowers the residual by construction; 42 of 155 galaxies remain above 0.10 dex and the mean is 0.11. Not a confirmed prediction.

Rung	Statement	Status
7	The external-field cap confines the equilibrium where the internal field falls to the external one (Milky Way ≈ 6 kpc; solar pairs unbound; clusters core-confined)	PROPOSED, WITH ONE RECORDED FAIL. G003 V5 is a failure in that lane's own reading: the capped phantom gives a Milky Way mass $17 \times$ below the measured $1.3 \times 10^{12} M_{\odot}$ within 100 kpc and a confined fraction 0.161 against the ledger's 0.14. The rescue, free cold dust outside 6 kpc, is a cold dark matter halo carrying 94% of the Milky Way's dark mass. The 6 kpc break is a prediction for Gaia DR4, not a confirmation. Section 5 adds that the cap lies below every radius the weak-lensing data reach.
8	Clusters: the baryon-steepened isothermal gives residual slope -1.478 against the certified -1.53 (G008); the field solve supplies $2.8\text{--}3.1 \times$ baryons at R_{500} , robust to the kernel swap (G016, G017); the bulk of the cluster residual is free dust	PARTIAL. The shape and temperature are the reading's own; the amplitude is the framework's known $\approx 2 \times$ cluster residual and is carried by cold collisionless matter, as in every other completion in the programme.
9	$n = 2$ is a measurement; the count-statistics route (G009), the dimensional route (L239), the EFT route and four structural searches do not derive it	EMPIRICAL
10	Flat $a_0(z)$ ($w = -1$): the deep-MOND Tully–Fisher zero point at $z \approx 2.5$, 0.00 dex against +0.33 for a rising scale, ± 0.13 decides at 20:1 (G011; pre-registered, DOI 10.5281/zenodo.22563139)	REGISTERED TEST
11	The solar neighbourhood: near-Newton wide binaries ($\gamma_v \approx 1.00\text{--}1.05$) with a period–separation signature (G006, G014, G018)	REGISTERED TEST (Gaia DR4)

Five checks in the track's lanes pass on a literal True and are not evidence (G001 V8, and one check each in G014, G015, G017, G020); the pass counts quoted in the lane outputs include them. The G003 lane prints 7/7 PASS while its own reading records V5 as a failure; the failure is the finding.

3. What is dead: the force-law pincer

Every relativistic *force-law* completion of the parameter-free curve is closed under an existing constraint, and these closures stand independently of the reading above:

- Modified gravity (AQUAL/QUMOND): the Cassini external-field quadrupole is $6.44 \times / 7.63 \times$ the Park 2026 ceiling on the two footings (L243, exact AQUAL; $5.45 \times / 6.29 \times$ on the anchored DHF instrument, G004/G005). The derived length $\xi = r_M$ does not rescue it (G005).
- Modified inertia: lensing-dead by conformal cancellation (L241).
- Disformal/vector (TeVes/AeST class): preferred-frame $\alpha_1 = O(1)$, kernel-independent (L244; DC-013/DC-019).
- Bimetric/composite: lensing-dead by exact frame algebra (G007, 11 Lean theorems): the conformal lever cancels in the lensing sum, the disformal lever is inert in both channels, the scalar's stress channel is 2×10^{-6} short.
- The photocount mechanism: the variance floor it predicts (≈ 0.3 dex) is three times the observed scatter (G009).
- The strictly bound solar cloud: excluded by DR3 wide binaries (G006).
- Pure power-law cluster fluids: acausal at the edge (G008).
- A clock-derived κ : the Clockmaker's Dilemma (G001, 9 Lean theorems).

4. The equilibrium architecture, and its recorded tensions

The reading's architecture is one field equation in three regimes: galaxy outskirts (the deep branch, the relation itself), clusters (the transition branch, supplying $2.8\text{--}3.1 \times$ baryons at R_{500} with the observed slope, the rest carried by free dust), and the solar system (inert, corrections of order 10^{-10}). Two tensions are recorded with their instruments, exactly like the successes:

- Growth (G020): the reading inherits, through its field equation, a raise of σ_8 by 1–3% over Planck- Λ CDM ($S_8 = 0.843\text{--}0.847$ against 0.834) and of $f\sigma_8$ by 1–4% at $z = 0.3\text{--}1.0$. That is opposite in sign to the direct-lensing S_8 values; the gate is open and killable by KiDS, DES and DESI.
- Cluster amplitude (G012, G016): uncapped, the pure isothermal phantom over-supplies by 1.9–2.2 $\times$; capped, it under-supplies by the known residual. The cap is doing the work, and what it leaves is cold collisionless matter.

5. The companion result, and the test that decided it

Lane L247 exhibits a covariant, non-barotropic constitutive law for the cold sector, $p = P(a)$ with $a^\mu = u^\nu \nabla_\nu u^\mu$ the medium's own proper acceleration, matched so that its hydrostatic state in a spherical baryon field is any kernel's phantom exactly (Lean, with the kernel left undetermined). Its deep limit is $p = g^2/8\pi G$, the gravitational field's energy density; hydrostatic equilibrium together with Poisson then forces the flat curve $g = C/r$ with C free, and $p/\rho = C/2$, so the isothermal identification of rung 5 becomes a consequence rather than a postulate. On the homogeneous background the pressure is exactly zero and second order at linear order ($w_{\text{eff}} \approx 10^{-9}$ at recombination), so the cosmology is

Λ CDM's with nothing tuned. It does not fix the amplitude: a supply-limited inner edge gives Tully–Fisher exponent 3, excluded at $\geq 5\sigma$ on 140 SPARC curves, so the a_0 kink must be imposed and κ remains the measured boundary condition of the zero-mode theorem.

That lane also named the test that would decide it. Lane L248 has now run that test, against the lane's own candidate.

The closed form. If the phantom producing a galaxy's weak-lensing signal is real mass drawn from a finite budget $B = M_{\text{dark}}/M_{\text{bar}}$, the enclosed mass freezes once the budget is spent, at $r_t = (1 + B) r_M$. The baryonic acceleration there is

$$g_{\text{bar},t} = \frac{a_0}{(1 + B)^2},$$

independent of the galaxy's mass, so every galaxy in a stacked measurement must turn at the same acceleration. Beyond the turn the frozen mass gives $g_{\text{obs}} = (1 + B) g_{\text{bar}}$: the relation passes from log-slope $\frac{1}{2}$ to log-slope 1, continuously at $a_0/(1 + B)$, and strictly downward. All four statements are machine-checked.

The measurement. Against the published KiDS-1000 excess-surface-density profiles of Brouwer et al. (2021), converted by that paper's own relation — which is exact for the r^{-2} profile under test — and using the full covariance, the log–log slope over the eight bins below $10^{-13} \text{ m s}^{-2}$ is

$$0.537 \pm 0.026,$$

which is 17.6σ from the truncated value 1 and 1.4σ from $\frac{1}{2}$. The relation has not turned. Against the truncated model the comparison gives $\Delta\chi^2 = +349$ at the abundance-matching budget (Moster et al. 2013, $B = 32$), $+970$ at the cosmic dark-to-baryon ratio, and $+75$ even at a stress budget $B = 100$; both a_0 footings agree. Radius-resolved, the enclosed lensing mass inside 1 Mpc already exceeds the galaxies' entire abundance-matching budget by factors 1.5, 4.1 and 8.7 in the three narrow stellar-mass bins.

What it kills, our own candidate included. That excess is real mass in the universe — in the standard picture the two-halo term of correlated neighbours — but it is not bound to the lens, so no reading whose lensing phantom is the galaxy's own equilibrated sector may claim it. This removes L247's medium as the source of the lensing signal, and with it every reading in which the phantom carrying the lensing is the galaxy's own bounded sector. MOND as a force law is untouched: no mass is there, so no budget applies. The capped equilibrium reading of rung 7 is also untouched, and the same computation says why — the cap for a lens of the sample's median mass sits at 5.8 kpc, below the innermost lensing bin at 35 kpc, so it covers none of the measured range. That reading evades the test by making no prediction where the data are, at the stated price that ordinary cold collisionless matter carries all of the signal these data measure.

What the shape of the lensing relation tests (v2, L257). A prior question the programme had not asked: does the deep branch's agreement with the square-root law carry gravitational information at all? If the enclosed mass grows linearly, $M(r) = M_0 r$, then $g_{\text{obs}} = GM_0/r$ and $g_{\text{bar}} = GM_b/r^2$, so $g_{\text{obs}}^2/g_{\text{bar}} = GM_0^2/M_b$ is constant in r : the square-root branch holds identically for any theory delivering an isothermal mass run, with $a_{\text{eff}} = GM_0^2/M_b$ (machine-checked). In the measured variable, an excess surface density falling as $R^{1-\gamma}$ maps to a relation of log-slope $(\gamma - 1)/2$ with no dynamics anywhere. The four KiDS-1000 profiles' own power-law slopes predict a relation slope of 0.547 ± 0.024

from the profiles alone, against the 0.537 ± 0.026 measured on the relation: 0.3σ . What is left is the amplitude's scaling with baryonic mass, which a universal a_0 forces to be exactly $\frac{1}{2}$ (machine-checked); measured on the three narrow stellar-mass bins it is 0.382 ± 0.079 , 1.5σ from $\frac{1}{2}$ and 2.3σ from the two-halo expectation of 0.2, so the data do not presently choose. The implied a_{eff} is $2.4\times$ (canonical) and $2.0\times$ (alternative) the registered footing, the same unbound mass the truncation test measured. The lensing relation constrains this framework through the mass budget and the mass-scaling exponent, and not through the visual agreement of the curve's shape.

Two limits on this result. The parameter-free square-root branch itself fits these data poorly in absolute terms ($\chi^2/\text{dof} = 13.9$ canonical, 9.5 alternative), so every verdict here is a *relative* comparison of two models on identical data and covariance, not a claim that either describes the data well. And dropping the seven lowest-acceleration bins leaves only one bin below the turn, whereupon the discrimination collapses to $\Delta\chi^2 = +5$: the verdict rests on the low-acceleration bins, because that is the only place a mass-budget truncation can appear.

6. What can be claimed, and what cannot

Claimable: the parameter-free curve and the data's selection of $n = 2$ (with L230–L233); the complete force-law pincer, with the bimetric door closed by a Lean certificate; the cluster shape and temperature from the isothermal phantom; the external-field cap as a predictive architecture with a stated Milky Way break radius; the period–separation signature as a novel observable; from L247, a constitutive law from which the identification follows; and, from L248, the mass-independent truncation acceleration $a_0/(1+B)^2$ together with its measurement, which excludes the whole real-mass-phantom class as an explanation of weak lensing.

Not claimable: a relativistic force-law theory of gravity (proven unavailable on this evidence); a derivation of the equilibrium temperature (rung 4 is postulated); a derivation of $n = 2$ or of κ ; cluster dark-matter-free status (free dust carries the cluster bulk and the CMB third peak, the Lyman- α forest and the merging-cluster gate force the same cold collisionless object); the dissolution of the galactic double-counting liability (it would follow from rung 4, which is not established); wide-binary velocity boosts (excluded by DR3); that the equilibrated sector produces the weak-lensing signal (excluded in Section 5); and anything favouring this framework over ΛCDM . *Added in v2 (Section 8)*: a dark-energy fraction derived from a_0 ; a derived equilibrium temperature; a Lorentz-invariant completion that clears Cassini; a parameter count of zero; and any reading in which the equilibrated sector both sources the field equation and is the phantom.

7. The tests that decide it

Test	Prediction	Decides	When
Deep-MOND Tully–Fisher zero point, $z \approx 2.5$	0.00 dex (flat a_0) vs +0.33 (rising)	the a_0 – Λ tie, at 20:1	JWST/ALMA, registered
Gaia DR4 wide binaries	$\gamma_v \approx 1.00$ – 1.05 , binned profile stated (G018), period–separation signature	the unbound-cloud architecture	Dec 2026, registered

Test	Prediction	Decides	When
Gaia DR4 dark-density mapping	break at ≈ 6 kpc; inner profile the zero-parameter phantom	the external-field cap	Dec 2026+
S_8 and $f\sigma_8$	$S_8 = 0.843\text{--}0.847$, $f\sigma_8$ raised 1–4%	the growth gate	KiDS/DES/DESI, now
Lensing amplitude vs baryonic mass	exponent $\frac{1}{2}$ (universal a_0) vs 1 (one-halo) vs 0.2 (two-halo)	whether the deep branch carries a universal scale	RUN (L257): 0.38 ± 0.08 ; undecided, needs wider mass coverage
Stacked lensing profile, mass budget	turn at $g_{\text{bar}} = a_0/(1+B)^2$, slope $\frac{1}{2} \rightarrow 1$	every reading whose lensing phantom is the galaxy’s own bounded sector	RUN (L248): no turn, slope 0.537 ± 0.026 , class excluded

8. Audit of the closure claims made after v1 (2026-09-14 and 15)

In the day after v1 the same track, joined by four sibling agent tracks, promoted the reading to “a complete theory”: a board labelling every rung of its chain derived, a referee defence declaring the solar system, lensing and cosmology answered, and two new rungs, a derived temperature and a dark-energy fraction derived from a_0 . Lane L258 tests those promotions on the tracks’ own committed numbers and on this programme’s own lanes (2 of 11 checks pass; the failures are the findings). Both a_0 footings, 9.3619×10^{-11} and $1.1279 \times 10^{-10} \text{ m s}^{-2}$, are carried throughout.

The dark-energy fraction “from a_0 alone” is circular. The claim (G052, G058, Q002) is $\Omega_\Lambda = 32\pi a_0^2/(3H_0^2 c^2) = 0.6857$, “+0.07% of Planck, the MOND scale and the dark energy are one measurement”. The registered $a_0 = 9.3619 \times 10^{-11}$ is $\frac{1}{2}c\sqrt{G\rho_\Lambda}$ evaluated at $\Omega_\Lambda = 0.685$, $H_0 = 67.4$ (reproduced to +0.004%), and the identity applied to it returns its input exactly (symbolic residual zero). The advertised agreement is the ratio $(67.4/67.36)^2 \times (0.685/0.6847) = 1.0016$ between two roundings of the same Planck parameters; the Q002 lane’s own closing paragraph concedes that “ a_0 ’s canonical value was derived from ρ_Λ in the first place”. Fed an a_0 that was not built from Ω_Λ , the identity gives $\Omega_\Lambda = 1.21$ from this programme’s own free-scale fit at the selected integer (1.2457×10^{-10}) and 1.13 from the literature relation (1.20×10^{-10}), both above unity and 60 to 70σ from Planck; the relation’s $\pm 20\%$ systematic maps to $\Omega_\Lambda \in [0.72, 1.62]$. The “flatness residual” $\Omega_{\text{dm}} = 1 - \Omega_\Lambda - \Omega_b = 0.2649$ is Planck’s own Ω_{dm} fed back, and is negative over most of that band. The alternative footing’s “exclusion at 44σ ” is the statement that the critical density is not the dark-energy density: it is $\frac{1}{2}c\sqrt{G\rho_{\text{crit}}}$ to 0.23% and returns $\Omega = 1$ by construction. What survives is the long-known coincidence $a_0 \sim \sqrt{G\rho_\Lambda}$ at the 20 to 30 per cent level (Milgrom 1983); nothing sharper is delivered, and a “parameter count of zero” that omits the cold-dust abundance and the cluster and Milky-Way dust normalisations is not a parameter count.

The same identity in natural units (v3). Two later lanes of a sibling track present $a_0 = \Lambda_{\text{DE}}^2/(2M_{\text{Pl}})$ and $cH_0/a_0 = \sqrt{32\pi/3\Omega_\Lambda} = 6.994$ as derivations, the first with the $\frac{1}{2}$ identified as the graviton’s two transverse polarisations. Lane L260 checks each on its own terms: the first identity holds for every coefficient once $\rho_\Lambda \equiv \Lambda_{\text{DE}}^4$ and $M_{\text{Pl}} \equiv G^{-1/2}$ are substituted (the reported ratio of exactly one reproduces for $\kappa = \frac{1}{3}$, $1/2\pi$ and 0.461), its “predicted” Λ_{DE} is computed from an a_0 that was built from ρ_Λ

three lines earlier (a galactic a_0 misses by 13 to 15 per cent), the vacuum value $f(0) = -1$ is the zero mode of Section 8's first paragraph, and the deep slope of the family μ_n is n by construction while the static scalar equation contains no tensor polarisation, so the two integers are unrelated. The second is the programme's own definition $a_0 = cH_\Lambda/Z$ with $Z = \sqrt{32\pi/3}$, printed as 6.99 by an earlier lane and rounded to 7 elsewhere. Neither changes a label.

The temperature is not derived; the profile is assumed. Three lanes (G046, G056, Q001) promote rung 4 to “derived, three independent routes”, with $\sigma^2 = \sqrt{GM_b a_0}/2$ as the hydrostatic equilibrium of the cold sector in the scalar's deep force $g = C/r$, $C = \sqrt{GM_b a_0}$. Solving that balance in general, $d(\rho\sigma^2)/dr = -\rho C/r$ gives $\rho = A r^{-C/\sigma^2}$: the temperature is C/γ for every profile slope γ , with the amplitude A free. Hence $\sigma^2 = C/2$ is equivalent to $\gamma = 2$, which is the statement that the dust is the phantom, the identity of rung 5. G046 (parts 3 and 4) and G056 (the “log-potential virial $2\sigma^2 = gr$ ”) insert $\gamma = 2$ and read the temperature back out; Q001's identity $P/\rho = C/2$ uses the constitutive law of Section 5, which was constructed to yield the phantom and which that lane itself records as leaving the amplitude free. The lane cited first for the promotion, G046, passes 3 of its own 11 checks: the three it names as the derivation and its full-sphere integration fail in its own output, its dimensional block is off by 10^3 in r_M and 10^6 in density (a parsec-for-kiloparsec slip), and it prints “rung 4 promoted” regardless. The v1 label, postulated, stands.

The double count, in closed form. The equilibrium reading states that the same field equation which gives the baryons their force also mediates the dust's self-gravity, and that the dust is the phantom. The phantom $\rho = C/(4\pi G r^2)$ has $M(r) = Cr/G$, whose Newtonian pull alone is C/r , the deep force. Solving the μ_2 equation for a Milky-Way-mass well with that dust included as a source, the outer curve rises as $(r/r_M)^{1/4}$: $v = 1.56, 1.72$ and $1.94 v_{\text{flat}}$ at $2, 5$ and $10 r_M$ on both footings. Adding the dust's Newtonian pull to the baryons' boosted response instead gives $1.44 v_{\text{flat}}$, a Tully-Fisher mass normalisation off by four. The only flat version is Newtonian gravity with the dust and no scalar force on anything, which is a cold-dark-matter halo with a prescribed profile and a free amplitude. The v1 double-counting liability is therefore not dissolved by the identification; it is quantified by it.

The completion. The frozen-scalar completion (H011, G054, G055) states that its static law is the μ_2 AQUAL equation; it therefore inherits the Cassini external-field quadrupole of Section 3 unchanged, $6.44\times$ and $7.63\times$ the ceiling on the two footings. The referee defence answers the solar system with a different construction, the fixed-congruence “Horn A” (G032), whose own output states its cost as explicit local Lorentz violation and which a later lane (G043) shows is not a gauge choice, while it answers Lorentz invariance with the frozen scalar, which has no congruence and hence no such PPN result: two mutually exclusive theories are cited as one. The weak-lensing signal, the cluster bulk and 94 per cent of the Milky Way's dark mass are carried, in the tracks' own words, by “the dust as ordinary matter” (the Milky Way lane records the capped phantom a factor 17 short and prints PASS), with an abundance that is Planck's fed back or fitted. The honest description of this construction is Λ CDM plus a scalar whose one remaining job, the galactic relation, double-counts with the dust it names. On the certified μ_2 solve the Milky Way itself is not in trouble: a point mass of $7 \times 10^{10} M_\odot$ gives 221 (canonical) and 226 (alternative) km/s at the solar radius against the measured 232.5 ± 5 before the disc correction; a sibling draft's “three times the baryon budget” applies the deep-MOND Tully-Fisher amplitude at a radius where $g_N = 1.4 a_0$, and is unit-broken besides.

The certificates. All 36 Lean files of the agent tracks and this programme, 474 theorems, were recompiled with axiom prints. Every certificate of the equilibrium track and its siblings compiles with zero sorry on the standard axioms, and certifies algebra: the “one-constant closure” theorem is the existence and uniqueness of $\rho := 4a_0^2/(Gc^2)$ given a_0 , a definition; the identification is Milgrom's isothermal

identity; the sound-speed identity is $P/\rho = C/2$ for two r^{-2} laws. One sibling file does not compile (a missing import; 5 of its 22 theorems rest on `sorryAx`). This programme’s own lensing block had two broken proofs at audit time, one resting on `sorryAx`; both are repaired and the file compiles. The statement of `v1` stands: certified means the algebra is machine-checked, never more.

What stands from the same day. The cluster partition lane (G059) tested three candidate rules on the X-COP fields and none closes the amplitude, the kernel share delivering 0.57 and 0.62 of the deficit at 420 kpc on the two footings: an honest negative, correctly recorded after a units bug in its first run was caught. The external-field split in the extended corpus is not established ($p = 0.11$). The deep-regime sag of the relation, -0.13 to -0.17 dex per dex, resolves to an observational channel, the asymmetric-drift correction for gas pressure support, which closes it to -0.03 dex per dex if the gas dispersion is 10 to 16 km/s and leaves -0.10 if it is below 7 km/s: open on an independent HI dispersion measurement. These, the Lean algebra, and Section 5’s two lensing results are what the day added.

9. Reproduction

Lanes `glm53_push/G001-G027_*.py` with `.out` and `_results.json`; Lean certificates in `glm53_push/lean/` (EQUILIBRIUM_THEORY 12, G001 9, G002+G003 4, G007 11) and the companion blocks `medium_*` and the truncation theorems in `fable_independent_2026/lean_2026/Mondlean.lean` (185 theorems, zero sorry); the audit record and lanes L247–L248 in `fable_independent_2026/`; for `v3`, `L260_seesaw_identity_audit.py`; for `v2`, `L257_lensing_rar_degeneracy.py`, the audit `L258_closure_audit.py` with `L258_lean_inventory.py` and its JSON inventory of all 36 certificate files, and the same-day lanes L250 (voids), L254 (kinematic SZ) and L256 (force law versus assembly attractor, undecided by its pre-registered rule), each with `.out` and results files, in the same directory (Mondlean 189 theorems, zero sorry). The lensing data are the public Brouwer et al. (2021) release. Compile with `lake env lean` against the repository’s Mathlib.

One caveat on priority within the later lanes: the vertical-slab “column identity” reported there — that the vertical boost is exactly 2 where the baryon column equals $a_0/8\pi G$ — is a restatement of Milgrom’s critical surface density $a_0/2\pi G$ (1989), which already appears throughout this repository. It should not be presented as new.

This document states what is earned, what is measured, what is dead, and what will decide it. It does not state that a theory is closed.