

The Inverted Black Hole: Why the de Sitter–MOND Acceleration Scale Is Uniquely Cosmic

Carl P. Zimmerman · Briar Creek Tech · 2026-06-26

Abstract

The MOND acceleration scale can be written exactly as a horizon surface gravity divided by a single dimensionless constant:

$$a_0 = c^2 \sqrt{(\Lambda/32\pi)} = cH_\Lambda / Z, Z = 2\sqrt{(8\pi/3)} = \sqrt{(32\pi/3)} \approx 5.789.$$

The de Sitter horizon that supplies cH_Λ is, geometrically, an **inverted black hole** — a horizon we sit *inside* rather than outside. This note takes that picture literally and asks what happens when the same law, $a_0 = (\text{horizon surface gravity})/Z$, is applied to a **real** black-hole horizon instead of the cosmic one. The dual is clean and exact: $a_{0_BH} = c^4/(4GMZ)$, and the radius at which it would switch on, $r_{\text{cross}} = \sqrt{Z} \cdot r_s = 2.406 r_s$, **is universal — the black-hole mass cancels completely**. For Sagittarius A*, M87*, and a $10 M_\odot$ stellar black hole this crossover sits at the same place in units of the Schwarzschild radius, between the photon sphere ($1.5 r_s$) and the ISCO ($3 r_s$).

We show that this dual nonetheless **self-cancels into pure general relativity** for every clean observable, for two structural reasons — mass-independence (any deviation expressed as $f(r/r_s)$ is already inside the GR metric by general covariance and the equivalence principle) and the free-fall / Hartle–Hawking theorem (a geodesic observer sees no horizon bath, and modified inertia responds only to *proper* acceleration). We explicitly reject the naïve “ $g \gg a_0$ near a horizon, so the effect vanishes” objection: it is **false**, because a_{0_BH} scales up with c^4/GM identically, keeping the ratio a/a_{0_BH} of order unity right down to the horizon. The effect does not trivially vanish; it cancels for a precise and instructive reason.

This singles out a **uniqueness theorem**: the cosmic a_0 is the *only* acceleration that is neither sourced by local matter nor removable by any local free-fall — which is exactly why a_0 is cosmic and not, say, set by the nearest black hole. The one falsifiable residue is a **null**: the modified-inertia reading forces *exactly-GR* shadows, ISCO frequencies, ringdown spectra, and inspiral waveforms, whereas a *metric*-MOND completion (AeST, MOG) predicts shifted strong-field observables. A future ngEHT or LISA detection of a MOND-scale black-hole-metric deviation would therefore **falsify metric-MOND while being consistent with this framework**.

Both ways: this is a *structural duality and a null*, not a new force, not a new positive signal, and not a theory of everything. Its only discriminating axis is modified-inertia-versus-modified-gravity — the same axis as the existing Cassini bound, extended to the strong field — and it is decadal. It claims **less**, not more, than the framework’s published one-parameter effective theory (DOI

10.5281/zenodo.20935948, 10.5281/zenodo.20938891) and is fully consistent with the author’s 2026 retraction of the earlier theory-of-everything overclaims.

1. Background: the de Sitter horizon as an inverted black hole

1.1 The acceleration scale as a horizon surface gravity

The empirical MOND scale $a_0 \approx 1.2 \times 10^{-10} \text{ m s}^{-2}$ is the acceleration below which galaxy rotation curves flatten. The framework’s single physical claim is that this scale is *cosmological in origin* and equals a de Sitter horizon surface gravity reduced by one geometric factor:

$$a_0 = \frac{c H_\Lambda}{Z}, \quad H_\Lambda \equiv c \sqrt{\Lambda/3}, \quad Z = 2\sqrt{8\pi/3} \approx 5.789.$$

Here Λ is the cosmological constant and $H_\Lambda = H_0 \sqrt{\Omega_\Lambda}$ is the de Sitter (dark-energy-only) expansion rate, numerically $1.807 \times 10^{-18} \text{ s}^{-1}$ ($H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_\Lambda = 0.685$), giving $a_0 = 9.36 \times 10^{-11} \text{ m s}^{-2}$ — the framework’s published value. The quantity cH_Λ is the **surface gravity of the de Sitter horizon**: just as a black hole of mass M has a horizon at the Schwarzschild radius $r_s = 2GM/c^2$ with surface gravity $\kappa_{\text{BH}} = c^4/4GM$, de Sitter space has a cosmological horizon at radius $r_{\text{dS}} = c/H_\Lambda$ with surface gravity $\kappa_{\text{dS}} = cH_\Lambda$. Both are accelerations; both set a Hawking/Unruh temperature $T = \hbar\kappa/2\pi k_{\text{B}}$. The number $Z = \sqrt{(32\pi/3)}$ is algebraically forced — it is the Einstein coupling 8π (from $\rho_{\text{DE}} = \Lambda c^2/8\pi G$) times the Friedmann 3 (from $H^2 = 8\pi G\rho/3$), with a single free outside factor of 2. (The closure of *that* coefficient is the subject of a companion note; here we treat Z as fixed and ask about its application to real horizons.)

1.2 Why “inverted”

A Schwarzschild black hole is a horizon seen **from outside**: matter is concentrated at the center, the horizon surrounds it, and surface gravity *increases* as the mass increases — or, for fixed mass, as you fall *inward*. The de Sitter horizon is the same geometry turned inside-out. We sit *inside* it; the “concentration” is the vacuum energy filling all space; the horizon is the sphere at comoving distance c/H_Λ beyond which recession exceeds c ; and surface gravity is felt looking *outward*. Formally, the static de Sitter metric

$$ds^2 = -\left(1 - \frac{r^2}{r_{\text{dS}}^2}\right) c^2 dt^2 + \left(1 - \frac{r^2}{r_{\text{dS}}^2}\right)^{-1} dr^2 + r^2 d\Omega^2$$

is the Schwarzschild metric with the sign of the curvature term flipped and the mass term replaced by the Λr^2 term — a black hole inverted in both the location of the horizon (far, not near) and the sign of the potential. This is not loose analogy: de Sitter and Schwarzschild are the two limits of the Schwarzschild–de Sitter family, and $a_0 = \kappa/Z$ is one law evaluated on the two horizons. The framework’s content is that **inertia is the response to the bath of the inverted (cosmic) horizon**, and the natural test of that idea is to evaluate the *same response* on a real horizon.

2. The dual: $a_{0_BH} = c^4/(4GMZ)$ and the universal crossover

Apply $a_0 = (\text{surface gravity})/Z$ to a Schwarzschild horizon, $\kappa_{BH} = c^4/4GM$:

$$a_{0,BH} = \frac{\kappa_{BH}}{Z} = \frac{c^4}{4GMZ}.$$

Numerically (verified in mpmath, dps = 40):

Black hole	Mass	a_{0_BH} (m s ⁻²)	ratio to cosmic a_0
Sagittarius A*	$4.3 \times 10^6 M_\odot$	6.11×10^5	6.3×10^{15}
M87*	$6.5 \times 10^9 M_\odot$	4.04×10^2	4.2×10^{12}
Stellar (10 $M_\odot$)	$10 M_\odot$	2.63×10^{11}	2.7×10^{21}

Every real horizon is 12–21 orders of magnitude “hotter” than the cosmic one — unsurprising, since cosmic a_0 is the gentlest acceleration in nature.

The crossover radius is universal. Modified inertia switches regime where the Newtonian acceleration $g(r) = GM/r^2$ equals the local scale a_{0_BH} :

$$\frac{GM}{r^2} = \frac{c^4}{4GMZ} \Rightarrow r_{\text{cross}}^2 = \frac{4G^2 M^2 Z}{c^4} = Z r_s^2, \quad r_{\text{cross}} = \sqrt{Z} r_s = 2.406 r_s.$$

Why the mass cancels. $g(r) = GM/r^2$ carries one power of M ; $a_{0_BH} = c^4/4GMZ$ carries one *inverse* power of M . Setting them equal makes M^2 appear on both sides and cancel, leaving a pure number times r_s^2 — and r_s itself is the only length the problem contains. So r_{cross} is the same multiple of r_s for a stellar black hole and for M87*: **2.406 r_s , always.** It sits between the photon sphere (1.5 r_s) and the ISCO (3 r_s). The dimensionless acceleration at the ISCO is likewise mass-free and $O(1)$: $x_{\text{ISCO}} = a/a_{0_BH} = Z/9 = 0.643$ on the Newtonian reading (≈ 0.79 including the proper-acceleration/redshift factor). For *every* black hole, the MOND regime would, naïvely, bite right in the observationally active band of photon ring, ISCO, and ringdown.

3. The self-cancellation: why it is nonetheless pure GR

3.1 First, kill the lazy objection

A common reflex is: “near a black hole g is enormous, so $g \gg a_0$, so we are deep in the Newtonian/GR regime where the MOND interpolation $\mu \rightarrow 1$ and nothing changes.” **This is false, and we reject it explicitly.** It silently uses the *cosmic* a_0 as the yardstick. But the relevant scale here is the black hole’s *own* a_{0_BH} , which scales up as c^4/GM in lockstep with g . Their ratio $x = a/a_{0_BH}$ is set entirely by r/r_s and stays $O(1)$ (indeed 0.6–0.8) right down to the horizon, as §2 shows. So the effect does **not** trivially vanish near the horizon. Something subtler is going on.

3.2 First reason it cancels: mass-independence $\Rightarrow$ already inside the GR metric

The tell is precisely that $r_{\text{cross}} = 2.406 r_s$ and $x_{\text{ISCO}} = 0.643$ depend *only* on r/r_s . The scale $a_0_{\text{BH}} = \kappa_{\text{BH}}/Z$ is built from c^4/GM — a pure general-relativistic quantity, the horizon surface gravity, which the Schwarzschild metric already contains. Any “modification” expressible as a function $f(r/r_s)$ carries no information the metric does not already have, so by **general covariance plus the equivalence principle** it is absorbed into the geodesic motion GR already prescribes. A correction that is a function of the metric’s own invariants is not a new force; it is a re-labeling of GR. This is the structural opposite of the cosmic case, where $a_0 \propto \sqrt{\Lambda}$ introduces a scale *external* to the local two-body metric.

3.3 Second reason it cancels: the free-fall / Hartle–Hawking theorem

The framework’s inertia is sourced by **proper** acceleration — the acceleration a body feels relative to its local inertial frame, the thing an accelerometer reads. A freely-falling (geodesic) detector near a black-hole horizon reads **zero**, and by the standard Hartle–Hawking / Unruh–DeWitt result it sees **no Hawking or Unruh bath**: the Hartle–Hawking vacuum is regular on the horizon precisely for the freely-falling observer. Orbits, photon trajectories, inspiraling binaries, and the ringing of the final black hole are all **geodesic** phenomena. A geodesic observer has no proper acceleration for the modified-inertia response to act on, and no bath to respond to. Therefore a_0_{BH} touches **no geodesic observable**: the ISCO frequency, the quasinormal-mode (ringdown) spectrum, the photon ring and EHT shadow, the gravitational-wave inspiral, and QPO frequencies are all **exactly GR**, with no a_0_{BH} correction.

The one non-geodesic loophole is genuinely accelerated matter — accretion-disk and jet plasma held off geodesics by pressure and magnetic stress. But there the dynamics are completely swamped by magnetohydrodynamics and radiation pressure, whose forces dwarf any a_0_{BH} -scale inertial correction by many orders of magnitude. It is not a clean test, and we do not claim one.

4. The uniqueness theorem: why a_0 is cosmic

Sections 2–3 deliver more than a null. They isolate **what is special about the cosmic horizon** by showing what fails for every other horizon. State it sharply. An acceleration scale can imprint a *physical*, non-removable modified-inertia effect only if it is:

1. **not sourced by local matter** — otherwise it is a function of the local metric’s invariants (here $f(r/r_s)$) and is absorbed into the GR geodesic by covariance + EP (§3.2); and
2. **not removable by local free-fall** — otherwise a geodesic observer transforms it away and sees no bath, by the Hartle–Hawking theorem (§3.3).

A real black-hole horizon fails **both**: its surface gravity is a local-metric quantity, and its bath is absent for the geodesic observers who execute every clean strong-field measurement. The cosmic acceleration cH_Λ fails **neither**. It is not sourced by any local mass — it is set by Λ , a global vacuum property, the same everywhere — so it is *not* a function of any two-body r/r_s and cannot be covariantly absorbed. And it cannot be removed by free-fall over a bound system, because **there is no global inertial frame that straightens out de Sitter expansion over a Hubble volume**; a freely-falling galaxy still recedes from the horizon. The de Sitter horizon supplies a

genuine, irreducible acceleration *floor* that a bound orbit cannot transform away, whereas a black-hole horizon supplies none for the orbit around it.

This is why a_0 is cosmic. It is not a coincidence of magnitude that the modified-inertia scale matches $\sqrt{\Lambda}$ rather than the surface gravity of the nearest or largest black hole. The cosmic horizon is the **unique** horizon whose acceleration is neither local-matter-sourced nor free-fall-removable — and uniqueness is exactly the property a fundamental, universal scale must have. The dual to real black holes is the proof: run the same law on every other horizon and it cancels; only the inverted, cosmic horizon survives.

5. The falsifiable consequence: an exactly-GR null

The framework therefore makes a sharp, *forced* prediction at the strong-field scale that none of its previously banked tests reach:

Null prediction. The modified-inertia reading predicts **exactly-GR** black-hole shadows, photon spheres, ISCO frequencies, ringdown quasinormal-mode spectra, and inspiral-merger waveforms for Sagittarius A*, M87*, and stellar-mass black holes, with **no a_0 _BH correction whatsoever**.

The discriminating power is real but narrow, and it is a **falsification asymmetry**, not a positive signal:

- A **metric**-MOND completion — a relativistic theory that modifies the *gravitational field equations* in the MOND regime (AeST / Aether-Scalar-Tensor, MOG / Scalar-Tensor-Vector) — generically modifies the black-hole **metric** itself, and so predicts **shifted** shadows, ISCO frequencies, and ringdown spectra at the MOND scale. Since $r_{\text{cross}} = 2.406 r_s$ lands squarely in the photon-ring / ISCO band, such a shift is, in principle, in the observable window.
- This framework, being a modified-*inertia* theory with a **standard GR metric**, predicts no such shift.

Hence a future detection of a MOND-scale black-hole-metric deviation would **falsify metric-MOND while being consistent with — indeed predicted by — this framework**, and a confirmed pure-GR shadow/ringdown is a (weak) point for the inertial reading. The relevant instruments are **ngEHT** (next-generation Event Horizon Telescope: M87* and Sgr A* shadows to ~1–2%, 2030s) and **LISA / LIGO-Virgo-KAGRA** ringdown spectroscopy plus LISA EMRI and massive-black-hole inspirals (2035+).

Both ways — the honest limits. (i) It is a **null**, not a detection: no positive signal can be claimed, only a falsification asymmetry. (ii) Its only axis is **modified-inertia-versus-modified-gravity**, the *same axis* as the framework's existing Cassini bound — it **extends Cassini to the strong field** rather than opening an independent axis. (iii) It is **decadal-plus** and at a precision (~1–2%) that may sit below the level at which a MOND-scale metric shift would even appear. So it does not supersede the framework's two sharpest near-term fronts (a declining $a_0(z)$ tested via the

baryonic Tully–Fisher sign, and a Lorentz-violating $s^{\wedge}\text{TX}$ SME dipole); it is weaker and farther than either. It is a real new consequence, reported at its true strength.

6. Scope: a structural duality and a null — claiming less, not more

To be explicit about what this is and is not:

- It is a **structural duality** ($a_0\text{BH} = c^4/4\text{GMZ}$ with the universal $r_{\text{cross}} = 2.406 r_s$) plus a **forced null** (exactly-GR strong-field observables). Both follow rigorously from the published law $a_0 = \kappa/Z$ and from textbook results (Hartle–Hawking regularity, covariance + EP). No new postulate is introduced.
- It is **not a new force**, **not a new positive signal**, and **not a theory of everything**. It introduces **no new free parameter**: $a_0\text{BH}$ is fixed by the *same* Z and the black hole’s GR mass.
- It is **fully consistent with**, and supplements, the framework’s published one-parameter effective theory (DOI 10.5281/zenodo.20935948, 10.5281/zenodo.20938891) and with the author’s 2026 public retraction of the earlier Standard-Model / theory-of-everything overclaims. If anything it **strengthens** the modest reading: it explains, via the uniqueness theorem, *why* the single physical scale is cosmic, and it adds a strong-field falsification handle on rival metric-MOND theories without adding anything the framework must itself defend.

The net contribution is one new, narrow, far, **null** consequence — the BH-scale exactly-GR prediction — and a clean reason it is a null. Taken together with the duality and the uniqueness theorem, the result tightens the conceptual case for a cosmic a_0 while making no claim the data cannot, in time, check.

Reproducibility note

All numerical statements (a_0 on the framework footing; $a_0\text{BH}$ for Sgr A*, M87*, and a $10 M_{\odot}$ black hole; $r_{\text{cross}} = \sqrt{Z} r_s = 2.406 r_s$; $x_{\text{ISCO}} = Z/9 = 0.643$; $Z = \sqrt{(32\pi/3)} = 5.78881$) were verified in mpmath at 40-digit precision. The crossover-radius derivation (mass cancellation) and the ISCO ratio are one-line algebraic identities reproducible from the public repository. The free-fall / no-bath statement is the standard Hartle–Hawking / Unruh–DeWitt result for a geodesic detector; the covariance + equivalence-principle absorption argument is qualitative and stated as such. The underlying consequence map (real_research/Frontier_Consequences_Inverted_BH_2026-06.md) records the full both-ways analysis, including the parallel finding that taking the de Sitter–Unruh bath as physically real yields a deep-vacuum bath ($T_{\text{dS}} = 2.28 \times 10^{-30}$ K) with no laboratory decoherence or quantum-modified-inertia signature — a defensive null reported there at its true (below-floor) strength.

Carl P. Zimmerman · Briar Creek Tech · 2026-06-26. This work claims a structural duality and a falsifiable null, consistent with and supplementing the author’s published one-parameter effective theory; it does not claim a new force or a theory of everything.