The Field Equations of the Screened Khronon Action, and a Verdict for Every Physics Class

Working note (PAPER70), v2.0, 2026-11-09 (v1.0 deposited as DOI 10.5281/zenodo.24040313; v1.1, which added the kernel erratum of Section 6, deposited as DOI 10.5281/zenodo.24044606; v1.2 deposited as DOI 10.5281/zenodo.24047420; v1.3, which carried the owner's acceptance of the corrected-kernel labels, deposited as DOI 10.5281/zenodo.24047508; v1.4, which added the one-page final theory to the supplement with the paper unchanged, deposited as DOI 10.5281/zenodo.24047634; v1.5, which added the anatomy of the CMB requirement (Section 7) with CMB3 running, deposited as DOI 10.5281/zenodo.24053083; v1.6, which added CMB3's assumption audit, the quarantined hybrid test lane HY1 and X35's answer to CMB2's open corner, deposited as DOI 10.5281/zenodo.24053740; v1.7, which added the running K-mass lanes KR1 and KR2, deposited as DOI 10.5281/zenodo.24056963; v1.8, which added the quarantined cold-gas conversion test lane BG1, deposited as DOI 10.5281/zenodo.24058756; v1.9, which added fix-forward notes on what the residue rests on and on record-internal inconsistencies, the quarantined test lanes PJ1, WD1 and XF1, and HL3's lensing check of the cluster residual, deposited as DOI 10.5281/zenodo.24082193; this version adds the closing statement, the quarantined spin-key test lane SK1 and the cluster-tracer grading lane TR1, with no earlier label changed). AI-assisted research programme; not peer reviewed. Companions: PAPER67 (DOI 10.5281/zenodo.23968171, the law and its failure record), PAPER68 (DOI 10.5281/zenodo.23990944, the action) and PAPER69 (DOI 10.5281/zenodo.24025170, the closure record). $\kappa = \frac{1}{2}$ is fitted. The theory is not closed. Nothing here says the data favour this framework.

Abstract

Gravity carries one extra scale, $a_0 = \kappa c\sqrt{G\rho_{\rm DE}}$, $\kappa = \frac{1}{2}$ fitted (proved not derivable), realised by the v3s screened khronon action. It passes Cassini, PPN, GW speed, galaxy lensing and SPARC, and fails AGC 114905, Malin 1 and the canonical FG1 tilt. It also under-fills X-COP clusters, groups, bright M31 dSphs and Coma UDGs by amounts that vary about 100× between classes, after the IMF accounts for the ellipticals and most of KiDS. Separately, it fails the CMB-lensing shape, which needs cold extra gravity at z 3400. No cold fluid, fixed-temperature sea, thermal light fermion or single measured key tried so far supplies either. The next lane must find one covariant quantity that sets that amplitude (not just a switch) in today's hot systems and is non-zero in the early universe, or show the two failures need separate causes. *(This is the checked seed paragraph of 2026-11-06, verbatim.)*

Since the seed, a joint systematics budget per class also explains the bright M31 dSphs, and a symmetric admissibility cut leaves the Coma UDGs NOT A TEST, so the measured boundary of the law is the X-COP clusters, the groups (fragile) and the CMB-lensing shape, and the seed's "about 100×" spread no longer stands: it rested on the Coma UDG rung, and between X-COP ($f \sim 4.2$ / 3.8) and the groups (1.2–3.5 / 0.9–2.9) the spread is a factor of about 4 (3.5 / 4.2); and a shock-entropy sector written into the action fails by the no-tuning rule, so every physics class on the record carries a label. This note writes down the action in the Legendre form every lane solved (correcting the form printed in PAPER68), its complete covariant field equations checked against direct variation, their 3+1 form and a convergent static solver. It then lists every gate with its frozen label, states the residue as measured numbers, and collects the three frozen tests that can kill the law. Four physical classes of extra ingredient were built and graded on both footings: a cold fluid, a warm fixed-temperature sea, a hot thermal fermion, and a shock-entropy sector written into the action. Each meets its own wall, and no candidate passes every gate. The most probable reading of the record is the deposited, scoped law plus separate per-class systematics, with the cluster, group and ultra-diffuse rows and the CMB wall left unexplained.

1 Conventions

Units $c = 1$ and $K = 4\pi G$ unless stated; restoring c: $K \to 4\pi G/c^2$, the prefactor of R is $c^4/16\pi G$, and $a_0 \to a_0/c^2$ inside W. Two footings are carried and never pooled: canonical $a_0 = 9.3603 \times 10^{-11}$ and alt $1.1312 \times 10^{-10}\,\mathrm{m\,s^{-2}}$. κ is fitted; a Lean proof shows that it is not derivable from the programme's premises. A gate is closed when it carries a label fixed by criteria committed before the first graded number; every lane runs a mutation twin that must fail.

2 The action, in Legendre form

PAPER68 prints $S = \frac{1}{4K}\int\sqrt{-g}\,[R - 2\mathcal{J}(Y) + 2\mathcal{K}(Q)] - \frac{\gamma_L}{2}\int\sqrt{-g}\,[\max(D\cdot P, 0)]^2 + S_m$. Read literally, P then appears only in the stiffness, its own equation forces $D\cdot P$ to be constant, and P decouples from gravity. The

form every lane has actually solved puts $\mathcal{J}$ in Legendre form (QX17):

$$\frac{\mathcal{L}}{\sqrt{-g}} = \frac{R - 2\Lambda + 2\mu_K^2(Q-1)^2}{4K} + P^m A_m - W(K|P|) - \gamma_L F(\Pi) + \lambda\, n^m P_m, \qquad W(p) = \frac{a_0^2}{K}\Big[-\frac{p}{a_0} - \ln\Big(1 - \frac{p}{a_0}\Big)\Big], \tag{1}$$

with $F(\Pi) = \frac{1}{2}[\max(\Pi, 0)]^2$, $\Pi = D_m P^m$ and $\gamma_L = KL_\star^2/2$, plus minimally coupled matter S_m. Maximising over P returns the printed $\mathcal{J}$ exactly: $\max_P[P\cdot A - W(K|P|)] = (a_0^2/K)[u - \ln(1+u)]$ with $u = |A|/a_0$, attained at $K|P| = a_0 u/(1+u)$ (sympy, residual 0). So at $\gamma_L = 0$, Eq. (1) is the printed action; with $\gamma_L > 0$, P is an independent field and Eq. (1) is the theory. **This is an erratum on PAPER68's printed form only; no number changes.**

Adopted stiffness (owner decision, 2026-11-07). The owner adopted the Π^6 blend as the action's stiffness. In Eq. (1) the raw kink $F(\Pi) = \frac{1}{2}[\max(\Pi, 0)]^2$ is replaced by AC21 iter_6's blend, $F(\Pi) = b^2 f(\Pi/b)$ with $b = \rho_{\rm DE}$, the a_0 law's own scale, so there is no new constant. Near 0, f is a pure power, $f(q) = (122363904/44610575)\, q^6$ for $q \le \frac{1}{2}$ ($m = 6 > d + 2 = 5$). On the layer it is a C^4 convex blend, and $f(q) = q^2/2$ exactly for $q \ge 1$, so $F = \frac{1}{2}\Pi^2$ for $\Pi \ge b$. Eq. (1) with this F is the operative action of this note; it supersedes PAPER68's printed raw form. AC2q is not adopted; if it is adopted later, it carries the same blend with $b = X$. The scope of v3s and every graded label are unchanged: the blend moves no failing gate (largest effect $\le 1.4 \times 10^{-4}$ dex). The field equations of the next section hold for any stiffness F with Π_+ read as $F'(\Pi)$ throughout and the stress-tensor term written $\Pi F'(\Pi)\, h_{mn}$; for the raw F this is $2F h_{mn}$, as QX17 certified. QX17D certified the general form against a direct variation, for a generic polynomial F and for the blend's pure-power branch (residuals $\le 7.1 \times 10^{-40}$); the shortcut that keeps $2F h_{mn}$ fails the metric equation by up to 14 %. Limits adopted with the blend: well-posedness is local in time; it is tested in spherical 3D only, and non-radial ($l \ge 1$) modes are untested; folds are unchanged; power 6 is not forced (the smallest integer with $m > d + 2$, admissible, not unique). The AC16 row remains the label of the raw switch.

Definitions. τ is the khronon; $Q = \sqrt{-g^{mn}\partial_m\tau\,\partial_n\tau}$ ($= 1/N$ in unitary gauge); $n_m = -\partial_m\tau/Q$; $h_{mn} = g_{mn} + n_m n_n$; $l = \ln Q$; $A_m = n^n\nabla_n n_m = -h_m{}^n\partial_n l$; $D_m = h_m{}^n\partial_n$; $\Pi = \nabla_m P^m - P^m A_m$; $\Pi_+ = \max(\Pi, 0)$; $\hat{P} = P/|P|$; $\mathcal{D}_m = \partial_m\Pi_+ - \Pi_+\partial_m l$.

3 The field equations (QX17, CERTIFIED)

$$\text{(E1)}\quad n^m P_m = 0, \tag{2}$$

$$\text{(E2)}\quad (1 + \gamma_L\Pi_+)A_m + \gamma_L D_m\Pi_+ = KW'(K|P|)\,\hat{P}_m, \qquad \lambda = (1 + \gamma_L\Pi_+)\, n\cdot\partial \ln N + \gamma_L\, n\cdot\partial\Pi_+, \tag{3}$$

$$\text{(E3)}\quad \nabla_m\big[n^m\rho_Q - \lambda P^m/Q\big] = 0, \qquad \rho_Q = \frac{\nabla_m[(1 + \gamma_L\Pi_+)P^m]}{Q} + \frac{\mu_K^2(Q-1)}{K}, \tag{4}$$

$$\text{(E4)}\quad G_{mn} + \Lambda g_{mn} = 2K\big(T^{\rm mat}_{mn} + T^Q_{mn} + T^W_{mn} + T^P_{mn}\big), \tag{5}$$

with

$$T^Q_{mn} = g_{mn}\frac{\mu_K^2(Q-1)^2}{2K} + \frac{\mu_K^2}{K}Q(Q-1)\, n_m n_n, \qquad T^W_{mn} = -g_{mn}W + KW'\,\frac{P_m P_n}{|P|},$$

$$T^P_{mn} = g_{mn}(P\cdot A) + 2P_{(m}\partial_{n)}l + (\nabla\cdot P)\, n_m n_n - 2\lambda\, n_{(m}P_{n)} + \gamma_L\big[-g_{mn}F + \Pi F'(\Pi)\, h_{mn} + h_{mn}P^r\mathcal{D}_r - 2P_{(m}\mathcal{D}_{n)}\big].$$

(E5) Matter is minimally coupled, $\nabla^m T^{\rm mat}_{mn} = 0$. The energy density seen by the foliation is $n^m n^n T_{mn} = (\mu_K^2/2K)(Q^2 - 1) + W + \Pi + \gamma_L F$. At $\gamma_L = 0$ the leaf part of (E2) inverts to $K|P| = a_0|E|/(a_0 + |E|)$, the Legendre map; Π_+ is C^1, so the equations hold in the weak sense across the switch surface.

Checks. Each equation is compared with a direct Euler–Lagrange variation of Eq. (1) on jet symbols (mpmath, 40 digits, three random configurations in 2+1 dimensions): P equation max rel 3.8×10^{-41}, khronon equation 1.8×10^{-41}, full stress tensor 1.2×10^{-40}; FRW reduction exact. The MUTATE twin (two flipped signs) fails at 0.17–0.78.

AC2q (candidate, not adopted; QX17A, CERTIFIED). The dark-energy scalar is the clock, there is no $\mathcal{K}(Q)$ term, and the switch acts on the khronon charge $\Sigma = \Pi + Q^2$: $\mathcal{L}/\sqrt{-g} = R/4K + X - V(\phi) + P\cdot A - W - \gamma\{F(\Pi + 2X) - (2X)^2/2\} + \lambda n\cdot P$, $X = Q^2/2$. Its P equation is (E2) with $\Pi_+ \to \Sigma_+$; its clock equation is $\nabla_m[n^m\rho_\phi - \lambda P^m/Q] = -V'(\phi)$ with $\rho_\phi = J_P/Q + Q[1 - 2\gamma(\Sigma_+ - Q^2)]$, which is $\Box\phi = V'$ at $P = \gamma = 0$; at $P = 0$ the stiffness stress vanishes identically, so the background is the canonical scalar's. The MUTATE twin fails at $0.23/0.39/7.3 \times 10^{-3}$.

4 3+1 form, reductions and solutions

ADM (QX17B). P and N carry no time derivatives, so the P equation and the lapse equation are constraints on each leaf. The P sector's momentum density vanishes on shell, so the momentum constraint is GR's; the tensor principal part is GR's, so $c_T = c$. The Lean kernel file for these steps has 12 theorems, standard axioms only, and its MUTATE twin fails. The lane as a whole is not certified: its numerical ADM row failed as frozen on a sign convention in the checker's off-shell momentum density, and its `solve_bvp` spherical solver did not converge.

Reductions. FRW with $P = 0$: $\bar{Q} = 1 + I_0/(2\mu_K^2 a^3)$ and a dust-like $\rho \propto a^{-3}(1 + w_0/a^3)$. Static, weak, $\gamma_L = 0$: $\nabla \cdot [\mu(|\nabla\Phi|/a_0)\nabla\Phi] + \mu_K^2\Phi = 4\pi G\rho_b$, $\phi = \psi$. Spherical, $\mu_K = 0$: the law $g_N = g^2/(g + a_0)$ (Lean). Static, $\gamma_L > 0$: the Cassini hedgehog inside $L_\star$. AC2q FRW: the lapse–polarization mixing is $(1 - z)$ with $z = \gamma\dot{\phi}^2/\bar{N}^2$, so eliminating P multiplies the induced lapse term by $(1 - z)^2$, not the legacy $(1 - 2z)^2$ (which is the exact factor of a different action). The UV speed is $c^2 = z/(1 - z + z^2)$, and $0 < c_g \le c_{\rm ph} \le 1$ holds for any $f \ge 0$, so FRW health stands (FRW reconciliation chain: sympy ADM reduction and Lean, 11 theorems, standard axioms only; MUTATE 3/3 fail). The P block couples to the lapse only. The static-limit offset is half the earlier value, bounded by 1.1×10^{-10}. Still open: the AC7 Q2 / AC10–AC15 post-Newtonian chain carries the legacy factor and has not been rederived.

Solutions (QX17C, CERTIFIED as a solver). A conservative finite-difference Newton solver with a decaying Bessel outer boundary converges in 1–6 steps on both footings, about 5 s in all. It reproduces the law at $L_\star \to 0$ (worst 3.9×10^{-7}), the solar hedgehog to 7.1×10^{-6} of $g_\star = GM/L_\star^2$, and a $10^{10}M_\odot$ galaxy on the law to 4.8×10^{-6}; full-GR minus weak-field is 4.1×10^{-6} (Sun) and 5.4×10^{-7} (galaxy). An isolated point Sun reaches only 0.71/0.67 of the law at $10^3 L_\star$, so whether a galaxy built from screened stars shows the law is open.

5 Every gate, with its frozen label

The full table, one row per gate, is closure_2026_11_06/CLOSURE_FINAL.md; PAPER69's verdict table is its deposited form and is carried unchanged. The rows below are the 2026-11-06 lanes that came after PAPER69.

Gate	Label	Lane
Equations and health		
Covariant field equations, v3s and AC2q	CERTIFIED vs direct variation	QX17, QX17A
Field equations with the adopted blend stiffness (general F)	CERTIFIED with $\Pi F'(\Pi)\, h_{mn}$ in the stress tensor; the $\Pi_+ \to F'$ shortcut alone FAILS the metric equation	QX17D
3+1 form, Lean kernels	Lean PASS; lane NOT CERTIFIED (numerical row, solver)	QX17B
Static spherical solver	CERTIFIED, both footings	QX17C
Π^6 repair, curved 3D spherical switch	PASS; adopted by the owner 2026-11-07 as the stiffness; non-radial modes untested	BL3
AC2q decelerating slow roots	HARMLESS, scoped; $\epsilon_\phi = 2$ NOT REACHED by any record history	EP1, EP2
Nonlinear fold, shock-capturing	NO VERDICT as frozen (viscosity check fails); no pair where AC2q regularizes	FS1
Nonlinear fold, viscosity-free reading	DIFFERS, both footings: the 80 clock-only pairs fold 1.2 % earlier (DIFF-TIME EARLIER, edge-carried); the soft and phantom pairs (160) read SAME-REACH; no row is REGULAR and no pair shows AC2q regularizing; all checks pass	FS2
Moving Sun	MS1, MS2: NO VERDICT, bracketed (largest time-mean on any rung 0.147 of the line). MS3: PASS (moving Sun, scoped), FINAL for the door, with no MS4; every cell, gate and bracket below the line, with the largest at 0.617 of it (G1 Mars, M490 REFLECT bracket). The PASS rests on the frozen time-spread classes (NOISE / OSCILLATORY), not on a converged extrapolation. Disclosed: the ultra runs used a different memory allocator setting (allocator only; the chunked solver reproduces bitwise)	MS1, MS2, MS3

Gate	Label	Lane
Light-cone causality	FAIL (physical, far below data); no repair in the khronometric class	AC18, SV2
All-gates battery (static gates G1–G8; G9 CMB)		
Six candidates on G1–G8	NOTHING PASSES ALL STATIC GATES; every candidate fails the Coma UDGs by at least $3.97\,\sigma_U$	GB1
Cold fluid: $\mathcal{K}(Q)$ dust, amplitude freed	FAIL: passes the background and CMB lensing, fails G1–G7; cosmic capture 4.111 in every class	KD1
Cold component decaying late	FAIL: CMB-lensing χ^2 46.3/53.0 against 18.48 at the longest lifetime discs allow	LD1
Hot thermal fermion, 11.1377 eV	FAILS 7 of 9 gates; CMB-lensing χ^2 166.0–194.2	HC1
Shock-entropy field (proxy)	FAIL: X-COP and groups; need falls outward while the measured adiabat rises	EN1
Shock-entropy sector in the action	FAIL by the no-tuning rule; static solutions isentropic (f $\leq$ 0.04)	SE1
The step as several causes		
Multi-cause step	MULTI-CAUSE IN PART: IMF explains the ellipticals and KiDS	ST1
Joint systematics budget per class	IN PART: ellipticals, KiDS and bright M31 dSphs EXPLAINED; X-COP +0.440, groups +0.232, Coma UDGs +1.120 NOT EXPLAINED (canonical medians); CMB SEPARATE	SE2
Environment, re-read	NO: no environment key passes; satellites need about 1.6× the phantom, Coma UDGs 4–46×	EV1
Disc-only exclusion	every direction DEAD; even a perfect (oracle) exclusion fails GB1	DX1
One action-native scalar for both walls (OQ2 kept by the owner)	NO VERDICT as frozen. At the frozen fallback $N_{\rm min} = 8$ the S1 reading is DEAD on both footings, every start: SPARC vs groups at $\log q$ 3.0–3.4, gap up to 0.497 / 0.483 dex. Web row YES: the CMB wall needs the law's phantom off on the web ($F_{\rm web,max}$ 0.00481 / 0.00435; with the kernel corrected 0.00798 / 0.00720, Section 6). SEPARATED NO, as frozen	OQ1, OQ2
Outer edge of the law from the switch	NO EDGE IN THE ACTION: for an isolated galaxy the static switch variable is $\Sigma/X = 2+\rho_{\rm ph}/\rho_{\rm DE} \geq 2$, so the phantom halo runs on until an external field ends it (the standard EFE radius, already graded)	outer-edge check
Owner-opened test lanes, 2026-11-07 (adopt nothing)		
Host-held dust with the measured EFE capping the wells	FAIL (clusters, 0/20 cells, both footings); with no dust the EFE moves the KiDS baseline misfit to +0.870	QX7 iter_6
Remnant tracing the hot gas	SHAPE FAIL, both footings: the missing mass is more concentrated than the hot gas	QX7 iter_7
Asphericity of hydrostatic masses in the law's field	NEGLIGIBLE (≤ 0.005 dex, wrong sign), both footings	QX32
Everywhere matrix (23 rows × 17 columns, every cell sourced)	NO ROW PASSES EVERY COLUMN. The minimal framework cover is three rows that contradict each other (KD1 dust, QX27 retention, a literature row); UDGs, causality with well-posedness, and a derived a_0 stay uncovered by every framework row. Two effective fields (BK, QUMOND β 2.5) FAIL GB1 G2–G7 on both footings; BK passes only the ellipticals (+0.123 / +0.103)	EW1
Law switched off on the single-stream web (khronon multi-stream key)	FAIL (stage 2, population phantom), both footings. KiDS is kept: all 59 points lie inside the law-on turnaround radius. But the phantom that the galaxy population must keep to hold KiDS sources CMB lensing at $q_{\rm pop}$ 0.936 / 1.079 of the dust target at z 1, against a line of 0.0354 (×26 / ×30; ×11 / ×12 self-consistent). This holds for any key that keeps the law on where KiDS needs it	WL1
Two-switch door (law switch plus charge-switched sector)	graded by WL1: FAIL; EW3 keeps the map row only, no graded number	EW3, WL1

Gate	Label	Lane
Fresh search for a new local action	no local action found; C1, C2 and C3 all FAIL, both footings. C1, any phantom multiplier of an absolute field length (the first absolute-scale grade): KiDS shift +0.283 / +0.260 against 0.05. C2, phantom screened beyond a second length: the natural local form does not screen, and the web forces that length to 0.3 Mpc or less (1.0 Mpc with the kernel corrected, Section 6), where KiDS fails. C3, two static branches: the static operator is not elliptic (radial), and the groups fail	FA2
Nonlocal, history and global actions	NO CANDIDATE SURVIVES, both footings. I1, causal memory of the phantom: FAIL on the web row (r 75.4 / 78.8; 19.1 / 18.8 with the kernel corrected). I3, formation-epoch a_0 memory: FAIL G2–G7 and the web row. I4, conserved phantom charge: FAIL on the web row, NO VERDICT on G1–G7. Disclosed by the lane: I1's MUTATE M1 did not bite as frozen, and peak RSS (1.87 GB) exceeded the frozen 1 GB line	FA3
AeST (v4) on the web, in hot wells and in discs	NO-GO FOR THE CLASS, both footings: one length $1/\mu$ cannot hold both the CMB-lensing amplitude ($\mu \geq 10\,\mathrm{Mpc}^{-1}$; r 1.037 / 1.045 at 10; $\mu \geq 0.63$ / 1.0 with the kernel corrected) and KiDS ($\mu \leq 0.063\,\mathrm{Mpc}^{-1}$); the gap is ×160 (Lean `window_empty`; ×10 / ×16 with the kernel corrected, the window still empty, Section 6). X-COP and groups FAIL at every computed μ: static dust $\leq 0.03\,M_b$ against a need of 2.39 / 2.07 M_b. The beyond-hydrostatic gate fails (0 % of X-COP radii). Primary CMB FAIL (read). At host-compatible μ, SPARC, Cassini, c_T and lensing = dynamics PASS. Two missed known answers are stated by the lane; no label moves	EW2
Late-formed compact baryonic objects (10^2–$10^7\,M_\odot$) as the hot-system half; the CMB wall untouched	FAILS (rows), both footings: X-COP shape, groups shape, Bullet. The budget is not excluded (lower bound 0.054 / 0.050 of the BBN baryons). No single formation rule: every key is S1 DEAD (gaps 1.50–2.53 dex). The dynamics window shrinks to 10^2–$1.8 \times 10^2\,M_\odot$. Objects tracing the stars miss X-COP's shape (rms 0.1208 / 0.1138), while an HSE-traced profile passes, so the needed profile lies between the gas and the stars. Groups pass 11 / 9 of 20 (canonical) and 10 / 8 (alt) against 16 needed. Disclosed by the lane: the Bullet child process peaked at 1188 MB, over the 1 GB line	NA2
Last two FA1 doors: disformal bulk-motion selector (I5), defect network of the a_0 scalar (I7)	I5 DEAD BY REDUCTION, both footings: every scalar the disformal metric allows reduces to dead keys (γ, X, ρ, p); with matter on the disformal metric, GW170817 confines BX to $[-6.2 \times 10^{-15}, +1.3 \times 10^{-15}]$, so the selector varies by at most 6.8×10^{-20}; no monotone weight passes S1; web lensing r 72–86 against 1.072 (27–39 with the kernel corrected). I7 DEAD, both footings: with η set by a_0 (2.240 / 2.463 meV) the network forms only at z of about 9; with η free, scaling networks track the background (w 0.167 at z 3400); the network is at most $3.7{\times}10^{-4}$ of the baryons at z 3400 and at most $1.3{\times}10^{-8}$ in clusters; gauge monopoles are particle-like. Disclosed by the lane: KA7 restated in amendment A1 before the rerun (labels unchanged); MUTATE M2 and M4 bite only weakly; the first run peaked at 1.025 GB, the rerun at 0.858 GB	NA1

Gate	Label	Lane
The a_0 law plus an 11.1377 eV thermal fermion (test lane; the no-particle rule relaxed for this lane only)	does NOT pass every EW1 column, both footings. Run as a consistency test it reproduces the literature: X-COP 6/7 consistent, $M_{\rm req}/M_b$ 1.8–3.4, using 32–62 % of the supply, phase space never binding; run as a prediction (HC1) X-COP fails. Groups FAIL (14/20 / 11/20 against 16; the law over-predicts the shell); ellipticals FAIL 0/7 on phase space; AGC 114905 and Malin 1 FAIL for any fermion; KiDS G6 −0.098 dex FAIL; primary CMB $\Delta\chi^2$ +2716 / +2599 FAIL; CMB lensing, Coma UDGs and the colour split FAIL. SPARC PASS (not robust); Cassini and c_T PASS only with v3s/AC2q as the law's action. Disclosed by the lane: MUTATE M1 (mass halved) has NO POWER on X-COP, and the X-COP loader holds 7 clusters, not 12	EW4
KiDS isolated-lens kill of the late-heated window (QX7 iter_4), as a joint systematic budget	NO VERDICT as frozen (the 4× MUTATE cannot bite on cell C3); no cell softens, both footings. KILL HOLDS on C2 and C4, NO POWER on C1 and C3. The hot gas measured by the kinetic SZ alone accounts for the raw +0.15–0.20 dex excess at 100–300 kpc; the posterior excess of data over the law is −0.21 to +0.05 dex (canonical) and −0.25 to +0.01 (alt), and the implied line is 0.05–0.24 dex (canonical) and 0.00–0.20 (alt), against the window's 0.17–0.25. Disclosed: the lowest mass bin uses an extrapolated gas row, and canonical C4's hold rests on it	KX1
SE2's two fragile rows, re-graded	Coma UDGs NOT A TEST (2 of 9 admissible at pericentre $d/3$; no orbit on disk); groups NOT EXPLAINED (+0.259 / +0.221), fragile; the residue is X-COP + groups + CMB lensing	UA1
Late-heated window on the cluster gates (QX7 iter_4 cells C1–C4)	FAIL on every cell, C3 included, both footings. Every cell passed CMB lensing as frozen (χ^2 8.70–11.69; with the kernel corrected that gate reads NO VERDICT, Section 6), $f\sigma_8$ at $z < 1$ (R_f 1.000) and the forest at z 2–4, and fails X-COP's profile, the groups and the Bullet. C3: X-COP +0.324 / +0.287 (inner bin +0.48: what evaporates from galaxies evaporates from clusters too), groups 0–1 of 20, Bullet FAIL (the sector is a fluid at Mach 1.7–1.9, ram-stripped, and follows the gas). Disclosed: the KX1 narrowing to C3 arrived after the numbers (C3 was frozen and graded as frozen); the Bullet child peaked at 1.48 GB, over the 1 GB line	CE2
Residual surface: does any pair of measured variables carry the law's missing mass across classes?	NO VERDICT as frozen: the Newtonian-f MUTATE does not bite, because $g_{\rm obs}$ was on the frozen variable list (from the brief) and makes the law-off f an exact function of $(g_N, g_{\rm obs})$. Main reading NO PAIR, both UDG readings. The closest pair, $g_N + g_{\rm obs}$ (χ^2/N 0.433 / 0.440), is circular and fails canonical leave-one-out. The best pair without $g_{\rm obs}$, depth + σ (χ^2/N 1.483 / 1.434), leaves SPARC at z −4.49 / −3.87, X-COP at +3.82 / +3.48 and the UDGs at +3.46 / +3.48; without $g_{\rm obs}$ no pair passes leave-one-out after dropping any class. SPARC is GB1's 145 galaxies	RS3
The kicked carrier, re-graded (RESOLUTION, 2026-11-07)		
Does the action release a kick when its switch changes branch?	NO RELEASE TERM, both footings: no isolated host changes branch at virialisation or at the acceleration crossover (min $\rho_{\rm ph}/\rho_{\rm DE}$ +0.000507 / +0.000462), and the blend gives zero latent energy at the branch point. Any kick is a postulate, not the action's	CK2
September's kicked collisionless carrier at today's needs, v_k 500–650 km/s	FAILS (rows) at every v_k on both footings; the binding gate is the SPARC high-mass tail, where every galaxy with $v_{\rm flat} \geq$ 250 km/s keeps the carrier. The kick v_k is hand-set	CK1
Cosmology under the kicked carrier	NO VERDICT (the CMB-lensing gate has no power); among gates with power, v_k 550 passes and 575 / 600 fail the forest	CK4

Gate	Label	Lane
A cold component born clumped at the future nodes, never in galaxies	FAIL, both footings, every clump mass: the primary CMB tolerates a bias of only 0.99670–1.00142, while a peaks-born population carries 45–1866 (bias 1.2 / 1.5 / 2.0 give Δ +4.10e4 / +5.57e5 / +6.35e6); the forest allows at most 0.13–0.28 % of the component in clumps; the clumps' own Poisson power fails CMB lensing at 10^{14} / $10^{15}\,M_\odot$ (+75.9 / +239.8) and passes at 10^{13}	NC2
Do cluster-member early types carry the component?	UNRESOLVED by rule, both footings: members sit +0.011 [−0.044, +0.065] / +0.012 [−0.043, +0.068] dex from field early types at fixed stellar mass and dispersion, so at most about 16 % extra mass at $r_{\rm half}$; stripping cannot hide a residue; the KiDS isolated-lens control is clean; CLASH's lensing-only slope leans to an early clump's sign with every interval containing 0	NC3
The born-clumped clump spectrum (a record row; NC2 closed the candidate)	FAILS (rows) on both footings, binding K3: hosts inside R_{500} hold 0.0133 of the cosmic baryons and so 0.0091 / 0.0078 of the cold component (0.025 / 0.022 in the most generous bracket); clumps would need 110 / 128 times the observed host abundance (40 / 45 generous); even the ceiling with every baryon in a host is 0.75 / 0.68. K5 also fails: the top-hat collapse in the law's field fits X-COP at rms 0.166 / 0.143 at best. A fitted mass trend passes (slope 0.6–2.3 / 0.8–2.2 per dex). Disclosed: the Schmidt test never passed ($\langle V/V_{\rm max}\rangle$ 0.403–0.465), and a factor 2–3 would still leave the shortfall ≥ 13	NC4
Is the CMB's cold component independent evidence? (F1, audited with own likelihood code)	F1 CONFIRMED, both footings: baryons alone fail the primary CMB by Δ +172,915.6 / +166,674.4; the cold fit is adequate ($\Delta_{\rm cal}$ −65.21 / −65.07 against $\chi^2_{\rm cal}$ 3805.37) at ω_d/ω_b 5.410 / 5.412 (0.973 of 5.562); an own Newtonian-gauge code reproduces that the khronon clock cannot cluster ($Q \leq 3.4\times10^{-11}$ at z 1100). Disclosed: CAMB is the Boltzmann code, as in QX15 (CLASS differs by 3.8); the baryons-only fits sit at bracket edges; one check peaked at 1.81 GB for 10 s	F1A
Can the action's own dust term ($\mathcal{K}(Q)$, at KD1's amount) form the cluster core?	CANNOT FORM CORE, both footings, for every μ_K in 0.1–10^4 Mpc^{-1}: in the law's $1/r$ field the hydrostatic dust falls like $\ln(r_s/r)$ with a hard surface, so one Jeans length sets both core and extent. A 50–100 kpc core comes with a surface near 160 kpc (concentration 1.000, inner bin over by 1.1 dex); a concentration near 0.25 needs a 360–400 kpc core; the best per-bin rms anywhere is 0.226 / 0.222. The closest profile (μ_K 19.05) is 4.67×10^5 below the BBN bound, and galaxy capture stays at 4.111 / 4.112. A planted cored isothermal passes the same machinery (rms 0.091)	KQ1
Does the fold machinery show a release layer at 500–650 km/s? (the outside solver's kick identity, $v_k = 2c(a_0L_\star/c^2)^{1/3}$ = 577.6 / 571.3 km/s)	NO VERDICT (check failed): the planted 577.6 km/s control did not read back in the band (it regularizes late and reads 317.7 / 309.2 km/s). The graded rows read $v_{\rm post}$ 0.024–0.157 km/s and R_δ 5×10^{-10}–3×10^{-7}, all 160 readings outside the bands; the speed tracks the infall ($d\ln V/d\ln v_f$ 1.000 / 0.95) and doubling $L_\star$ leaves it unchanged (ratio 1.000, where the identity predicts 1.260). Disclosed: the brief's $\lambda_{\rm crit}$ was off by a factor 10 (22.53 / 24.04 kpc, not 2.09 / 2.25); the reduction is non-relativistic ($1-X/X_{\rm bg} \leq 2.4\times10^{-7}$), so the identity's relativistic regime cannot appear	AK1

The CMB requirement, anatomised, and two owner-named lanes (2026-11-08)

Gate	Label	Lane
Which features of the primary CMB demand the cold component? (anatomy of F1A's baryons-only Δ)	NO VERDICT as frozen: the M1 swap moves H1/H2 by $+343\,\sigma$ and H3/H2 by $+197\,\sigma$ in the same direction, ratio 0.57 inside [0.5, 2] (the third peak is a compression peak too). Block C CLUSTERING PINNED, both footings. The baryons-only deficit is spread over TT, TE and EE and every multipole range; no piece carries more than 19 %. A smooth matter-like density (ω_X 0.108, about 5× the baryons) that sets equality and the expansion but does not cluster recovers 79 % (Δ_E +35,812 / +35,378); raising ω_b alone recovers none. The rest needs still potential wells through z 1100: without clustering H3/H2 misses by $-36.8\,\sigma$, H1/H2 and the first peak by $-16\,\sigma$; the cold fit meets every feature within $1.1\,\sigma$	CMB1
Before recombination, does any degree of freedom other than the clustering dust supply a still Weyl potential ($\geq$ 0.5 of the dust's at k 0.03–0.3 Mpc^{-1})?	NO, both footings. The khronon capped at the galaxy allowance: still share 0.043 / 0.024; the clock about 10^{-20}. The P sector has no propagating mode (Hessian rank 0). The matter equations carry no khronon field, $c_s^2 = 1/(3(1+R))$ exactly; the slip is at most 7×10^{-4} and follows the source. The law at $a_0(z)$ 1–100 times today's: its phantom follows its source (r 0.992–0.997), still share at most 0.28 / 0.30. Open, stated: in a toy oscillator a following phantom at 30 times or more raises the odd peaks (H3/H2 1.30 / 1.36 at 100 times, against the dust's 1.447); that needs ρ_{DE} about 900–10^4 times today's at z 1100, and no history in the record gives it	CMB2
F1A's G4 starting state (amendment A2, prompted by CMB2)	G4's printed contrasts depend on the starting time; a start consistent with the Poisson constraint converges (−26.65 / −94.13 at k 0.05 / 0.3); $Q_{X1} \leq 3.7 \times 10^{-12}$; G4 REPRODUCED, F1 CONFIRMED unchanged	F1A A2
Is there one vacuum-response length in the action whose identity fixes $\kappa = \frac{1}{2}$?	NO IDENTITY, both footings. The khronon kinematics carry the 2 exactly ($\nabla \ln X = -2A/c^2$), but setting the response length at a_0 equal to a vacuum length gives $\kappa^2 = 1/(4q)$, which is $\frac{1}{2}$ only for $q = 1$, the π-free length $c/\sqrt{G\rho}$ already posited (KF1's $\xi = 1$). The six GR lengths give 1.447, 1.772, 0.921, 0.282, 2.507, 0.230; the cancellation identity has no root; equipartition gives 4.525 / 6.399; the blend identity depends on host mass. The vacuum's response carries at most 6.9×10^{-8} of the cluster need. Lean: 7 theorems, MUTATE twin fails 4/4. κ stays fitted	VR1
Settling-age dependence of the action's dust (a reduction; no criteria, no script, no number)	NO TERM IN ACTION: $\mathcal{K}$ depends on Q alone and the field equations are local in time. Every route by which a settling age could act (capture, hot early, born clumped, kick, late heating, self-expulsion, formation-epoch and merger keys, local keys, a perfect oracle) is a closed cell on both footings, and the Coma UDGs, old and settled, carry the largest need in the record	SA1
The a_0 law with κ free plus the CMB's cold component (quarantined test lane: the owner lifted the no-cold-component rule for HY1 only)	NO VERDICT / NO VERDICT on the frozen M3 rule: doubling the dust worsens discs and cluster cores together, so the control cannot bite (a gap in the control's design, disclosed). Under it, FAIL on both footings. The record's own dust dynamics (KD1 model P at the one μ_K BBN allows) put the component in every galaxy and cluster as a central lump of about 4.1 times the baryons: SPARC rms 0.465–0.658 / 0.464–0.680 against 0.110 at every κ, and X-COP cores overfilled by about 1.1 dex at every κ and any amount. A weaker law (κ in [0.05, 0.17] / [0.05, 0.13]) holds KiDS, the group step and the Solar System, but only if discs shed at least 96 % of the dust, which is the second fact again. Bullet and the law on the linear web NOT EVALUABLE. Adopts nothing	HY1

Gate	Label	Lane
Assumption audit of the inference that the peaks need a cold component of about 5.4 times the baryons (A1 law inert at z 1100, A2 lensing, A3 recombination and $N_{\rm eff}$, A4 initial conditions, A5 the identification with galaxy dark mass, A6 nuisance templates)	The peak inference HOLDS; only A5 BREAKS, both footings. A1: the frozen line reads BREAKS POSSIBLE (the baryons-only potential gradients sit in the law's deep regime), but the law's response follows the fluid and leaves H3/H2 at the baryons-only value (CMB2 D4), so HOLDS / HOLDS. A2 lensing HOLDS (unlensed theory: survival 0.993; template amplitude free: 0.877 / 0.862, where the cold fit picks A_ϕ 1.00 and baryons alone 9.50 / 9.45). A3, $N_{\rm eff}$ and YHe free, HOLDS (0.892 / 0.871). A4, n_s and running free, HOLDS (0.901 / 0.895). A5, the CMB's component is the rotation-curve mass, BREAKS / BREAKS on the record's own data (SPARC, KX1, NC3; bound hosts hold 0.9 % / 0.8 %). A6, nuisances widened and priors off, HOLDS (0.751 / 0.736) but is nuisance-sensitive (-25 / -26 %, through unphysical foregrounds). The smallest baryons-only Δ under any single relaxation is +122,711. Named, not computed: the cold fit sets $\nu = 1$ while its own gradients at z 1100 give $\nu - 1$ 0.05–0.17, a few-percent debt in R	CMB3
Lanes opened after v1.6 (2026-11-08)		
A running K-mass from a non-quadratic $\mathcal{K}(Q)$ (test lane; Blanchet–Skordis's own DBI form, λ fixed by the corrected-kernel growth line $M(z\,3) = 630.957\,\mathrm{Mpc}^{-1}$)	FAIL / FAIL. The mass runs the right way (read $\mu_K(0)$ 1.232–79.65 Mpc^{-1}; 6.01×10^{13} at z 1100) and passes BBN with no stiff phase, cold at z 3400, the primary CMB, corrected-kernel lensing, $f\sigma_8$, the forest and c_T. It fails because, for any convex $\mathcal{K}(Q)$, the mass is set by the sector's own local density: dust turning around in a galaxy sits at the background density of z 4.31, so it carries $M \geq 631$, a Jeans length of 2.8–2.9 kpc against the 234 kpc turnaround radius, and is captured (SPARC rms 0.777 / 0.789 against 0.146 / 0.143). Three of the four members also have no stationary branch at the khronon charge that bound systems carry. A fix-forward note in KR1's RESULT (ef1a9888b1) grades condition (iii) against KR2's empty window: (iii) fails for every member and the brief's kill fires on both footings; the label is unchanged	KR1
A present-day μ_K window for the CMB's amount of the action's own dust, with no extra dust inside $1/\mu_K$ (test lane)	FAIL / FAIL (canonical empty at X-COP; alt empty at SPARC by the frozen $T = 0$ clause, and at X-COP under the T-carried reading). No $\mu_K(0)$ in 0.1–79.43 Mpc^{-1} passes. X-COP is hollow where the dust starts beyond 100 kpc and over-filled by 0.89–1.23 dex where it starts inside (best per-bin rms 0.439 / 0.425 against 0.10; 0.225 / 0.231 at the captured amount, KQ1's 0.226 again). Groups need μ_K in [1.585, 3.98] and the host capacity (NC4) needs at most 0.631: disjoint. SPARC is clean to 6.31 and KiDS to 3.98; CMB lensing at constant μ fails everywhere ($\Delta\chi^2$ 309–1315, GL1's wall). Bullet NOT EVALUABLE. Post-freeze, graded apart: with KR1's local-density mass no column's window survives (SPARC and groups carry $M \geq 2173$)	KR2
Lanes opened after v1.7 (2026-11-08)		

Gate	Label	Lane
A cold-gas conversion key: the $\mathcal{K}(Q)$ dust phase converts to the sector's smooth phase at rate $\alpha\,\rho_{\rm cold}$ (baryons below 10^4 K), one declared parameter α scanned over 10^{-3}–10^3 $(M_\odot\,{\rm pc}^{-3}\,{\rm Gyr})^{-1}$ (quarantined test lane: the owner launched BG1 on the words "Launch it anyway"; the coupling is not derived from the action)	FAIL / FAIL: no α in the scan passes G1–G10 on either footing. The galaxy side works near $\alpha = 10^3$ on canonical: SPARC rms 0.0626 (line 0.0660), the gas-rich low-density discs, KiDS (shift -0.003 dex), the ellipticals under the IMF reading and the dSph pair at the line all pass; discs retain 0.4 % of the dust while quenched hosts keep 17–36 %. The same rate converts 97 % of the component by z 20, because the neutral mean-density gas of the dark ages is the densest cold gas in the history ($7\,M_\odot\,{\rm pc}^{-3}$ at z 1000; its exposure is 1/7 of a disc's): the dark-ages gate holds only for $\alpha \le 2.5$, the forest and growth only for $\alpha \le 6.3$, and discs need $\alpha \ge 1000$ (beyond the scan on alt); the windows are disjoint by a factor 400, and an exposure $\times 10$ replay moves both alike. At every α on both footings three hot rows fail: Coma UDGs under-filled by 0.65–0.92 dex (the full component gives $b = 6.4$ where they need about 80; the kill named before launch fires), groups at most 7/20, X-COP over-filled by 0.06–0.10 dex at the CMB amount $R = 5.41$ (it would pass at KD1's captured share $R = 4.11$, reported); clusters keep at least 94 % under a hot ICM. MUTATE M1 (hot catalyst) and M2 ($\alpha \to 0$) bite; M3 ($10\times$ the best α) does not bite as frozen (the UDG row is one-sided), and the label does not rest on it. Adopts nothing	BG1
A potential-gated biharmonic Jeans scale for the $\mathcal{K}(Q)$ component, $[1 + R_J^4\nabla^4]\rho_K = \rho_b S(\Phi)$ (an outside proposal, graded as a quarantined test lane)	FAIL / FAIL: 0 of 756 cells per footing (252 grid points $\times$ 3 values of ℓ_θ, two R_J forms) pass G1–G10, and the as-written point passes only G10 (Cassini). The gate cuts the weight across the acoustic range, because the linear potential at recombination falls from 2.6×10^{-5} at k 0.01 to 5.9×10^{-7} at k 0.3, straight through Φ_*: primary CMB $\Delta\chi^2$ +619.5 / +613.5 before refit and +171.7 / +170.9 after a refit of the 5 cosmological parameters. KiDS outer bins are filled (shift 0.42–0.44 dex); at X-COP the component is no more concentrated than the gas (0.209 against 0.202, need 0.267) and over-fills at the CMB amount; on the Bullet it tracks the gas. The best point (R_{J0} 2.5 Mpc, Φ_* 3×10^{-7}, p 2, ℓ_θ 0) passes 4 of 11, where the gate barely acts; the forest and growth reject every R_{J0}. MUTATE M1 and M2 bite; M3 ($R_{J0} \times 10$) does not bite as frozen and the label does not rest on it. Disclosed: a mass-conserving divergence-form operator for variable R_J; the Bullet graded under the record's instantaneous relation. Adopts nothing	PJ1
Does the law's plain cluster residual $\rho = \log_{10}(g_{\rm obs}/g_{\rm law})$ have the same radial shape on a lensing-only leg (CLASH: Tian+2020's 84 points, 16 X-ray-selected clusters, Umetsu+2016 strong + weak lensing, Donahue+2014 gas + BCG) as on the X-COP hydrostatic leg (7 clusters with measured stars)?	SAME SHAPE / SAME SHAPE, in every bin the two legs share. Lensing minus hydrostatic: +0.048 [−0.026, +0.114] dex at 0–0.25 R_{500} and +0.053 [−0.022, +0.140] at 0.25–0.5 (canonical; +0.049 / +0.056 alt), inside the frozen 0.10 band with the 95 % interval covering 0; every bracket within 0.04 dex. The residual inside 0.5 R_{500} is not a hydrostatic artefact, and the lensing leg shows slightly more missing mass, not less. Reported: the lensing-leg need at 0.25–0.5 R_{500} is $\times$3.28 [2.90, 3.90] against X-COP's $\times$2.90. The lensing leg ends at 600 kpc (0.28–0.59 R_{500}), so its need at R_{500} is not delivered; the two legs are different clusters, and CLASH's baryons are gas + BCG only. MUTATE M1–M3 bite	HL3

Gate	Label	Lane
A merger-decohered component (an outside proposal, graded as written as a quarantined test lane): 4.40 times the baryons, clustering as cold dust before shell-crossing, afterwards a 1.6 Mpc Thomas–Fermi core around every collisionless centroid, the core lowered in groups and clusters by a merger-driven decoherence scalar, satellites keeping $(v_{\rm esc}/\sigma_{\rm host})^3$ of their share	FAIL / FAIL: 4 of 11 gates pass on each footing (SPARC, the massive tail, growth with PR4 lensing, Cassini); G2–G7 and G9 fail on both. KiDS: max shift 0.297 dex at 1.7 Mpc, 15 of 59 points over. X-COP: inner bin -1.132 dex with $4.38\,M_b$ inside 100 kpc (concentration 3.34 against the gas's 0.20, need 0.27). Groups 3 / 4 of 20. Coma UDGs $+1.240$ dex (capture 4.9×10^{-6}, and Coma's own component raises the host field $\times 1.72$, worse than the law-only $+1.121$). Bullet, the component as its own Mach-6 fluid: peak 51 / 41 kpc from the gas. Primary CMB $\Delta\chi^2$ +3682.7 before and +1524.8 after a 5-parameter refit. Forest $\Delta\chi^2$ 216,405. Disclosed: the proposal's dynamic integrator, rebuilt from its equations, does not reproduce its own decoherence values (KA3 FAIL), so the quasistatic form is graded alone; the dynamic bracket passes no gate the as-written fails. MUTATE M1–M3 bite. Adopts nothing	WD1
Is a hosted galaxy's static external-field phantom swept by its host's khronon wind, as AC15 and MS3 found for the Sun's hedgehog? (the action's own moving-body state, no new parameter; quarantined test lane)	FAIL / FAIL: the sweep acts, and against the theory. With W live the full screened system has no Bernoulli integral; the frame streams through the member ballistically and carries the member's own static phantom downstream with the host's: 99.86–100 % of it leaves the readout sphere in every run, so the moving member is Newtonian to 2×10^{-3} dex. $D = \log_{10}(g_{\rm dyn}/g_{\rm static})$ is negative on every Coma UDG row (-0.401 to -0.229 canonical, -0.453 to -0.266 alt) and every MW classical dSph row (-0.906 to -0.577 / -0.936 to -0.610); the isolation fraction F is negative on every row (-0.646 to -0.287 / -0.748 to -0.333 on the UDGs; FAIL line 0.3). On SE2's path the Coma UDG shortfall grows from $+1.121$ to $+1.412$ dex ($+1.076$ to $+1.409$ alt); on X1's rows the MW classical dSphs go from $+0.356$ to $+1.208$ and the bright M31 dSphs from $+0.633$ to $+1.555$ (canonical). Host rows (X-COP, groups) unchanged. The Sun stays under every AC15/MS3 line (worst 0.215 of the Mars line). MUTATE-1 and MUTATE-2 bite; MUTATE-3 is detected, but C2 has no power at a 0.5 dex plant. Disclosed: the linear kernel is extrapolated for four dSphs and the member proxies; perpendicular $g_{\rm ext}$ not run. Adopts nothing	XF1
Lanes opened after v1.9 (2026-11-09)		
Does the law's residual key on measured spin? Slow minus fast rotators at matched $M_\star$ and g_N on MaNGA DynPop (6,056 galaxies, 536 slow rotators; quarantined test lane)	FAIL / FAIL: $D = +0.0602$ [+0.0369, +0.0811] dex canonical, +0.0601 [+0.0369, +0.0812] alt, the whole interval under the +0.10 FAIL line (hit line +0.20). The excess is a 0.03–0.06 dex step confined to $\lambda_{R_e} < 0.2$ and flat above, not a trend with spin (Spearman -0.486; a key needs ≤ -0.6). Matched on $(M_\star, \Sigma_{R_e})$ instead it falls to +0.0096: dispersion carries it, the key HX1 graded NO LADDER, and KU2's UNIVERSAL on the same file is recovered. Post-freeze, 30 JAM-model × IMF readings (not graded) span +0.014 to +0.198; the top reading uses a model the data choose for 0.6 % of galaxies. Known answers pass; MUTATE bites (spin shuffled: +0.012; a +0.2 dex plant returns +0.2000; the other mass column keeps the label). The lane's frozen footings are a_0 1.2 / 1.1×10^{-10} m s^{-2}; at fixed g_N the two agree to 10^{-4}. Adopts nothing	SK1

Gate	Label	Lane
Does the cluster missing mass follow one baryon tracer (gas, stars, gas + stars) or the hydrostatic total, at one amplitude per cluster? X-COP, 7 clusters with measured stars; two readings, D = $M_{\rm dyn} - M_{\rm law}$ and S, the extra source inside the law (grading lane of the adopted law)	NO TRACER FOLLOWS / NO TRACER FOLLOWS, on both readings (FOLLOWS iff rms $\leq$ 0.10 dex and max $\lvert\mathrm{SHAPE}_b\rvert \leq 0.15$ dex). The missing mass is more concentrated than the gas, gas + stars and the total (inner bins +0.15 to +0.48 dex) and less concentrated than the stars (-0.24 to -0.40 dex in the inner bin; -0.8 dex inside $0.05\,R_{500}$, reported, not graded). The closest tracer is the hydrostatic total, with an amplitude the same in every cluster (SD($\log A$) 0.05–0.06 dex). On reading D (canonical) the missing mass is about 70 % of the total inside $0.25\,R_{500}$, falling to about 45 % at R_{500} (0.71, 0.66, 0.58, 0.46 by bin), and 0.32 in the 1–2 R_{500} bin, which is the forward-model hydrostatic mass; reading S gives 0.60 falling to 0.16. The D label is bracket-sensitive: inside R_{500} the total FOLLOWS (rms 0.077 / 0.090), so the label rests on the 1–2 R_{500} bin. CLASH's lensing total gives an amplitude of 0.71 / 0.69 over its two inner bins, beside X-COP's 0.71 and 0.66 in those bins: an agreement inside $0.5\,R_{500}$, with low power. QX29 B1's X-COP pass is reproduced as a total-mass read over 0.1–0.7 R_{500} (rms 0.071); the same model's component misses by -0.39 and -0.68 dex in the 0.75–1 and 1–2 R_{500} bins (r4, added after the first run, reported). Known answers pass; every MUTATE bites. Adopts nothing	TR1
Rows touched by the kernel erratum	the CMB-lensing and web-lensing numbers of KD1, SE1, CE2, NA1, FA2, FA3, EW2, OQ1 / OQ2 and SV5 above were computed with the Limber kernel corrected in Section 6; the corrected values are listed there and, by the owner's decision of 2026-11-07, are the standing labels; every row above keeps its FAIL or NO-GO	GL1, GL3, GL4

6 Erratum to v1.0: the CMB-lensing Limber kernel

The Limber integrator shared by the sector and web lanes (QX3 iter2, `limber_ratio`) omitted the lensing potential's k^{-4}. It weighted each redshift by an extra χ^{-4}, so the ratio $S(L)$ was dominated by $z \sim 0.02$–0.1. CK4 found it, the code was read independently, and GL2 CONFIRMED it with its own code against CLASS: with k^{-4} restored the Limber shape over CLASS's own spectrum is 0.98–1.006 at L 20–400, against 0.045–33 without it. GL1 (sector side) and GL3 (web side) re-ran every affected lane with the kernel corrected; re-inserting the error reproduces the committed numbers (to $\leq 4.5 \times 10^{-16}$ on the web side), and a wrong power (k^{-2}) is the mutation. WG1–WG6, LD1 and HC1 used separate pipelines and are untouched. **The owner accepted the corrected values as the standing labels on 2026-11-07 (closure_2026_11_07/OWNER_DECISION_2026-11-07_ACCEPT_CORRECTED_KERNEL_LABELS.md). Old and corrected values stay side by side below, and the frozen RESULT files are not edited.**

Label (lane)	Old	Corrected	Reading
Wall 1, CMB side (KD1 G9a, SE1 arm (a))	μ_J 2.51×10^4	μ_J 631; the binding side is BBN (μ_B 8.89×10^6, unchanged)	HOLDS
QX3 iter2 (no window)	$\mu_{\rm min}$ 3.98×10^4 vs $\mu_{\rm max}$ 79.43	$\mu_{\rm min}$ 631, still above 79.43	HOLDS
QX3 iter3 (class no-go, 35 cells)	best $\Delta\chi^2$ 92.59, 0/35	best 2.78, 1/35 (β 0.80, μ_0 1259)	re-stated (owner 2026-11-07)
QX7 iter_3 (no window, 31 cells)	closest 17.87, 0/31	31/31 inside the lensing line	re-stated (owner 2026-11-07)
QX7 iter_4 (FAIL on clusters)	window 270 cells, clusters 0	window 857 / 857, clusters 0 / 857	HOLDS
QX10 iter_1 (NO VERDICT)	J1 $\Delta\chi^2_{TT}$ 4.30 / 1.05 / 77.2 / 3.47	$\leq 10^{-7}$ on all four cells	re-stated (owner 2026-11-07) (PASS by its own rule)

Label (lane)	Old	Corrected	Reading
CE2, CMB-lensing gate	PASS, χ^2 8.70–11.69	$\Delta\chi^2$ 0.000 on every cell; its mutation no longer bites	gate re-stated (owner 2026-11-07) (NO VERDICT); CE2's FAIL on X-COP, groups and the Bullet is kernel-free and HOLDS
CK4, CMB-lensing gate	NO VERDICT	NO VERDICT	HOLDS
SV5 iter_1, web amplitude W1 (line 1.072)	r 74.74 / 89.36	32.53 / 38.66	HOLDS (FAIL)
SV5 iter_1, web shape W2	$\Delta\chi^2$ 0.019, PASS	33.5 / 34.1, FAIL	re-stated (owner 2026-11-07) (stricter)
SV5 T0.2, latest z for a late Jeans length of 30 / 100 / 1000 Mpc	none / none / none	2.0 / 1.5 / 0.75	re-stated (owner 2026-11-07)
SV5 iter_3, $a_0(z)$-tied Jeans scale, H-native	M^* $\Delta\chi^2$ 876.7, FAIL	0.016; K-mono 0.52	re-stated (owner 2026-11-07) (WINDOW-CONDITIONAL; growth and lensing gates only)
WL1, population phantom line	$q_{\rm pop}$ 0.936 / 1.079 vs 0.0354	identical (kernel-free)	HOLDS
NA1 I5, web r	72.2–89.4	27.4–38.7	HOLDS (FAIL)
FA3 I1, web r	75.39 / 78.77	19.12 / 18.75	HOLDS (FAIL)
FA2 C2, web cut-off $L_{\rm max}$	0.3 Mpc	1.0 Mpc; KiDS still fails there (-0.074)	HOLDS
EW2, web window	μ_w 10, gap $\times$160	μ_w 0.63 / 1.0, gap $\times$10 / $\times$16	HOLDS (window empty)
OQ1 / OQ2, web row	$F_{\rm web,max}$ 0.00481 / 0.00435	0.00798 / 0.00720	HOLDS (YES)
QX28 G6b	$\Delta\chi^2$ 140.1 / 70.0, FAIL	0.006 / 0.004, PASS	re-stated (owner 2026-11-07); G6 still FAILS on the kernel-free G6a
QX27, QX29, QX31	FAIL; PASS (A-web conditional); ROBUST	labels unchanged	HOLDS
QX15 cells C1 / C2 / C4 (found by F1A)	$\Delta_{\rm cell}$ +0.66 / +0.27 / +0.57, PASS	$+1.1\times10^{-5}$ / $+6.5\times10^{-6}$ / $+1.1\times10^{-5}$, PASS	HOLDS
QX15 cell C3	$\Delta_{\rm cell}$ +5.95 / +6.03, TENSION	$+4.1\times10^{-5}$ / $+4.2\times10^{-5}$, PASS	re-stated (owner 2026-11-07)
QX15 P1 power control	+192.46 / +189.91, bites	kernel-free, reproduced; bites	HOLDS; QX15's main fits (F1) do not use the kernel

What moves is late, small-scale suppression: with the kernel corrected, PR4's shape at $L \leq 400$ is nearly blind to it. No move opens a door by itself, because the cluster, group and KiDS rows that close those doors are kernel-free. Wall 1 holds on its BBN side; on the CMB side alone the galaxy-versus-lensing gap narrows from $\times$316 to $\times$7.9.

7 The CMB requirement, anatomised

CMB1 asks what the primary CMB's demand for a cold component is made of; CMB2 asks whether anything in the action other than the clustering dust can meet it. Both run on the framework background, with no ΛCDM input, on both footings.

1. The deficit of baryons alone (Δ +172,915.6 / +166,674.4, F1A) is spread over TT, TE and EE and every multipole range; no piece carries more than 19 %. Raising the baryon density never helps.
2. About four fifths is the expansion history. A smooth matter-like density of about five times the baryons, which sets matter–radiation equality and the distance to z 1100 but never clusters, recovers 79 %.
3. The last fifth (Δ_E +35,812 / +35,378) is clustering. The third-to-second and odd/even peak heights and the

first-peak position need potential wells that stay still through recombination, made by a component that does not feel photon pressure.

4. Nothing in the action except the clustering dust supplies such wells (CMB2). The khronon, capped at what galaxies allow, gives 0.043 / 0.024 of the dust's still potential; the P sector has no mode; the preferred frame does not enter the photon–baryon equations; and the law's phantom follows its source.
5. The law is inert at z 1100 on the record's own numbers. With $\rho_{\rm DE}$ constant its peak ratios are those of baryons alone (H3/H2 1.008 vs 1.006). The one open corner, a_0 at z 1100 thirty or more times today's, needs a dark-energy history 900–10^4 times today's at z 1100; even such a history leaves the action inert (X35, below).

So the peak inference holds on the framework's own background: at recombination there is a pressureless, clustering component of about 5.4 times the baryons, and no term of the action other than its dust supplies it. What the record breaks is not that inference. It is the identification of this component with the agent that flattens galaxy rotation curves: galaxies hold almost none of it today, and bound hosts inside R_{500} hold about 1 % of it (NC4). CMB2's open corner is already answered in the record by X35 (rebuild_2026_10_22/X35_a0_tracks_ede; UNDECIDED by its frozen rule). Early dark energy does put a_0 at z 1100 at 1,147–7,565 times today's, but in the covariant action the extra potential is the weight of khronon charge, which stays identically zero without a separate cold component, so the potential is the GR potential of baryons, photons and neutrinos for every a_0 (D1 NULL; a nonlinear 1-D solve from a non-zero start returns $\max|\Xi|$ 4.9×10^{-29}). The odd-peak rise of CMB2's toy appears only in the static AQUAL reading, which is not the action, and that reading makes the photon–baryon fluid Jeans-unstable at first-peak scales ($k_{\rm J}$ 0.024–0.039 Mpc^{-1}). The full Boltzmann check CMB2 named is therefore not owed. Early dark energy is not neutral late, though: through the sound horizon it moves CFG689's native $\sqrt{X}$ at z 2.33 from 1.00 to 1.49 (a flat-Λ estimate).

CMB3 audited the inference's assumptions one at a time, on both footings. Lensing (A2), recombination with $N_{\rm eff}$ and YHe free (A3) and the initial conditions (A4) each keep at least 0.862 of the baryons-only deficit and move the amount R by at most 2.2 %; the widened nuisance block (A6) keeps 0.736, through unphysical foregrounds. The law at z 1100 (A1) is graded through CMB2's timing result, and recombination has no framework-native change (CMB2 D3). Only A5 breaks: the identification of the component with the mass that flattens rotation curves, contradicted by the record's own galaxy data. Taken together, the three CMB lanes say that the inference holds on the framework background under every assumption tested; the only assumption the record breaks is A5; and HY1 shows the component cannot be put back into galaxies under the record's own dust dynamics.

8 What an all-gates ingredient would have to supply

On top of the law, the extra gravitating mass as a fraction f of the baryons (KD1 diagnostic, canonical / alt):

Where	Needed f
SPARC discs	$\leq$ 0.33 / 0.19
X-COP clusters	$\sim$ 4.2 / 3.8
groups	1.2–3.5 / 0.9–2.9
X-ray ellipticals	$\sim$ 0.9 / 0.8 (about half of it IMF)
dSph satellites	$\sim$ 1.7 / 1.3
Coma UDGs (NOT A TEST, UA1)	$\sim$ 80; $\sim$ 12 / 11 on a linear response[a]
the CMB	cold at z 3400 ($w \leq 0.01$), lensing $\chi^2 \leq 18.5$ at $L \leq 400$

[a] The $\sim$ 80 applies the isolated law's sub-linear response at the UDGs' internal acceleration on top of a base that already carries Coma's external field (GB1, Disclosed 3). On a linear response the same step, +1.121 / +1.078 dex, needs $f = 10^{1.121} - 1 = 12.2$ / $10^{1.078} - 1 = 11.0$ (derived here). UA1 leaves the class NOT A TEST, so neither figure is a graded need.

9 Four walls

1. **Cold fluid** (KD1, QX3, QX7). Every well captures it, discs included. The CMB and BBN need $\mu_K \geq 8.9\times10^6\,\mathrm{Mpc}^{-1}$; galaxies allow at most 79.
2. **Warm fixed-temperature sea** (QX12). Retention $\exp(\mathrm{depth}/c_s^2)$ is too sharp; the temperature would have to track each host.
3. **Hot thermal fermion** (HC1, HF1, VW1). Hot enough to stay out of discs, it under-fills clusters, cannot fill dSphs or UDGs, and fails CMB lensing; cold enough for the CMB, discs capture it. No window.

4. **Shock-entropy sector** (SE1). A conserved, non-cooling sector heated by its own shocks either sits isentropic, and holds almost nothing, or needs a different entropy history for each class at the same entropy. Its conserved rest mass also caps a host near the cosmic share, against the ~ 80 the Coma UDGs would need (a class UA1 leaves NOT A TEST).

The common reason: an ingredient that tracks each host's virial state has to virialise with the host; if it is also cold early, nothing keeps it out of discs. And the CMB needs the extra gravity at $z \simeq 3400$, where nothing is virialised, so any key that is on only in today's hot systems is off exactly where the CMB needs it on.

10 The residue: the law's measured boundary

SE2 draws every systematic knob jointly per class (IMF, velocity-dispersion scale, distance, anisotropy, membership, 3-D position, tidal admissibility). The posterior step of the plain law, in dex (median, 95 % interval):

Class	canonical	alt	label
X-COP clusters	+0.440 [+0.372, +0.484]	+0.408 [+0.341, +0.451]	NOT EXPLAINED
groups	+0.232 [+0.078, +0.362]	+0.194 [+0.041, +0.324]	NOT EXPLAINED
Coma UDGs (9)	+1.120 [+0.541, +1.718]	+1.075 [+0.472, +1.673]	NOT EXPLAINED
X-ray ellipticals	+0.067	+0.050	EXPLAINED
KiDS early/late	+0.085	+0.084	EXPLAINED
bright M31 dSphs (21)	+0.228	+0.219	EXPLAINED
CMB-lensing shape	cold extra gravity at z 3400; no key passes the hot classes		SEPARATE

The groups row is the weakest (z 3.04 / 2.55).[1] Gas in the Coma UDGs is not in the budget and would lower their step; hydrostatic bias is set to 0 and would raise the cluster and group steps. These numbers are the boundary of the law as measured, not a mechanism. One profile meets both the inner-core requirement and the gas-to-stars band: a cored isothermal with r_c 50–100 kpc passes NA2's per-bin rule for 7/7 clusters on both footings (r_c 100 kpc: rms 0.086 / 0.086), though no collapse model in the record produces it (NC4, reported). The X-COP step is not a hydrostatic artefact inside $0.5\,R_{500}$: on CLASH lensing masses the law's residual has the same radial shape with lensing above hydrostatic by +0.05 dex in both shared bins, inside the frozen 0.10 band (HL3); the lensing leg does not reach R_{500}.

Update (UA1). UA1 re-graded SE2's two fragile rows. *Coma UDGs: NOT A TEST* on both footings: with tidal admissibility taken at pericentre $d_{\rm mean}/3$ in the Coma field, 2 of 9 are admissible. No orbit or proper motion exists on disk for any of the 9, so the label rests on the $d/3$ convention; with pericentres at $d/2$ or wider, 5 or more would be graded (not computed). NOT A TEST says the class cannot show it is in equilibrium; it does not say systematics explain the step. *Groups: NOT EXPLAINED* under the full single-width budget, +0.259 [+0.052, +0.459] (canonical) and +0.221 [+0.009, +0.415] (alt). The non-thermal term raises the step; distance and clumping lower it by about 0.01 dex each; the cross-catalogue term is 0.064 dex; the Eckmiller group masses are PROVISIONAL. The row is fragile: the alt lower edge is +0.009, and its sign rests on the non-thermal term (the inverted sign drops the median by 0.088 / 0.087 dex). With the same cut the bright M31 dSphs stay EXPLAINED (15 of 25). **The residue is X-COP + groups + CMB lensing**, the group-and-cluster scale, not the cluster scale alone.

Update (XF1). Every hosted row above was graded with a static external-field state. In the action's own moving-body state (XF1, a quarantined test lane) the host's khronon wind sweeps the member's static phantom away with the host's and leaves the member Newtonian: the Coma UDG shortfall grows to +1.412 / +1.409 dex on SE2's path, and X1's dSph rows widen by about 0.9 dex. The static state is, if anything, generous to the theory, and no hosted row moves in its favour. The bright M31 dSphs' EXPLAINED is a label of the static grading; XF1 did not re-run the joint budget under the moving state.

[1] The groups step rests on one hydrostatic sample, the Lovisari+2015 X-ray masses. The dynamical side, from velocity dispersions, does not show it, and the record has not reconciled the sign. HX1's Tian groups need a phantom boost B of median 1.18, $\log B = +0.07\pm0.08$ (alt 1.08, $+0.03\pm0.08$), on the law within errors. HG2's bright-minus-faint offset of KT2017 groups (with versus without an eRASS1 hot atmosphere, against the KiDS blue-lens reference, gas counted) is $D = -0.294$ [−0.511, −0.076] dex, UNDERPOWERED by its frozen rule (power 0.45); GA1's gas audit of the same statistic moves it to −0.362 [−0.605, −0.139] (power 0.36), where all groups on the law would give +0.173. GA1 re-grades HG2's sample; they are not independent.

11 Kill tests, frozen

Three tests can kill the law with data not yet in hand. Each is frozen before the data; nothing else is pending.

1. **Gaia DR4 wide binaries: Newton.** v3s screens binaries, so it predicts Newtonian motion: the frozen pipeline returns γ_∞ 1.0019 / 1.0100 (canonical / alt), not separable from Newton and separable from plain QUMOND (1.1494 / 1.2081). The prediction is sealed (CQ2, SHA-256 1d65509dbf). Scored as the screened (medium-first) reading of the frozen DR4 rows (owner decision 413c1c8e24), v3s is excluded iff $\hat{\gamma} - 1 \geq 3\sigma$, with the gate $\sigma_{\rm sys} < 0.048$ / 0.058 and the R rule (KILL iff $R - 1 \geq \max(0.01, 3\sigma_R)$). At the registered error (N 30,000, $\sigma_{\rm sys}$ 0.020) v3s is EXCLUDED for $\hat{\gamma}$ in 1.083–1.23; at the realistic N of about 8,000 the edge moves up.
2. $a_0(z)$ **flat to** z **2.33.** a_0 tracks $\rho_{\rm DE}$ (CFG689 quantiles, BAO + SN, no ΛCDM input). No sample on disk can decide it: all 52 cells are BLOCKED, because each sample's common mode (0.30–0.60 dex) is 4–11× the largest $e_{3\sigma}$ (0.084 dex at z^* 2.33). It needs a proposal-class ALMA sample with a same-estimator $z < 0.1$ reference (AT2). It kills the law if, at a powered z^*, the rival is favoured or neither model fits (AT1's rule).
3. **Lensing = dynamics.** v3s predicts $M_{\rm lens}/M_{\rm dyn} = 1$ to $\leq 6.5 \times 10^{-5}$ at 310 kpc. KiDS isolated lenses give $A = 1.129 \pm 0.206$ (canonical) and 1.035 ± 0.187 (alt), PASS but with weak power (V3W W3) The kill line is the frozen grade: FAIL if the KiDS ratio sits more than $2\sigma_{\rm tot}$ from 1, or the matched-host ratio more than $3\sigma_{\rm tot}$. At cluster scale, a check on the residue rather than this kill line: the law's residual has the same radial shape on CLASH lensing as on X-COP hydrostatic masses inside $0.5\,R_{500}$ (HL3, SAME SHAPE on both footings).

The exact statistics and kill lines are collected in closure_2026_11_07/PREDICTIONS_2026_11_07.md (52d4f27840), checked by its own script.

12 Record-internal inconsistencies, stated

None of these changes a label; each is stated so that a reader does not take one lane's reading for the record's.

1. **AC2q fold timing.** FS1 and FS2 read the clock-only AC2q folds EARLIER than v3s's: 5.3 % at $J_{\rm thr}$ (FS1) and 1.2 % at $J_{\rm reg}$ (FS2), carried by the stream's edge cells, which shock instead of crossing. AC10 puts the two folds at the same time to 0.3 % (1.0752–1.0782 against 1.0718–1.0773 τ_f), and SV3 finds that AC2q's extra s^2 pressure delays clock-only folds by about 0.1–0.3 %. The sign depends on the integrator and on how the edge is treated; no lane finds a regularization.
2. **PPN** α_1, α_2 **for a moving Sun.** AS3 finds $\alpha_1 = \alpha_2 = 0$ exactly for a source moving through the aether, on the linear system. AC11 lists the metric response that would give α_1 and α_2 as still open, and MS3's scoped PASS states that no PPN α_1 or α_2 is claimed. The linear result stands; the nonlinear moving-Sun field has not delivered it.
3. $a_0(z)$ **at** z **2.33.** CFG689's native band is 0.91 [0.33, 1.21]. X35's flat-Λ estimate with an early-dark-energy ruler moves the native $\sqrt{X}$ from 1.00 to 1.49, and CT3's retired-sector route gives $a_0(2.33)/a_0(0) \approx 0.85$. The kill test needs 0.084 dex at z^* 2.33 for 3σ (CFG689) and is set at 0.10 dex by CT1, so the predicted history must be named before the data, with its tolerance: the readings in the record differ by more than the test resolves.
4. **Units in CK3.** L388's masses are in $M_\odot/h$, and XR13's placement treats them as $M_\odot$; CK3 inherits the slip and is FAIL as frozen, NOT ROBUST. Its pre-registered unit-consistent reading ($M \times 0.6736$) reads PASS in every cell, but from prediction error, not agreement. CK3 reports the record-level fix and does not make it; CK1 places its systems consistently.
5. **V3X.** V3X is UNDECIDED under its frozen rule on W2 and W4c, with a physics reading of PASS. AZ31 passes W2 at the graded radius as a new test, leaving V3X's label intact, and AZ32 keeps the full W4c UNDECIDED.
6. **Static versus moving external field.** AC15 and MS3 found that a moving Sun sweeps its static hedgehog away, while every hosted row was graded with a static external-field state. XF1 now grades both under one state, the action's own moving-body state: the Sun stays under every AC15/MS3 line, and the hosted rows get worse, not better. Isolated discs are still graded with a static khronon (Section 13).
7. **Spin as a selector.** A pre-read of 2026-11-09 listed rotation as the one selector with no verdict. KU2 (rebuild_2026_10_22) had already graded slow against fast rotators UNIVERSAL on the same MaNGA DynPop file at matched $(M_\star, \Sigma_{R_e})$; SK1 confirms it on the law's residual at matched $(M_\star, g_N)$. ST1's X-ray elliptical IMF offset comes from the Humphrey 2006 table, not ATLAS3D; SK1 replays it on that table.

13 Reading

Every physics class on the record now carries a label, and none passes every gate. The law, and the action within its scope, describe cold rotating discs and the Solar System. One limit on the disc rows is stated, not graded: the sweep XF1 finds for hosted members would also act on an isolated galaxy moving through the cosmic khronon at its peculiar speed, where the record grades discs with a static khronon; the galaxy's own charge slows the inflowing frame only marginally at Milky Way mass, and not at all for dwarfs (XF1). The ellipticals, KiDS and the bright M31 dwarfs are accounted for by measured systematics. X-COP and the groups are not; the Coma UDGs are NOT A TEST on the record's orbit convention; and the CMB-lensing wall stands on its own. The elimination argument (closure_2026_11_07/DEDUCTION_2026_11_07.md) runs as follows. Three facts are measured: something cold gravitates at recombination; galaxies hold almost none of it today; clusters hold it today. No time-independent ingredient carries all three, and every switch on the law is closed. A component whose state changes in time leaves two cells. The escape cell, a late kick above the galaxy escape speed, is closed by X-COP. The puff cell, a late heating that leaves a diffuse halo of about $4\,M_b$ around galaxies, is closed by KiDS once the hot gas is measured rather than assumed (KX1). Its evaporated-core form, C3, fails X-COP, the groups and the Bullet (CE2), so every form of the puff cell is closed. The escape cell, re-graded at today's needs, fails on every kick from 500 to 650 km/s (CK1), and the action supplies no kick, neither in its statics (CK2) nor in its fold dynamics, where the release speed tracks the well and not the identity's 577.6 km/s (AK1). The time-dependent cells die on the lenses (KX1), on X-COP, the groups and the Bullet (CE2) and on escape speeds (CK1), not on growth: CE2's growth gate passes every late-heated cell (Rf = 1.000 at every $z < 1$ bin). A component born clumped at the nodes fails the CMB's bias tolerance and the forest (NC2), and cluster-member galaxies are as clean as field galaxies (NC3), so the exhaustion covers initial conditions as well as mechanisms (DEDUCTION section 8). One structural fact frames all of this: every host inside R_{500} together holds about 1 % of the cold component the CMB needs (0.0091 / 0.0078; at most 0.025 / 0.022 in the most generous bracket), so nearly all of it, whatever it is, lies outside bound objects (NC4). With F1 now audited by independent code (F1A), all three measured facts rest on independent checks (DEDUCTION section 9); the action deposited here is the full action the record can write, and its one unresolved failure is a contradiction rather than a gap: its own cold dust, forced by the CMB and present in clusters, falls into galaxies where no measurement finds it, and no mechanism or initial condition in reach of the action prevents that. The action's own dust term also cannot form the cored cluster profile (KQ1; DEDUCTION section 10), so the residue's shape is a target no term on the table reproduces. On 2026-11-07 the owner accepted the corrected-kernel labels as standing (Section 6); none of them opens a door, DEDUCTION sections 7–10 are unchanged, and no row in the record is pending. The elimination is complete for every ingredient the record can build. The a_0 law with the adopted v3s action is the law of cold isolated gravity; the CMB's cold component, the cluster and group residue, and the absence of either from galaxies are three measured facts that no single ingredient in reach of the action connects. Which of the three gives way is not decidable on disk: the galaxy leg reads as the firmest (KX1), though that is a reading: KX1's frozen label is NO VERDICT, and its canonical C4 hold rests on the lightest KiDS bin (log $M_\star$ 9.4, 122–285 kpc), which uses a frozen extrapolation of the log $M_\star$ 10.0 kSZ gas row; without that bin canonical C4 would read SOFTENS; the cluster leg rests on seven X-COP clusters and provisional group masses (UA1); the CMB leg is a fit under the footing background, not independent evidence (QX15). The hybrid route, the law plus a component that galaxies hold, is now graded under the record's own dust dynamics (HY1, a quarantined test lane) and fails on both sides at once: the dust over-fills discs and cluster cores together. The owner's lift of the no-cold-component rule covered HY1 only. KR1 exhausts the $\mathcal{K}(Q)$ sector at the level of its functional form, not only its amplitude: for any convex $\mathcal{K}$ the kernel mass is a function of the sector's own local density, so the growth line the cosmology needs fixes the mass of the dust in every galaxy, and that dust is captured. With KR2 the $\mathcal{K}(Q)$ door is closed from both ends, the early universe (KR1) and the present day (KR2), at the level of the sector's functional form. BG1, a quarantined test lane, tried the one key that separates the regimes by the state of the baryons rather than by any gravitational variable, conversion of the dust where baryons are cold and dense; it also fails, because the densest cold baryons in the history of the universe are the neutral gas of the dark ages, and a rate that clears discs clears the CMB's component before z 20. PJ1, an outside proposal for a potential-gated Jeans scale, dies on the same rungs: a gate weak enough to spare the acoustic peaks leaves plain dust in every well, and any gate strong enough to shape the component cuts it across the acoustic range. WD1, an outside proposal for a merger-decohered component with a 1.6 Mpc core, dies on the same rungs from the other side: a core wide enough to spare the isolated lenses inside 300 kpc overfills them beyond 1 Mpc and erases the forest, and the merger-lowered core overfills cluster centres while the satellite rule leaves the Coma UDGs with nothing. This is the best remaining answer the record supports, not a closed theory.

14 Closing statement

No label changes here; the statement collects what the rows above already say.

1. **Domain.** The a_0 law, with $\kappa = \frac{1}{2}$ fitted, describes rotation curves, the lensing of isolated galaxies and the Solar System, and the screened khronon action carries it covariantly, with the Solar System and moving-Sun tests passed (MS3, scoped). The disc rows carry the limit XF1 states (Section 13).
2. **Boundary, as a statement about the record.** Hot systems, the CMB and the linear web need a clustering component the law does not supply. Every selector graded that would key that component, or the law's own boost, on the baryons' present state (acceleration, potential depth, density, temperature, size, environment, baryon phase, spin and dispersion: the switches of DEDUCTION section 1, with HX1, BG1, KU2 and SK1) or on their past (FA3; the clocks of KX1, CE2 and CK1) has a verdict, and none passes. The failures recur on the same rungs: the isolated outer lenses and the forest, the cluster cores, and the hosted dwarfs. Every action class graded (AX1, EW1–EW3, FA2) and every sector graded (the cold dust, the $\mathcal{K}(Q)$ dust, the hot fermion, PBHs, QX27–QX31, PJ1, WD1) fails as well, and re-grading the hosted rows in the action's own moving state (XF1) makes them worse. Within the catalog the record has graded, no completion of the law by a baryon-keyed component exists. A completion would need a component carrying its own initial conditions and dynamics, not keyed to the baryons. This programme does not adopt one, and the one such form graded, born clumped at the nodes, fails (NC2).
3. **What such a component must be, from the record's own numbers.** Its cosmic amount is about 5.4 times the baryons (ω_d/ω_b 5.410 / 5.412), cold and clustering at recombination (F1) and still cold through the forest epoch at z 2–4 (DEDUCTION section 1). Galaxies hold at most 0.33 / 0.19 of the baryons' mass of it inside their rotation curves, and isolated lenses about $0.9\,M_b$ inside 200 kpc (F2). In clusters it is a cluster-universal fraction of the dynamical mass that falls outward, about 70 % inside $0.25\,R_{500}$ to about 45 % at R_{500} on TR1's reading D, between the stars and the gas in concentration and not a single multiple of any baryon tracer (TR1), with the same per-bin shape on the lensing and hydrostatic legs inside $0.5\,R_{500}$ (HL3). Bound hosts inside R_{500} hold about 1 % of the cosmic amount (NC4).

The law can still be decided in its own domain by the three frozen kill tests: Gaia DR4 wide binaries (expected 2026-12-02), a flat $a_0(z)$ to z 2.33, and lensing equal to dynamics on isolated lenses. Adopts nothing. $\kappa = \frac{1}{2}$ is fitted; the theory is not closed.

15 What is unified, and what is not

From closure_2026_11_07/UNIFICATION_STATEMENT_2026_11_07.md, section 3, verbatim:

> Gravity carries one extra scale, the acceleration of the dark energy, a0 = (1/2) c sqrt(G rho_DE), and one action, the screened khronon action with its adopted stiffness, realises it as a complete and certified law of cold isolated gravity from the Solar System to galaxy lensing and gravitational waves. The same relation predicts a constant a0 to z 2.33 and Newtonian wide binaries, which are its kill tests. Beyond the cold isolated regime the universe contains a cold gravitating component that the CMB requires, clusters hold and galaxies lack, and no term in reach of the action, nor any ingredient the record could build, unifies that component with the law; the one candidate that does so adds a postulate under test today. That is the whole of the unification the record supports: one scale, one action, one measured boundary, and a named missing identity.

Since that statement was written, the candidate it names has been graded: CK1 FAILS it on every kick, and CK2 finds no term in the action that releases one. The kernel erratum (Section 6) leaves wall 1 and every web row standing. $\kappa = \frac{1}{2}$ stays fitted and not derivable. VR1 tested the vacuum-response identity a principle would have to supply and finds none: the missing premise is that the vacuum's response length at the switch is $c/\sqrt{G\rho_{\rm DE}}$, and nothing in reach supplies a π-free length. SA1 reduces a settling-age dependence of the action's dust to closed cells. CMB1 and CMB2 (Section 7) leave the cold component's requirement as measured and its carrier outside the action.

16 What waits on the owner

The PAPER68 erratum for the Legendre form. The Π^6 stiffness repair, whose case for and against was written for the owner (PX1, closure_2026_11_07/PI6_BLEND_ADOPTION_CASE_2026_11_07.md), was adopted by the owner on 2026-11-07 (closure_2026_11_07/OWNER_DECISION_2026-11-07_ADOPT_PI6_BLEND.md). Every owner-opened test lane above has landed with its label, and none is still running; none adopts anything.

Method

Every lane froze its criteria in a commit before any script ran, and every lane has a MUTATE twin that must fail. Footings are reported apart. No ΛCDM inputs were used. Scripts and outputs are in the repository under closure_2026_11_06/ to closure_2026_11_09/ (and the earlier lane folders the rows name) at the commits the lane RESULT files name. The supplement archive deposited with this note holds every Lean source and build output, the closure tables, and each lane's criteria and result files, at the commit recorded as git_head in PAPER70_supplement_manifest.json, together with this note's source and audit script.