


MACQUARIE UNIVERSITY

Grokking MATH135

A very unofficial textbook.



SID	Name	Initials	Email
4196 3539	CARMICHAEL, Adam Justin	AJC	adam.carmichael@ieee.org
xxxx xxxx	HUNT, Nathaniel K	NHK	
yyyy yyyy	RACE, Alison	AR	

Document compiled Monday 17th December, 2012 at 00:05

CHALLENGE
ACCEPTED

CONTENTS

Contents	1
1 Trigonometry Review	5
1.1 Radian & Arc Length	5
1.2 The Unit Circle	6
1.3 Trigonometric Functions	6
1.4 Pythagorean Identities	7
1.5 Properties of Trigonometric Functions	8
1.6 Trigonometric Identities	9
2 Polynomials	11
2.1 Polynomial Equations	11
2.2 Polynomial Division	12
2.3 Polynomial Roots	14
2.4 Fundamental Theorem of Algebra	14
2.5 Roots of Real Polynomials	15
2.6 More Polynomial Roots	15
2.7 Rational Functions	15
2.8 Greatest Common Divisor	15
3 Complex Numbers	17
3.1 Arithmetic of Complex Numbers	19
3.2 Complex Conjugates	19
4 Complex Plane	21
4.1 Argand Diagrams	21
4.2 Polar Coordinates	23
4.3 Modulus	23
4.4 Euler's Formula	24
5 de Moivre's Theorem & Roots of Complex Numbers	25
5.1 de Moivre's Theorem	25
5.2 Finding Roots of Complex Numbers	26
5.3 Analytic Geometry	26
6 Linear Equations	27
6.1 System of Linear Equations as Matrices	27
6.2 Elementary Row Operations	27
6.3 Row Echelon Form	27

6.4	Reduced Row Echelon Form	27
6.5	Solving a System of Linear Equations	27
6.6	Homogeneous Systems	28
7	Matrices & Linear Equations	29
8	Application of Linear Equations	31
8.1	Application to Network Flow	31
8.2	Application to Electric Circuits	31
8.3	Application to Economics	31
8.4	Application to Chemistry	31
8.5	Application to Mechanics	31
9	Arithmetic of Matrices	33
10	Determinant of a Matrix	35
11	Vectors	37
12	Vectors & Geometry	39
13	Combinatorics & Binomial Theorem	41
14	Real Numbers	43
15	Induction	45
16	Functions	47
17	Limits	49
18	Continuity	51
19	Differentiability	53
20	Stationary Points	55
21	Curve Sketching	57
22	Applications of the Derivative	59
23	Antiderivatives	61
24	Techniques of Integration	63
25	Separable Differentiable Equations	65
26	Area Under a Curve	67
27	Volume of Solids of Revolution	69
28	More Applications of Integration	71

CONTENTS

3

29 Glossary

73

TRIGONOMETRY REVIEW

Further reading: Chen & Duong ‘‘mtfym01.pdf’’, <http://goo.gl/wyojl>

Big Ideas:

- Radians and arc length are interrelated. There are 2π radians in a circle.
- An angle subtended by θ will have arc length θ .
- $\cos(\theta) = y\text{-ordinate}$, $\sin(\theta) = x\text{-ordinate}$
- $\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)} = \frac{y}{x}$
- **Pythagorean Identities** (page 7) are worth learning!

1.1 Radian & Arc Length

Further reading: Chen & Duong ‘‘mtfym01.pdf’’ pp 1-2, <http://goo.gl/wyojl>

Radians and arc length are interrelated. There are 2π radians in a circle, that is a semi-circle has about 3.14 radians (π radians to be exact).

In areas of maths, science and engineering radians are used as the main unit of measurement of angles rather than degrees, as such it can be handy to know how to convert between the two units.

$$\text{degrees} = \text{radians} \times \frac{180}{\pi} \quad (1.1)$$

$$\text{radians} = \text{degrees} \times \frac{\pi}{180} \quad (1.2)$$

Aside from the conversion formula, it is also useful to remember the following special values:

$$\frac{\pi}{6} = 30^\circ \quad (1.3)$$

$$\frac{\pi}{4} = 45^\circ \quad (1.4)$$

$$\frac{\pi}{3} = 60^\circ \quad (1.5)$$

$$\frac{\pi}{2} = 90^\circ \quad (1.6)$$

$$\pi = 180^\circ \quad (1.7)$$

$$2\pi = 360^\circ \quad (1.8)$$

1.2 The Unit Circle

In high school we learned that trigonometry was about angles of triangles, and while this remains true, we look at it in the context of circles. Consider a special circle called *the unit circle*. See (ref fig 1.2 page 7). If we run a vertical line from each point in the unit circle to the line $y = 0$, the triangle is formed.

1.3 Trigonometric Functions

Further reading: Chen & Duong “mtfym01.pdf” pp 2-4, <http://goo.gl/wyobjl>

Looking at the unit circle, an angle is defined by the two points, the first at the origin (0,0), the second is $(\cos\theta, \sin\theta)$. That is to say

$$x = \cos\theta \quad \text{and} \quad y = \sin\theta \quad (1.9)$$

Additionally, we can define $\tan\theta$:

$$\tan\theta = \frac{\sin\theta}{\cos\theta} = \frac{y}{x} \quad (1.10)$$

Finally there are the inverse trigonometric functions:

$$\sec\theta = \frac{1}{\cos\theta} = \frac{1}{x} \quad \text{and} \quad \csc\theta = \frac{1}{\sin\theta} = \frac{1}{y} \quad (1.11)$$

$$\cot\theta = \frac{\cos\theta}{\sin\theta} = \frac{x}{y} \quad (1.12)$$

Note: $\tan\theta$ and $\sec\theta$ are defined only when $\cos\theta \neq 0$ and $\cot\theta$ and $\csc\theta$ are defined only when $\sin\theta \neq 0$.




Figure 1.1: The Unit Circle, courtesy of Supreme Ayal, TiKZ Examples [?]]

1.4 Pythagorean Identities

Further reading: Chen & Duong ‘mtfym01.pdf’ pp 4-5, <http://goo.gl/wyoj1>

$$\cos^2 \theta + \sin^2 \theta = 1 \quad (1.13)$$

By dividing 1.13 by $\cos^2 \theta$ we get:

$$1 + \tan^2 \theta = \sec^2 \theta \quad (1.14)$$

By dividing 1.13 by $\sin^2 \theta$ we get:

$$1 + \cot^2 \theta = \csc^2 \theta \quad (1.15)$$

1.5 Properties of Trigonometric Functions

Further reading: Chen & Duong ‘mtfym01.pdf’ pp 5-10, <http://goo.gl/wyojl>

Properties of $\sin \theta$

$$\cos(\theta + 2\pi) = \cos \theta \quad (1.16)$$

$$\cos(-\theta) = -\cos \theta \quad (1.17)$$

$$\cos(\theta + \pi) = -\cos \theta \quad (1.18)$$

$$\cos(\pi - \theta) = \cos \theta \quad (1.19)$$




Figure 1.2: $f(x) = \cos \theta$

Properties of $\cos \theta$

$$\sin(\theta + 2\pi) = \sin \theta \quad (1.20)$$

$$\sin(-\theta) = -\sin \theta \quad (1.21)$$

$$\sin(\theta + \pi) = -\sin \theta \quad (1.22)$$



$$\sin(\pi - \theta) = \sin \theta \quad (1.23)$$

Properties of $\tan \theta$

$$\tan(\theta + \pi) = \tan \theta \quad (1.24)$$

$$\tan(-\theta) = -\tan \theta \quad (1.25)$$

$$\tan(\pi - \theta) = -\tan \theta \quad (1.26)$$

Figure 1.3: $f(x) = \sin \theta$ Figure 1.4: $f(x) = \tan \theta$

1.6 Trigonometric Identities

Further reading: Chen & Duong ‘mtfym01.pdf’ pp 11, <http://goo.gl/wy0jl>

Sine Rule

$$\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)} \quad (1.27)$$

Cosine Rule

$$a^2 = b^2 + c^2 - 2bc \cos(A) \quad (1.28)$$

$$b^2 = a^2 + c^2 - 2ac \cos(B) \quad (1.29)$$

$$c^2 = a^2 + b^2 - 2ab \cos(C) \quad (1.30)$$

Angle Sum & Difference Identities

Further reading: Chen & Duong “mtfym01.pdf” pp 12, <http://goo.gl/wy0jl>

$$\sin(A \pm B) = \sin(A) \cos(B) \pm \cos(A) \sin(B) \quad (1.31)$$

$$\cos(A \pm B) = \cos(A) \cos(B) \mp \sin(A) \sin(B) \quad (1.32)$$

Double Angle Identity

Further reading: Chen & Duong “mtfym01.pdf” pp 12, <http://goo.gl/wy0jl>

$$\sin(2\theta) = 2 \sin(\theta) \cos(\theta) \quad (1.33)$$

$$\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta) \quad (1.34)$$

$$= 1 - 2 \sin^2(\theta) \quad (1.35)$$

$$= 2 \cos^2(\theta) - 1 \quad (1.36)$$

Half Angle Identity

Further reading: Chen & Duong “mtfym01.pdf” pp 13-15, <http://goo.gl/wy0jl>

$$\sin^2\left(\frac{\theta}{2}\right) = \frac{1 - \cos(\theta)}{2} \quad (1.37)$$

$$\cos^2\left(\frac{\theta}{2}\right) = \frac{1 + \cos(\theta)}{2} \quad (1.38)$$

CHAPTER 2

POLYNOMIALS

Further reading: Chen & Duong “mtfym02.pdf”, <http://goo.gl/jHCXo>

Big Ideas:

- An expression of finite length constructed from variables and constants using only addition, subtraction, multiplication and non-negative integer powers.
- The highest power of a variable indicates the polynomials *degree*.
- Determining the *roots* of a polynomial tells us where a polynomial equation intersects the x -axis.
- Polynomials of degree 2 are called *quadratic*.
- Polynomials of degree 3 are called *cubic*.

2.1 Polynomial Equations

Further reading: Chen & Duong “mtfym02.pdf” pp 3-4.

An equation in the form $ax + b = 0$ where $a, b \in \mathbb{F}$ and $a \neq 0$ is called a *linear equation* or *linear polynomial equation*. It has a degree of 1 (that is, the power of x is 1). We can determine the value of x :

$$ax + b = 0 \quad (2.1)$$

$$ax = -b \quad (2.2)$$

$$x = -\frac{b}{a} \quad (2.3)$$

An equation in the form $ax^2 + bx + c = 0$ where $a, b, c \in \mathbb{F}$ are constants and $a \neq 0$ is called a *quadratic equation*. To solve such an equation we use the *quadratic formula*:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad (2.4)$$

Where $b^2 - 4ac$ (called the *determinant*) is greater than or equal to 0. This will yield two solutions:

$$x = \frac{-b + \sqrt{b^2 - 4ac}}{2a}$$

and

$$x = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$$

However, equation 2.4 is the most compact and easiest to remember.

If the determinant < 0 , then it indicates there are no roots exist in the real plane (\mathbb{R}) and that a complex plane (\mathbb{C}) solution exists.

For polynomials of degree 3 or more there is no general formula for a solution, however division of polynomials allows us to reduce polynomials of a higher degree to a factor and a lesser degree. By iteratively reducing polynomials we can eventually find all roots.

2.2 Polynomial Division

Further reading: Chen & Duong “mtfym02.pdf” pp 4-6.

Remember long division from primary school? By the time we get to university most of us have forgotten. Checkout <http://goo.gl/TvDoI> for a Kahn Academy video on how to do it using just plain numbers. It is well worth the refresher even if you think you can remember how it works.

There is a simple recursive algorithm for polynomial long division, consisting of 3 steps which are repeated until either a divide by zero, or a remainder occurs.¹

Jaye's Algorithm:

1. Divide the next term numerator (the part under the long line) by the first term in the denominator. Write the result above the long line.
2. Multiply the result by the denominator and then by -1 . Write the result underneath the term(s) operated on, underlining any new terms.
3. Perform a simple addition, carrying anything down below the line.

Rinse and repeat these 3 steps underneath your new long line.

The following example comes from Purple Math, <http://goo.gl/GPi0z>: Suppose we wish to divide $\frac{x^2 - 9x - 10}{x + 1}$:

1. First write the polynomial in the form:

$$\begin{array}{r} x + 1 \overline{) x^2 - 9x - 10} \end{array}$$

We will look only at the leading x in the divisor and the leading x^2 in the numerator.

¹I call this algorithm Jaye's Algorithm because Jaye, a 9 year old at the time, helped me memorise it a few weeks before writing it down.

2. We want to divide the x^2 in the numerator by the x in the denominator. $\frac{x^2}{x} = x$ so we write the resulting x above the line.

$$\begin{array}{r} x \\ x+1 \overline{) x^2 - 9x - 10} \end{array}$$

3. Now we must take care of the $+1$ in the denominator. Multiply the result above the line by -1 and multiply again by the denominator:

$$\begin{array}{r} x \\ x+1 \overline{) x^2 - 9x - 10} \\ \underline{-x^2 \quad -x} \end{array}$$

4. Perform the subtraction as we would a normal sum carrying anything to the right down:

$$\begin{array}{r} x \\ x+1 \overline{) x^2 - 9x - 10} \\ \underline{-x^2 \quad -x} \\ -10x - 10 \end{array}$$

5. Divide $-10x - 10$ by x goes -10 times, so write -10 above the line

$$\begin{array}{r} x-10 \\ x+1 \overline{) x^2 - 9x - 10} \\ \underline{-x^2 \quad -x} \\ -10x - 10 \end{array}$$

6. Multiply -10 by -1 and then again by $(x+1)$:

$$\begin{array}{r} x-10 \\ x+1 \overline{) x^2 - 9x - 10} \\ \underline{-x^2 \quad -x} \\ -10x - 10 \\ 10x + 10 \\ \hline 0 \end{array}$$

7. **The zero means there is no remainder, and we are done.**

Another example of polynomial long division, taken from the same site as above, to divide $\frac{x^2+9x+14}{x+7}$

1. Write out in the standard form:

$$\begin{array}{r} x+7 \overline{) x^2 + 9x + 14} \end{array}$$

2. Next divide $\frac{x^2}{x} = x$ and write above the line.

$$\begin{array}{r} x \\ x+7 \overline{) x^2 + 9x + 14} \end{array}$$

3. Multiply $-x$ by $x+7$ and write underneath:

$$\begin{array}{r} x \\ x+7 \overline{) x^2 + 9x + 14} \\ \underline{-x^2 - 7x} \end{array}$$

4. Perform the addition:

$$\begin{array}{r}
 x \\
 x+7 \overline{) x^2+9x+14} \\
 \underline{-x^2-7x} \\
 2x+14
 \end{array}$$

5. Divide
- $\frac{2x}{x} = 2$

$$\begin{array}{r}
 x+2 \\
 x+7 \overline{) x^2+9x+14} \\
 \underline{-x^2-7x} \\
 2x+14
 \end{array}$$

6. Multiply
- -2
- by
- $x+7$
- and write underneath, and we can see there is no remainder.

$$\begin{array}{r}
 x+2 \\
 x+7 \overline{) x^2+9x+14} \\
 \underline{-x^2-7x} \\
 2x+14 \\
 \underline{-2x-14} \\
 0
 \end{array}$$

Sometimes there are tricky ones where you might have some terms with a coefficient of zero, such as $\frac{2x^3-9x^2+15}{2x-5}$. Here x^1 does not appear as a part of numerator, but we can rewrite it as $\frac{2x^3-9x^2+0x+15}{2x-5}$. With problems such as these, it is easiest to keep your writing neat and in columns. Where you might see $0x$ you can just as easily substitute a bit of whitespace.

$$\begin{array}{r}
 x^2-2x-5 \\
 2x-5 \overline{) 2x^3-9x^2 +15} \\
 \underline{-2x^3+5x^2} \\
 -4x^2 \\
 \underline{4x^2-10x} \\
 -10x+15 \\
 \underline{10x-25} \\
 -10
 \end{array}$$

This example also shows a remainder of -10 . The final answer could also

be written as one of two ways, either:

1. $x^2-2x-5-\left(\frac{10}{2x-5}\right)$ – or –
2. $x^2-2x-5 \text{ r } 10$

Though the first method is probably better as it is less confusing sans the letter “r”.

2.3 Polynomial Roots

Further reading: Chen & Duong “mtfym02.pdf” pp 6–8.

2.4 Fundamental Theorem of Algebra

Further reading: Chen & Duong “mtfym02.pdf” pp 7.

2.5 Roots of Real Polynomials

Further reading: Chen & Duong “mtfym02.pdf” pp 7-8.

2.6 More Polynomial Roots

Further reading: Chen & Duong “mtfym02.pdf” pp 8.

2.7 Rational Functions

Further reading: Chen & Duong “mtfym02.pdf” pp 8-11.

2.8 Greatest Common Divisor

Further reading: Chen & Duong “mtfym02.pdf” pp 12-13.

Further reading: Kenneth H ROSEN “Discrete Mathematics and Its Applications 6e”, Chapter Primes and Greatest Common Divisors, pp 210-227.

Definition: Let a and b be integers not both zero. The largest integer d such that $d|a$ and $d|b$ is called the greatest common divisor of a and b , denoted as $\gcd(a,b)$.

^aread as d divides a

There are many ways to determine the gcd of two numbers, including a products of primes and Euclid's algorithm. Euclid's algorithm is by far the most efficient and is discussed in the next subsection.

Euclid's Algorithm

Further reading: Chen & Duong “mtfym02.pdf” pp 12-13.

Further reading: Rosen “Discrete Mathematics and Its Applications”, 6e, pp 227-229

Euclid's algorithm is recursive, and consists of 2 steps:

Euclid's algorithm:

1. Divide the larger of the two numbers by the smaller of the two numbers, and add the remainder onto the end.
2. Find the gcd of the remainder and the smaller of the two original numbers.

In summary:

$$\text{Let } a = bq + r \quad | \quad a, b, q, r \in \mathbb{Z} \quad \text{Then } \gcd(a, b) = \gcd(b, r) \quad (2.5)$$

The following example comes from ROSEN p 229:

$$662 = 414 \cdot 1 + 248 \quad (2.6)$$

$$414 = 248 \cdot 1 + 166 \quad (2.7)$$

$$248 = 166 \cdot 1 + 82 \quad (2.8)$$

$$166 = 82 \cdot 2 + 2 \quad (2.9)$$

$$82 = 2 \cdot 41 \quad (2.10)$$

2 is the last non-zero remainder, therefore $\gcd(414, 662) = 2$.

COMPLEX NUMBERS

Further reading: Chen & Duong “em09-cn.pdf”, <http://goo.gl/C5rfq>

Further reading: Chen & Duong “fyc01.pdf”, <http://goo.gl/e0E7y>

Further reading: Chris Cooper “LINALG03 Complex Numbers”, <http://goo.gl/N85u0>

Big Ideas:

- Complex numbers are created to solve what would otherwise be unsolvable equations because we cannot square any number in \mathbb{R} to get a negative number.
- Complex numbers are made of two parts, a real part and an imaginary part (i).
- i exhibits ring-like behaviour when raised to powers.
- A complex number resembles the form:

$$z = x + yi \quad | \quad x, y \in \mathbb{R} \quad (3.1)$$

- Rules exist for adding, subtracting, multiplying and dividing complex numbers.
- Complex numbers fall into their own special set of numbers \mathbb{C} .
- Complex numbers have a “twin” called a *complex conjugate*, suppose we have:

$$z = (x + yi)$$

then the conjugate is:

$$\bar{z} = (x - yi)$$

When we square any number in the real number plane, we *always* get a positive number. This begs the question, how do we square root a negative number? Previously, a problem such as

$$i^2 + 1 = 0 \quad (3.2)$$

$$i^2 = -1 \quad (3.3)$$

Would be left with “no real solution” as an answer in a paper. While this is true, we **can in fact find a solution**. It just happens to be outside of the real plane and is defined exactly as:

$$i = \sqrt{-1} \quad (3.4)$$

This gives rise to a very interesting property... What if we were to square i ? We already know, the answer from equation 3.3

$$i^2 = -1 \quad (3.5)$$

What about if we cube i ?

$$i^3 = i * i^2 \quad (3.6)$$

$$= i * -1 \quad (3.7)$$

$$= -i \quad (3.8)$$

What if we go again?

$$i^4 = i^2 * i^2 \quad (3.9)$$

$$= -1 * -1 \quad (3.10)$$

$$= 1 \quad (3.11)$$

And again?

$$i^5 = i * i^4 \quad (3.12)$$

$$= i * 1 \quad (3.13)$$

$$= i \quad (3.14)$$

This repetitive behaviour is called a *ring* and is cyclic for every 4th power of i . Remember this fact, it will become useful in later trigonometry.

Ring behaviour of powers of i :

$$i^0 = 1 \quad (3.15)$$

$$i^1 = i \quad (3.16)$$

$$i^2 = -1 \quad (3.17)$$

$$i^3 = -i \quad (3.18)$$

$$i^4 = 1 \quad (3.19)$$

$$i^5 = i \quad (3.20)$$

$$i^6 = -1 \quad (3.21)$$

$$i^7 = -i \quad (3.22)$$

$$i^8 = 1 \quad (3.23)$$

Why does this work so neatly? Consider our basic index laws and apply to numbers between 0 and 3 inclusive:

$$i^{a+b} = i^a * i^b \quad | \quad a, b \in [0, 3] \quad (3.24)$$

3.1 Arithmetic of Complex Numbers

Further reading: Chen & Duong “em09-cn.pdf” pp 1-2.

Further reading: Chen & Duong “fyc01.pdf” pp 6-8.

Addition Rule:

$$(a + bi) + (c + di) = (a + c) + (b + d)i \quad (3.25)$$

Subtraction Rule:

$$(a + bi) - (c + di) = (a - c) + (b - d)i \quad (3.26)$$

Multiplication Rule:

$$(a + bi) \cdot (c + di) = (ac - bd) + (ad + bc)i \quad (3.27)$$

Division Rule:

$$\frac{(a + bi)}{(c + di)} = x + yi \quad | \quad x, y \in \mathbb{R} \quad (3.28)$$

such that

$$a + bi = (c + di)(x + yi) = (cx - dy) + (cy + dx)i \quad (3.29)$$

It follows that

$$a = cx - dy \quad (3.30)$$

$$b = cy + dx \quad (3.31)$$

With the solution:

$$x = \frac{ac + bd}{c^2 + d^2} \quad \text{and} \quad y = \frac{bc - ad}{c^2 + d^2} \quad (3.32)$$

3.2 Complex Conjugates

Complex numbers have a “twin” called a *complex conjugate*, suppose we have:

$$z = (x + yi)$$

then the conjugate is:

$$\bar{z} = (x - yi)$$

Conjugates exhibit special properties, they are:

$$\overline{z + w} = \bar{z} + \bar{w} \quad (3.33)$$

and

$$\overline{z \cdot w} = \bar{z} \cdot \bar{w} \quad (3.34)$$

Proof of equation 3.33 is as follows:

$$\text{Let: } z = x + yi \quad | \quad x, y \in \mathbb{R} \quad (3.35)$$

$$\text{Let: } w = u + vi \quad | \quad u, v \in \mathbb{R} \quad (3.36)$$

$$\overline{z + w} = \overline{(x + u) + (y + v)i} \quad (3.37)$$

$$= (x + u) - (y + v)i \quad (3.38)$$

$$= (x - yi) + (u - vi) \quad (3.39)$$

$$= \bar{z} + \bar{w} \quad (3.40)$$

□

Proof of equation 3.34 is as follows:

$$\text{Let: } z = x + yi \quad | \quad x, y \in \mathbb{R} \quad (3.41)$$

$$\text{Let: } w = u + vi \quad | \quad u, v \in \mathbb{R} \quad (3.42)$$

$$\overline{z \cdot w} = \overline{(x + yi)(u + vi)} \quad (3.43)$$

$$= \overline{(xu - yv) + (xv + yu)i} \quad (3.44)$$

$$= (xu - yv) - (xv + yu)i \quad (3.45)$$

$$= (x - yi)(u - vi) \quad (3.46)$$

$$= \bar{z} \cdot \bar{w} \quad (3.47)$$

□

COMPLEX PLANE

Further reading: Chen & Duong “em09-cn.pdf” pp 3-4.

Further reading: Chen & Duong “fyc01.pdf” pp 8-11.

Big Ideas:

- Complex numbers take the form $(a + bi)$.
- We can represent x and y ordinates in the real plane (\mathbb{R}^2) in the form $(x + yi)$.
- x is called the real part (represented by $\Re[z]$)
- y is called the imaginary part (represented by $\Im[z]$)
- Plotting these on an x, y axis is called an *Argand diagram*.
- Polar coordinates use two variables, (r, θ) to describe a point.
- θ is an angle measured in radians
- r is a length.
- Conversion between polar coordinates and Cartesian coordinates is done using Pythagoras theorem or trigonometry.

4.1 Argand Diagrams

Complex numbers consist of two parts, such as $z = (x + yi)$. Suppose we were to plot both x and y ordinates on an x, y plane as seen in figure 4.1.

Following our addition rules in section 3.1 (page 19), we can add, two complex numbers as seen in figure 4.2.



Figure 4.1: Argand Diagram, z , and complex conjugate \bar{z} .

Figure 4.2: Addition of two complex numbers in an Argand Diagram

The working for 4.2 is as follows:

$$\text{Let: } z = x + yi \quad (4.1)$$

$$\text{Let: } w = u + vi \quad (4.2)$$

By the addition rule:

$$(x + yi) + (u + vi) = x + u + yi + vi \quad (4.3)$$

$$= (x + u) + (y + v)i \quad (4.4)$$

4.2 Polar Coordinates

1. Polar coordinates use two variables, (r, θ) to describe a point.
2. r is a length.
3. θ is an angle




Figure 4.3: Two polar coordinates. Note how r , the radius, is specified first followed by the angle.

Converting between polar coordinates and cartesian coordinates is a matter of applying some trigonometry but is summarised as follows:

$$x = r \cos \theta \quad (4.5)$$

$$y = r \sin \theta \quad (4.6)$$

Iff $r \geq 0$ and $-\pi \leq \theta < \pi$ we can also convert to Cartesian coordinates as follows:

$$r = \sqrt{x^2 + y^2} \quad (4.7)$$

$$\theta = \begin{cases} \arctan\left(\frac{y}{x}\right) & \text{if } x > 0 \\ \arctan\left(\frac{y}{x}\right) + \pi & \text{if } x < 0 \text{ and } y \geq 0 \\ \arctan\left(\frac{y}{x}\right) - \pi & \text{if } x < 0 \text{ and } y < 0 \\ \frac{\pi}{2} & \text{if } x = 0 \text{ and } y > 0 \\ -\frac{\pi}{2} & \text{if } x = 0 \text{ and } y < 0 \\ 0 & \text{if } x = 0 \text{ and } y = 0 \end{cases} \quad (4.8)$$

4.3 Modulus

Consider diagram 4.4:




Figure 4.4: The modulus, $|z|$, is given by Pythagoras' theorem

4.4 Euler's Formula

Euler's Formula: Euler was this dude who had a formula:

$$e^{i\theta} = \cos\theta + i\sin\theta \quad (4.9)$$

In the context of the unit circle, projected onto an Argand diagram, it starts to form an important relationship shown in figure 4.5. This relationship is further extended in chapter 5 on de Moivre's Theorem (page 25).




Figure 4.5: Euler's Formula plotted on an Argand diagram

DE MOIVRE'S THEOREM & ROOTS OF COMPLEX NUMBERS

Further reading: Chen & Duong “fyc01.pdf”, <http://goo.gl/e0E7y>

Big Ideas:

- de Moivre was this dude who had a theorem named after him, though this specific theorem was never actually mentioned in his work.
- Derived from Euler's Formula (see section 4.4 p 24).

5.1 de Moivre's Theorem

Further reading: Chen & Duong “fyc01.pdf” pp 12.

$$e^{i\theta} = \cos\theta + i\sin\theta \quad (5.1)$$

apply an index law:

$$\left(e^{i\theta}\right)^n = e^{in\theta} \quad \text{iff } n \in \mathbb{Z} \quad (5.2)$$

then by Euler's formula:

$$e^{i(n\theta)} = \cos\theta + i\sin(n\theta) \quad (5.3)$$

Proof that this follows for all integers can be done through a proof by induction. The following proof comes courtesy of Wikipedia, <http://goo.gl/K8nRR>.

$$(\cos\theta + i\sin\theta)^n = \cos(n\theta) + i\sin(n\theta) \quad | \quad n \in \mathbb{Z} \quad (5.4)$$

For $n > 0$, we proceed by principle of mathematical induction. $S(1)$ is true. For $S(k)$ is true where $k \in \mathbb{N}$. That is we assume:

$$(\cos \theta + i \sin \theta)^k = \cos(k\theta) + i \sin(k\theta) \quad (5.5)$$

Consider $S(k+1)$:

$$(\cos \theta + i \sin \theta)^{k+1} = (\cos(\theta) + i \sin(\theta))^k (\cos \theta + i \sin \theta) \quad (5.6)$$

$$= [\cos(k\theta) + i \sin(k\theta)] (\cos \theta + i \sin \theta) \quad (5.7)$$

$$= \cos(k\theta) \cos \theta - \sin(k\theta) \sin \theta + i [\cos(k\theta) \sin \theta + \sin(k\theta) \cos \theta] \quad (5.8)$$

$$= \cos[(k+1)\theta] + i \sin[(k+1)\theta] \quad (5.9)$$

Because $S(k)$ implies $S(k+1)$, by POMI, it follows that the result is true for all \mathbb{N} . For negative integers, consider an index of $-n$ for \mathbb{N} :

$$(\cos \theta + i \sin \theta)^{-n} = [(\cos \theta + i \sin \theta)^n]^{-1} \quad (5.10)$$

$$= [\cos(n\theta) + i \sin(n\theta)]^{-1} \quad (5.11)$$

$$= \cos(-n\theta) + i \sin(-n\theta) \quad (5.12)$$

5.2 Finding Roots of Complex Numbers

Further reading: Chen & Duong “fyc01.pdf” pp 12-14.

5.3 Analytic Geometry

Further reading: Chen & Duong “fyc01.pdf” pp 14-15.

CHAPTER 6

LINEAR EQUATIONS

Further reading: Chen & Duong “1a01.pdf”, <http://goo.gl/N7S2s>

Big Ideas:

- Linear equations are often called a *system of linear equations* when equations are linked together.
- A system of equations can be represented in a *matrix*.

6.1 System of Linear Equations as Matrices

Further reading: Chen & Duong “1a01.pdf” pp 1-3.

6.2 Elementary Row Operations

Further reading: Chen & Duong “1a01.pdf” pp 3-6.

6.3 Row Echelon Form

Further reading: Chen & Duong “1a01.pdf” pp 6-10.

6.4 Reduced Row Echelon Form

Further reading: Chen & Duong “1a01.pdf” pp 11-12.

6.5 Solving a System of Linear Equations

Further reading: Chen & Duong “1a01.pdf” pp 12-15.

6.6 Homogeneous Systems

Further reading: Chen & Duong “la01.pdf” pp 15–16.

CHAPTER



7

MATRICES & LINEAR EQUATIONS

Big Ideas:

-

APPLICATION OF LINEAR EQUATIONS

Big Ideas:

-

8.1 Application to Network Flow

Further reading: Chen & Duong “la01.pdf” pp 16-18.

8.2 Application to Electric Circuits

Further reading: Chen & Duong “la01.pdf” pp 18-21.

8.3 Application to Economics

Further reading: Chen & Duong “la01.pdf” pp 21-22.

8.4 Application to Chemistry

Further reading: Chen & Duong “la01.pdf” pp 22-23.

8.5 Application to Mechanics

Further reading: Chen & Duong “la01.pdf” pp 23-25.

CHAPTER 9

ARITHMETIC OF MATRICES

Big Ideas:

-

DETERMINANT OF A MATRIX**Big Ideas:**

-

VECTORS

Big Ideas:

-

VECTORS & GEOMETRY**Big Ideas:**

-

COMBINATORICS & BINOMIAL THEOREM

Further reading: Chris Cooper “ALG03 The Binomial Theorem”, <http://goo.gl/bqTQH>

Big Ideas:

-

REAL NUMBERS**Big Ideas:**

-

INDUCTION

Big Ideas:

- POMI establishes that a given statement is true for all \mathbb{N} .
- Start by proving (or assuming if given) that the first statement is true. This statement is called the **basis** or **base case**
- Next prove that **any one** statement then so is the next statement. This is known as the **inductive step**.

CHAPTER

16

FUNCTIONS

Big Ideas:

-

CHAPTER

17

LIMITS

Big Ideas:

-

CONTINUITY

Big Ideas:

-

DIFFERENTIABILITY**Big Ideas:**

-

CHAPTER 20

STATIONARY POINTS

Big Ideas:

-

CHAPTER

21

CURVE SKETCHING

Big Ideas:

-

APPLICATIONS OF THE DERIVATIVE**Big Ideas:**

-

CHAPTER

23

ANTIDERIVATIVES

Big Ideas:

-

TECHNIQUES OF INTEGRATION

Big Ideas:

-

SEPARABLE DIFFERENTIABLE EQUATIONS**Big Ideas:**

-

CHAPTER 26

AREA UNDER A CURVE

Big Ideas:

-

VOLUME OF SOLIDS OF REVOLUTION**Big Ideas:**

-

CHAPTER 28

MORE APPLICATIONS OF INTEGRATION

Big Ideas:

-

CHAPTER

29

GLOSSARY

Symbol	Name	ℒ _T ℒ _X
A	Blackboard A	<code>\mathbb{A}</code>
B	Blackboard B	<code>\mathbb{B}</code>
C	Blackboard C	<code>\mathbb{C}</code>
D	Blackboard D	<code>\mathbb{D}</code>
E	Blackboard E	<code>\mathbb{E}</code>
F	Blackboard F	<code>\mathbb{F}</code>
G	Blackboard G	<code>\mathbb{G}</code>
H	Blackboard H	<code>\mathbb{H}</code>
I	Blackboard I	<code>\mathbb{I}</code>
J	Blackboard J	<code>\mathbb{J}</code>
K	Blackboard K	<code>\mathbb{K}</code>
L	Blackboard L	<code>\mathbb{L}</code>
M	Blackboard M	<code>\mathbb{M}</code>
N	Blackboard N	<code>\mathbb{N}</code>
O	Blackboard O	<code>\mathbb{O}</code>
P	Blackboard P	<code>\mathbb{P}</code>
Q	Blackboard Q	<code>\mathbb{Q}</code>
R	Blackboard R	<code>\mathbb{R}</code>
S	Blackboard S	<code>\mathbb{S}</code>
T	Blackboard T	<code>\mathbb{T}</code>
U	Blackboard U	<code>\mathbb{U}</code>
V	Blackboard V	<code>\mathbb{V}</code>
W	Blackboard W	<code>\mathbb{W}</code>
X	Blackboard X	<code>\mathbb{X}</code>
Y	Blackboard Y	<code>\mathbb{Y}</code>
Z	Blackboard Z	<code>\mathbb{Z}</code>

Table 29.1: Blackboard letters. Blackboard notation must be inside math mode.

Greek Uppercase	Lowercase	Name	\LaTeX (Upper)	\LaTeX (Lower)
A	α	Alpha	A	$\backslash\alpha$
B	β	Beta	B	$\backslash\beta$
Γ	γ	Gamma	$\backslash\Gamma$	$\backslash\gamma$
Δ	δ	Delta	$\backslash\Delta$	$\backslash\delta$
E	ϵ	Epsilon	E	$\backslash\epsilon$
	ε	Epsilon Variant		$\backslash\varepsilon$
Z	ζ	Zeta	Z	$\backslash\zeta$
H	η	Eta	H	$\backslash\eta$
Θ	θ	Theta	$\backslash\Theta$	$\backslash\theta$
	ϑ	Theta Variant		$\backslash\vartheta$
I	ι	Iota	I	$\backslash\iota$
K	κ	Kappa	K	$\backslash\kappa$
	\varkappa	Kappa Variant		$\backslash\varkappa$
Λ	λ	Lambda	$\backslash\Lambda$	$\backslash\lambda$
M	μ	Mu	M	$\backslash\mu$
N	ν	Nu	N	$\backslash\nu$
Ξ	ξ	Xi	$\backslash\Xi$	$\backslash\xi$
O	\omicron	Omicron	O	\omicron
Π	π	Pi	$\backslash\Pi$	$\backslash\pi$
	ϖ	Pi Variant		$\backslash\varpi$
P	ρ	Rho	P	$\backslash\rho$
	ϱ	Rho Variant		$\backslash\varrho$
Σ	σ	Sigma	$\backslash\Sigma$	$\backslash\sigma$
	ς	Sigma Variant		$\backslash\varsigma$
T	τ	Tau	T	$\backslash\tau$
Υ	υ	Upsilon	$\backslash\Upsilon$	$\backslash\upsilon$
Φ	ϕ	Phi	$\backslash\Phi$	$\backslash\phi$
	φ	Phi Variant		$\backslash\varphi$
X	χ	Chi	X	$\backslash\chi$
Ψ	ψ	Psi	$\backslash\Psi$	$\backslash\psi$
Ω	ω	Omega	$\backslash\Omega$	$\backslash\omega$

Table 29.2: The Greek alphabet.

Symbol	My \LaTeX	\LaTeX	Description
\Re	$\backslash\Re$	$\backslash\operatorname{\mathfrak{Re}}$	Real part of a number
\Im	$\backslash\Im$	$\backslash\operatorname{\mathfrak{Im}}$	Imaginary part of a number

Table 29.3: Real and Imaginary symbols