

Table of Contents

S.no	Group	Constraint	Layout	Title	Title Key	Population Set	Treatment Arm Set	Statistic Set
1	Core	Required	[D] (Default)	Table 15.8.1.3.1: Participant Disposition	t-s-disp	All Screened Participants	Planned Trt	Set 1 (N), Set 2 (n, %)
2	Core	Required	[D] (Default)	Table 15.8.3.1: Demographics and Baseline Characteristics	t-s-demog	ITT Analysis Set	Planned Trt	Set 2 (n, %), Ct [Nb] / Cn [n, Mean (SD), Median, Q1, Q3, Min, Max]
3	Oncology	Required	[D] (Default)	Table 15.9.1.x: Overall Survival (OS)	t-os	ITT Analysis Set	Planned Trt	{Stratified} Hazard Ratio {of A vs B} (95% CI), Set 4 [Q1, Median, Q3], Set 2 (n, %), Set 34 [n, Mean (StD), Median, Q1, Q3, Min, Max], Set 10 (P value 0.XXXX), Set 17 XX.X% (XX.X%, XX.X%)
4	Core	Required	[D1] (Default Layout 1)	Table 15.11.2.1.1: Treatment-Emergent Adverse Events: Overall Summary	t-s-aebrief	Safety Analysis Set	Actual Trt	Set 2 (n, %)
5	Core	Conditional	[D] (Default)	Table 15.10.1.1.x: Individual Data and Summary Statistics of Plasma Concentration (ng/mL) at Protocol-Specified Sampling Times	t-pkpc	PK Analysis Set	PK Sampling 0-24hrs	N [XX], Set 55 XX.X, Set 31 PK [n, Mean, SD, % CV], Set 39 XXX.XXX, Set 1 (N), Set 56 [XXX.XX]

TABLES: Disposition

PROGRAMMING NOTES

Table 15.8.1.3.1 Participant Disposition [Default],[Required]

1. For rows where the number of participants in each analysis set is presented, base the number of participants on applicable participant treatment grouping (eg, use TRTxP for the ITT Analysis Set or the FAS Analysis Set row and TRTxXA for the Safety Analysis Set or the PK Analysis Set row). Add rows to include all analysis sets specified in the SAP. Use the [ITT][Full] Analysis Set and TRTxP for the denominator in studies where the [ITT][Full] Analysis Set is the primary efficacy analysis set; otherwise, use the Safety Analysis Set and TRTxXA.
2. Reasons for Premature Discontinuation should match the study eCRF. As a default, categories with all zeros are omitted. When it adds value, all categories as listed on the eCRF may be displayed, including categories with all zeros.
3. Only display "Reason for [Premature] Discontinuation of [Any] Study [Drug] [Treatment]" section if any participants discontinued study drug. Only display "Reason for [Premature] Study Discontinuation" section if any participants discontinued the study.
4. "Continuing Study [Drug][Treatment]", "Continuing Study", and completion at specific time points may be added for interim analyses.
5. Denominator for categories under a treatment phase should be the number of participants who entered the respective treatment phase.
6. Sort the reasons for premature discontinuation by descending frequency under the total column. If the frequency is tied, then sort by alphabetical order.
7. The annotations of 'TRTxP(N)', 'TRTxXA(N)', 'AREL(N)' and other similar ones in the tables and listings indicate that either the character (eg, TRTxP, AREL) or the numeric (eg, TRTxPN, ARELN) variable can be used to create the column or row.

Table 15.8.1.3.1
Participant Disposition

(ADSL.SCRNFL == "Y")

All Screened Participants

	ADAM / ADSL / TRT01PN (ADSL.TRT01PN == 1) Drug A	ADAM / ADSL / TRT01PN (ADSL.TRT01PN == 2) Drug B
ADAM / ADSL / SCRNFL (ADSL.SCRNFL == "Y") Screened	XXX	XXX
ADAM / ADSL / ITTFL (ADSL.ITTFL == "Y") ITT Analysis Set	XXX	XXX
ADAM / ADSL / SAFFL (ADSL.SAFFL == "Y") Safety Analysis Set	XXX	XXX
ADAM / ADSL / PKFL (ADSL.PKFL == "Y") PK Analysis Set	XXX	XXX
ADAM / ADSL / COMTFL Study Drug Completion Status		
ADAM / ADSL / COMTFL (ADSL.COMTFL == "Y") Completed Study Drug	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / COMTFL (ADSL.COMTFL == "N")	XX (XX.X%)	XX (XX.X%)

Discontinued Study Drug		
ADAM / ADSL / DCTREAS Reason for Discontinuation of Study Drug		
ADAM / ADSL / DCTREAS (ADSL.DCTREAS == "ADVERSE EVENT") Adverse Event	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / DCTREAS (ADSL.DCTREAS == "LACK OF EFFICACY") Lack of Efficacy	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / COMSFL Study Completion Status		
ADAM / ADSL / COMSFL (ADSL.COMSFL == "Y") Completed Study	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / COMSFL (ADSL.COMSFL == "N") Prematurely Discontinued Study	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / DCSREAS Reason for Discontinuation of Study		
ADAM / ADSL / DCSREAS (ADSL.DCSREAS == "DEATH") Death	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / DCSREAS (ADSL.DCSREAS == "WITHDRAWAL BY SUBJECT") Withdrawal by Subject	XX (XX.X%)	XX (XX.X%)

Percentages for completion status were calculated based on the number of participants in the Safety Analysis Set.
Source: Listing 16.2.1.3.1

TABLES: Demographics

PROGRAMMING NOTES

Table 15.8.3.1 Demographics and Baseline Characteristics [Default],[Required]

1. ITT or All Enrolled analysis set may be used instead of Safety for studies that use the ITT or All Enrolled analysis set as the primary analysis set. For multiphase studies, [Study Specific] analysis set may be used based on the study design.
2. The following conventions will be used for the imputation of date of birth when it is partially missing or not collected:
 - If only month and year of birth is collected, then "15" will be imputed as the day of birth
 - If only year of birth is collected, then "01 July" will be imputed as the day and month of birth
 - If year of birth is missing, then date of birth will not be imputed.
- In general, age collected at Day 1 (in years {or specify unit}) will be used for analyses and presented in listings. If age at Day 1 is not available for a participant, then age derived based on date of birth and the Day 1 visit date will be used instead. If an enrolled participant was not dosed with any study drug, the randomization [enrollment] date will be used instead of the Day 1 visit date.
- For screen failures, the date the first informed consent was signed will be used for the age derivation. Age required for longitudinal and temporal calculations and analyses (e.g., estimates of creatinine clearance, age at date of AE) will be based on age derived from date of birth and the date of the measurement or event, unless otherwise specified. (Note, see SAP and discuss with Biometrics team).
3. Baseline characteristics (e.g., weight, height, and BMI) are presented in the same table with the demographics.
4. Height, weight, and/or body mass index (BMI) may also be displayed by sex if specified in the SAP. Derived BMI values will be reported instead of the collected value from eCRF.
5. Optional row for missing or unknown category may be added where applicable (e.g., interim analyses) and dropped for final analysis if all zero counts.
6. Categories for race and ethnicity should follow the order in the study eCRF.
7. As a default, categories with all zeros are omitted. If they add value, all categories as listed on the CRF may be displayed, including categories with all zeros.
8. Missing values may be included in percent calculations so percentages should add to 100%.
9. *Use TRTxX(A(N) when displaying the safety analysis set, and TRTxXP(N) otherwise.
10. The values of this variable cannot be used directly to satisfy the requirements of the table. An additional variable should be created in the table program to change to case or associate a format so that the display matches the mockup display.
11. Categories for age group and BMI group may be added if specified in the SAP. Up to a maximum of three age groups is permitted in the display.

Table 15.8.3.1
Demographics and Baseline Characteristics

(ADSL.ITTFL == "Y")

ITT Analysis Set

	ADAM / ADSL / TRT01PN (ADSL.TRT01PN == 1) Drug A	ADAM / ADSL / TRT01PN (ADSL.TRT01PN == 2) Drug B
ADAM / ADSL / AGE ADAM / ADSL / AGEU Age (years)		
n	XX	XX
Mean (StD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X
ADAM / ADSL / AGEGR1N Age Group		
ADAM / ADSL / AGEGR1N (ADSL.AGEGR1N == 1) < 65 years	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / AGEGR1N (ADSL.AGEGR1N == 2) ≥ 65 years	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / SEX Sex at Birth		
(ADSL.SEX == "M") Male	XX (XX.X%)	XX (XX.X%)

(ADSL.SEX == "F") Female	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / RACE Race		
(ADSL.RACE == "AMERICAN INDIAN OR ALASKA NATIVE") American Indian or Alaska Native	XX (XX.X%)	XX (XX.X%)
(ADSL.RACE == "ASIAN") Asian	XX (XX.X%)	XX (XX.X%)
(ADSL.RACE == "BLACK OR AFRICAN AMERICAN") Black	XX (XX.X%)	XX (XX.X%)
(ADSL.RACE == "NATIVE HAWAIIAN OR OTHER PACIFIC ISLANDER") Native Hawaiian or Pacific Islander	XX (XX.X%)	XX (XX.X%)
(ADSL.RACE == "WHITE") White	XX (XX.X%)	XX (XX.X%)
(ADSL.RACE == "OTHER") Other	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / ETHNIC Ethnicity		
(ADSL.ETHNIC == "NOT HISPANIC OR LATINO") Not Hispanic or Latino	XX (XX.X%)	XX (XX.X%)
(ADSL.ETHNIC == "HISPANIC OR LATINO") Hispanic or Latino	XX (XX.X%)	XX (XX.X%)
ADAM / ADSL / BWT Weight (kg)		
n	XX	XX
Mean (StD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X

ADAM / ADSL / BHT Height (cm)		
n	XX	XX
Mean (StD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X
ADAM / ADSL / BBMI Body Mass Index (kg/m^2)		
n	XX	XX
Mean (StD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X

Denominator for percentages was the ITT Analysis Set.

Age (in years{or specify unit}) was collected on date of Day 1 visit. [If age was not collected, age was derived using data of Day 1 visit if dose, date of [randomization] [enrollment] if not dosed, or the date of informed consent if not randomized.

Body Mass Index (BMI) (kg/m^2) = [Weight (kg)/Height (cm) ^2] * 10,000.

Source: Listing 16.2.4.1

TABLES: Efficacy Analysis

PROGRAMMING NOTES

Table 15.9.1.x Overall Survival (OS) [Default],[Required]

1. This table shell can be applied based on
 - IRC/INV assessment (PFS)
 - ITT/FAS/PP analysis set
 - Subgroup analysis
 - Sensitivity analysis
 - Unstratified/stratified analysis
2. There may be different categories under “Number (%) of participants censored” depending on specific study specifications. Zero rows in the pre-defined categories will be shown.
3. Refer to study statistician’s definition on the event/censoring rules, and also the censored categories based on different endpoints and analysis.
4. For HR and its 95% CI, please use the same parameter for number of decimal digits. If no values set for the parameter, please display 2 decimal digits by default.
5. For survival rates and its 95% CI, please use the same parameter for number of decimal digits. If no values set for the parameter, please display 3 decimal digits by default.
6. For median, Q1, Q3, and the corresponding 95% CI, please use the same parameter for number of decimal digits. If no values set for the parameter, please display 1 decimal digit.
7. Line “P value (Non-Inferiority Test)” is optional for the non-inferiority study. In that case, the next line for P value from the superiority test will be displayed only if the non-inferiority test is rejected.
8. If there are multiple treatment groups, in general, the pairwise treatment comparison will be conducted by only including the treatment groups which are compared with, unless otherwise specified by SAP. For example, if the comparison is for treatment A and B, then data from treatment group C will be excluded when doing the comparison.
9. The duration of OS/PFS follow-up only applies to the whole analysis set, not the sub-group analysis set.
10. The formula for calculating OS/PFS and the duration of OS/PFS follow-up are the same at individual participant level. At study level, the duration of OS follow-up is calculated using summary statistics (n, mean, std, median, Q1, Q3, minimum and maximum).
11. The reasons listed under “number of participants censored” should be mutually exclusive and modified based on the endpoint (PFS or OS) being analyzed and the type of analysis (primary or sensitivity). In the table template, the reasons for censoring for the primary analysis of the endpoint (PFS or OS) are listed.

Sample Code:

```
• For stratified log-rank test:
ods output HomTests={logrank};
proc lifetest data={indata} outsurv={_surv} timelist=xx reduceout;
  time {SurTime}*{Censored(1)};
  strata {strata1} {strata2}/ group={trtgrp} diff=all test=logrank;
run;
ods output close;
```

- For Un-stratified log-rank test and KM estimates at time points:

```

ods output HomTests={logrank};
proc lifetest data={indata} outsurv={_surv} timelist=xx reduceout;
  time {SurTime}*{Censored(1)};
  strata {trtgrp} / diff=all test=logrank;
run;
ods output close;

• Hazard ratio:
ods output HazardRatios={hr} estimates={test};
proc phreg data={indata};
  class {trtgrp} (ref = "{B}"); /*specify the reference arm in (ref='')*/
  strata {strata1} {strata2}; /*only needed for stratified test*/
  model {SurTime}*{Censored(1)}={trtgrp} / rl alpha=0.05 ties=efron;
  hazardratio {trtgrp} / alpha=0.05 diff=ref;
  estimate {trtgrp} {1} / testvalue=Ln_HR0; /*only needed for non-inferiority test. "estimate {trtgrp} 1" assigns 1 as the coefficient to be applied to the
non-inferiority statistical test assign Ln_HR0=ln(HR0) where HR0 is the non-inferiority test boundary*/
run;

```

Table 15.9.1.x
Overall Survival (OS)

(ADSL.ITTFL == "Y")

ITT Analysis Set

	ADAM / ADSL / TRT01PN (ADSL.TRT01PN == 1) Drug A	ADAM / ADSL / TRT01PN (ADSL.TRT01PN == 2) Drug B
ADAM / ADTTE / CNSR (ADTTE.CNSR == 0) Number (%) of Participants with Events (Death)	XX (XX.X%)	XX (XX.X%)
ADAM / ADTTE / CNSR (ADTTE.CNSR NE 1) Number (%) of Participants Censored	XX (XX.X%)	XX (XX.X%)
ADAM / ADTTE / AVAL Kaplan-Meier Estimate of OS (Months) (95% CI) [a]		
Q1	XX.X (XX.X, XX.X)	XX.X (XX.X, XX.X)
Median	XX.X (XX.X, XX.X)	XX.X (XX.X, XX.X)
Q3	XX.X (XX.X, XX.X)	XX.X (XX.X, XX.X)
ADAM / ADTTE / AVAL Kaplan-Meier Estimate of OS Rate (%) (95% CI)		
ADAM / ADTTE / AVAL At 6 Months	XX.X% (XX.X%, XX.X%)	XX.X% (XX.X%, XX.X%)
ADAM / ADTTE / AVAL At 12 Months	XX.X% (XX.X%, XX.X%)	XX.X% (XX.X%, XX.X%)

ADAM / ADTTE / AVAL At 18 Months	XX.X% (XX.X%, XX.X%)	XX.X% (XX.X%, XX.X%)
ADAM / ADTTE / AVAL Hazard Ratio of A vs B (95% CI) [b]	0.XX (0.XX, NEst)	
ADAM / ADTTE / AVAL P value [c]	0.XXXX	
ADAM / ADTTE / AVAL Duration of OS Follow-up (Months) [d]		
n	XX	XX
Mean (StD)	XX.X (XX.XX)	XX.X (XX.XX)
Median	XX.X	XX.X
Q1, Q3	XX.X, XX.X	XX.X, XX.X
Min, Max	XX.X, XX.X	XX.X, XX.X
ADAM / ADTTE / AVAL		
(ADTTE.AVAL GE 6) >= 6 Months	XX (XX.X%)	XX (XX.X%)
(ADTTE.AVAL GE 12) >= 12 Months	XX (XX.X%)	XX (XX.X%)
(ADTTE.AVAL GE 18) >= 18 Months	XX (XX.X%)	XX (XX.X%)

{Definition of Analysis set.}

NEst = Not Estimable.

a OS (months) = (date of death or censoring - date of {randomization/first dose} + 1)/30.4375, and the 95% CI is derived using log-log transformation.

b Hazard ratio and 95% CIs are calculated using the Cox proportional hazards model / statistical method.

c P value for comparing the overall survival functions from the two treatment groups is from.

d Duration of OS follow-up (months) = (date of death or censoring - date of randomization + 1)/30.4375.

Source: Listing 16.2.6.x

TABLES: Adverse Events

PROGRAMMING NOTES

Table 15.11.2.1.1 Treatment-Emergent Adverse Events: Overall Summary [Default Layout 1],[Required]

Refer to the TFL Designer 'GCDS Template Instructions' tab, under the 'Data Presentation' heading, for 'Adverse Events Tables Programming Notes'; these notes contain example tables for points 17 and 18, below, with the logic for analysis counts.

GENERAL:

1. The rows presented in the Overall AE Summary table should match the AE outputs presented in the study. A row for Death ADSL.DTHFL is required in the Overall AE Summary table unless a separate Death summary table is presented in the study.
2. If the Overall AE Summary table is generated using a Subgroup analysis, then each category of the subgroup will displayed on a separate page as the first row of that page.
3. For TEAEs of interest, optional rows may be added to the Overall AE Summary table. However, for multiple categories of TEAEs of interest a separate Overall AE Summary table for TEAEs of interest may be used, then each category of interest will be displayed on a separate page as the first row of that page.
4. Use the Safety ADSL.SAFFL Analysis Set for Adverse Event tables.
5. Only TEAEs will be summarized unless otherwise specified; TEAEs will be defined as any AEs that occurred on or after the first study drug dosing date and up to {XX} days {or study drug-specific therapeutic reach} after the date of the last dose of the study drug [and prior to the day of initiation of subsequent anticancer/antiviral/etc. therapy], unless otherwise specified. For interim analyses, refer to SAP for participants who are ongoing on study drug.
6. If the onset date of the AE is incomplete and the AE stop date is not prior to the first dosing date of study drug, then the month and year (or year alone if month is not recorded) of onset determine whether an AE is treatment emergent. The event is considered treatment emergent if both of the following 2 criteria are met:
 - The AE onset is the same as or after the month and year (or year) of the first dosing date of study drug, and
 - The AE onset date is the same as or before the month and year (or year) of the date corresponding to {XX} days {or study drug-specific therapeutic reach} after the date of the last dose of study drug [and prior to the day of initiation of subsequent anticancer/antiviral/etc. therapy].An AE with completely missing onset and stop dates, or with the onset date missing and a stop date later than the first dosing date of study drug, will be considered to be treatment emergent. In addition, an AE with the onset date missing and incomplete stop date with the same or later month and year (or year alone if month is not recorded) as the first dosing date of study drug will be considered treatment emergent.
7. Default sort order is alphabetically for SOC, [and HLT within SOC,] and then descending frequency of PTs in the total column. Summaries that include PT only, sort order is by descending frequency of PTs in the total column. Optionally, a different sort order maybe specified as based on the study design.
8. For studies that collect adverse event data for individual study drugs, the layout may be altered to summarize the individual study drugs separately.
9. A total column may be presented for studies where different doses of the same study drug are displayed.
10. Rows may be added to summarize the number of participants with missing data, as applicable.
11. Lower severity grades may be excluded from table summaries, as applicable.
12. A P value may be displayed if pre-specified in the SAP. Display P values in a column if 2 treatment groups are shown in the table, or in a row if more than 2 treatment groups are shown.
13. For Oncology studies, refer to the SAP when determining relevant custom grouping for lab abnormality AE terms (e.g., ADAE.AEDECOD are mapped: Neutrophil count decreased->Neutropenia, White blood cell count decreased->Leukopenia, Lymphocyte count decreased->Lymphopenia, Red blood cell count decreased->Anaemia, Platelet count decreased->Thrombocytopenia). Related footnote should be added to explain this.
14. For studies where the analysis is based on the Treatment Emergent within each period, then in such instances the variable ADAE.TREMxxFL can be utilized to summarize the AE's. [xx] in the variable name indicates each period and will be replaced by 01, 02, 03, etc.
15. Rows for treatment mean difference and P value are optional for Exposure-Adjusted Incidence Rate (EAIR) summaries. Refer to the SAP when determining relevant statistical methods.
16. For Oncology studies, the required displays are separate summaries of 'TEAE with Grade 3 or 4' and 'TEAE Leading to Death' (i.e., TEAE with Grade 5) instead of the combined summaries of 'TEAE with Grade 3 or Higher'. For Virology and Inflammation studies, the preferred display is the combined summaries of 'TEAE with Grade 3 or Higher' and separate summaries of 'TEAE with Grade 3 or 4' and 'TEAE Leading to Death' (i.e., TEAE with Grade 5) are optional only.
17. <Refer to 'Adverse Events Tables Programming Notes' in the 'GCDS Template Instructions' tab>
For Table 15.11.2.1.1 TEAE Overall Summary, the following rows are intended to count participants once only and at the highest severity grade experienced per participant throughout the study:
 - TEAE with Grade 3 or 4
 - TEAE Leading to DeathIf the highest severity grade experienced is a Grade 5, then those participants would be excluded from the "TEAE with Grade 3 or 4" row (but would be counted in the "TEAE Leading to Death" row).
For summaries related to study drug, subset records on relationship to study drug (ADAE.AREL='RELATED') first, before determining the highest severity grade experienced per participant. Participants are counted once only in the following rows at the highest severity grade experienced per participant of the drug-related TEAEs. If the highest severity grade of the drug-related TEAEs is a Grade 5, the participant is excluded from the "TEAE Related to Study Drug with Grade 3 or 4" row (but is counted in the "TEAE Related to Study Drug Leading to Death" row).
 - TEAE Related to Study Drug with Grade 3 or 4
 - TEAE Related to Study Drug Leading to Death

Example Study: In a 20-participant study, 20 participants have treatment-emergent maximum Grade 4 neutropenia and 2 participants have treatment-emergent maximum Grade 5 febrile neutropenia. All events are considered treatment-related except one case of Grade 5 febrile neutropenia. Refer the following summary table displays the expected participant counts:

18. <Refer to 'Adverse Events Tables Programming Notes' in the 'GCDS Template Instructions' tab>

For Table 15.11.2.2.2.1 Grade 3 or 4 TEAEs by SOC and PT, in the first summary row, include participants who experienced a maximum severity of Grade 3 or 4 throughout the study (ADAE.AOCCIFL='Y' and ADAE.ATOXGRN in (3, 4)) and exclude participants who experienced a maximum severity of Grade 5 - this matches the "TEAE with Grade 3 or 4" row in the Overall Summary table. Subsequent rows for SOC, exclude participants who experienced Grade 5 when counting the highest severity grade within this SOC (ADAE.AOCCSIFL='Y' and ADAE.ATOXGRN in (3, 4)). Subsequent rows for PT, exclude participants who experienced Grade 5 when counting the highest severity grade within this PT (ADAE.AOCCPIFL='Y' and ADAE.ATOXGRN in (3, 4)).

Additionally, for Table 15.11.2.3.3.1 Grade 3 or 4 Treatment-Related TEAEs by SOC and PT, subset the records on relationship to study drug (ADAE.AREL='RELATED') first, before determining the highest severity grade (i.e., Grade 3 or 4) per participant (first row - matches Overall Summary table) and for each SOC and PT when selecting records for the relevant 1st Max Sev./Int. and Related to Study Drug Occurrence Flag (ADAE.AOCCzzFL).

Table 15.11.2.1.1
Treatment-Emergent Adverse Events: Overall Summary
 (ADSL.SAFFL == "Y")
Safety Analysis Set

Number (%) of Participants with	ADAM / ADSL / TRT01AN (ADSL.TRT01AN == 1) Drug A	ADAM / ADSL / TRT01AN (ADSL.TRT01AN == 2) Drug B
ADAM / ADAE / TRTEMFL (ADAE.TRTEMFL == "Y") Any TEAE	XX (XX.X%)	XX (XX.X%)
ADAM / ADAE / ATOXGR (ADAE.ATOXGRN GE 3) TEAE with Grade 3 or Higher	XX (XX.X%)	XX (XX.X%)
ADAM / ADAE / ATOXGR (ADAE.ATOXGRN GE 2) TEAE with Grade 2 or Higher	XX (XX.X%)	XX (XX.X%)
ADAM / ADAE / AREL (ADAE.AREL == "RELATED") TEAE Related to Study Drug	XX (XX.X%)	XX (XX.X%)
ADAM / ADAE / ATOXGR (ADAE.AREL == "RELATED") && (ADAE.ATOXGRN GE 3) TEAE Related to Study Drug with Grade 3 or Higher	XX (XX.X%)	XX (XX.X%)
ADAM / ADAE / ATOXGR (ADAE.AREL == "RELATED") && (ADAE.ATOXGRN GE 2) TEAE Related to Study Drug with Grade 2 or Higher	XX (XX.X%)	XX (XX.X%)
ADAM / ADAE / AESER (ADAE.AESER == "Y") TE Serious AE	XX (XX.X%)	XX (XX.X%)
ADAM / ADAE / AESER (ADAE.AESER == "Y") && (ADAE.AREL == "RELATED") TE Serious AE Related to Study Drug	XX (XX.X%)	XX (XX.X%)
ADAM / ADAE / AEACN (ADAE.AEACN == "DRUG WITHDRAWN")	XX (XX.X%)	XX (XX.X%)

TEAE Leading to Discontinuation of Study Drug		
ADAM / ADAE / AEACN (ADAE.AEACN == "DOSE INTERRUPTED") TEAE Leading to Dose Interruption of Study Drug	XX (XX.X%)	XX (XX.X%)
ADAM / ADAE / AEACN (ADAE.AEACN == "DOSE REDUCED") TEAE Leading to Dose Reduction of Study Drug	XX (XX.X%)	XX (XX.X%)
ADAM / ADAE / TRTEMFL (ADAE.AESDTH == "Y") TEAE Leading to Death	XX (XX.X%)	XX (XX.X%)

TE = treatment-emergent; TEAE = treatment-emergent adverse event.

Adverse events were coded according to MedDRA Version {XX.X}.

Treatment-emergent events began on or after the study drug start date up to {XX} days {or study drug-specific therapeutic reach} after the date of the last dose of study drug [and prior to the day of initiation of subsequent anticancer/antiviral/etc therapy].

Severity grades were defined by [Gilead Grading Scale for Severity of Adverse Events and Laboratory Abnormalities][The CTCAE Version X][The DAIDS Version XX]. [Participants were counted only one based on their highest grade for each summary row.]

Death includes any death that occurred during the study. Percentages for TEAE Dose Limiting Toxicity was calculated based on the number of participants in the DLT Evaluable Set.

Source: Listing 16.2.7.1, 16.2.7.2.1, 16.2.7.2.2, 16.2.7.3, 16.2.7.4.x and 16.2.7.5.x

BIOSTAT NOTES: If not using the severity grading system, replace 'AE with Grade >= 3' with 'Severe AE' and delete severity grade footnote.

TABLES: Pharmacokinetic

PROGRAMMING NOTES

Table 15.10.1.1.x Individual Data and Summary Statistics of Plasma Concentration (ng/mL) at Protocol-Specified Sampling Times [Default],[Conditional]

1. Only present intensive PK sampling data in this listing.
2. A unique symbol with corresponding footnote should be used for each reason for data exclusion.
3. Geometric mean is optional should only be produced at the request of Clinical Pharmacology and is not presented as a default.
4. PK concentration values will be presented as provided and summary statistics will be presented to 3 significant digits. Please research how significant digits are done if you are not familiar with using them.
5. The Population Set may be displayed as "{Analyte}{Intensive} PK [Substudy] Analysis Set" using ADSL.PKFL or ADSL.PKyFL.
6. The by-groups for tables 15.10.1.1.x: and 15.10.1.2.x "Treatment: {GS-XXXX (XXmg)}" may be displayed using ADPC.TRTP(N) or ADSL.TRTxxP(N).

Table 15.10.1.1.x

Individual Data and Summary Statistics of Plasma Concentration (ng/mL) at Protocol-Specified Sampling Times

(ADSL.PKFL == "Y")

PK Analysis Set

ADAM / ADPC / TRT01AN
(ADPC.TR01AN == 1) || (ADPC.TR01AN == 2)

Treatment: Drug A

Participant ID	Sampling Time (h) Postdose								
	ADAM / ADPC / ATPT (ADPC.ATPT == "Pre-dose") Predose	ADAM / ADPC / ATPT (ADPC.ATPT == "1 Hour") 1h	ADAM / ADPC / ATPT (ADPC.ATPT == "2 Hours") 2h	ADAM / ADPC / ATPT (ADPC.ATPT == "4 Hours") 4h	ADAM / ADPC / ATPT (ADPC.ATPT == "8 Hours") 8h	ADAM / ADPC / ATPT (ADPC.ATPT == "12 Hours") 12h	ADAM / ADPC / ATPT (ADPC.ATPT == "16 Hours") 16h	ADAM / ADPC / ATPT (ADPC.ATPT == "18 Hours") 18h	ADAM / ADPC / ATPT (ADPC.ATPT == "24 Hours") 24h
ADAM / ADPC / SUBJID ADAM / ADPC / AVALC 001	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX
ADAM / ADSL / SUBJID ADAM / ADPC / AVALC 002	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX
ADAM / ADSL / SUBJID . . .	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX	XXX.XX
ADAM / ADPC / AVAL n	XX	XX	XX	XX	XX	XX	XX	XX	XX
ADAM / ADPC / AVAL	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX

Mean									
ADAM / ADPC / AVAL StD	XXX.XXXX	XXX.XXXX	XXX.XXXX	XXX.XXXX	XXX.XXXX	XXX.XXXX	XXX.XXXX	XXX.XXXX	XXX.XXXX
ADAM / ADPC / AVAL % CV	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X
90% CI (Lower)	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX
ADAM / ADPC / USUBJID 90% CI (Upper)	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX
ADAM / ADPC / AVAL Median	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX
ADAM / ADPC / AVAL Min	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX
ADAM / ADPC / AVAL Max	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX
ADAM / ADPC / AVAL Q1	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX
ADAM / ADPC / AVAL Q3	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX	XXX.XXX
ADAM / ADPC / AVAL N (LN scale)	XX	XX	XX	XX	XX	XX	XX	XX	XX
ADAM / ADPC / AVAL Geom. Mean	XXX	XXX	XXX	XXX	XXX	XXX	XXX	XXX	XXX
ADAM / ADPC / AVAL	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X

% GCV									
ADAM / ADPC / ASTRESC (ADPC.AVALC == "BLQ") # BLQ	XX	XX	XX	XX	XX	XX	XX	XX	XX
ADAM / ADPC / ASTRESC (ADPC.AVALC == "BLQ") >1/3 BLQ	XXX	XXX	XXX	XXX	XXX	XXX	XXX	XXX	XXX
ADAM / ADPC / ASTRESC (ADPC.AVALC == "BLQ") % BLQ	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X	XX.X

BLQ = below the limit of quantitation; LLOQ = lower limit of quantitation; CI = confidence interval.

Values BLQ were treated as 0 for predose and postdose.

If > 1/3 of postdose values are BLQ at a time point, then the mean, StD, and %CV are not calculated at that time point.

LLQ was defined as 5.0 ng/mL for analyte GS-XXXX.

For each participant in PK analysis set, intensive serial PK sample collection (samples over 24-hr postdose) was performed at Day 1.