

Table 15.8.1.3.1.1: Participant Disposition
All Screened Participants

	Drug A (N=50)	Drug B (N=50)	Total (N=100)
Screened			100
ITT Analysis Set	50	50	100
Safety Analysis Set	50	50	100
PK Analysis Set	50	50	100
Study Drug Completion Status			
Completed Study Drug	15 (30.0%)	18 (36.0%)	33 (33.0%)
Discontinued Study Drug	35 (70.0%)	32 (64.0%)	67 (67.0%)
Reason for Discontinuation of Study Drug			
Adverse Event	26 (52.0%)	25 (50.0%)	51 (51.0%)
Lack of Efficacy	9 (18.0%)	7 (14.0%)	16 (16.0%)
Study Completion Status			
Completed Study	20 (40.0%)	22 (44.0%)	42 (42.0%)
Prematurely Discontinued Study	30 (60.0%)	28 (56.0%)	58 (58.0%)
Reason for Discontinuation of Study			
Death	26 (52.0%)	25 (50.0%)	51 (51.0%)
Withdrawal by Subject	4 (8.0%)	3 (6.0%)	7 (7.0%)

Percentages for completion status were calculated based on the number of participants in the Safety Analysis Set.
Source: Listing 16.2.1.3.1

Data Extracted: ddmmmyyy

Source: .../final/draft1/prog/t-disp.sas v9.4 Output file: t-disp.pdf 30AUG2026:15:55