

Table 15.9.1.x: Overall Survival (OS)
ITT Analysis Set

	Drug A (N=50)	Drug B (N=50)
Number (%) of Participants with Events (Death)	26 (52.0%)	25 (50.0%)
Number (%) of Participants Censored	24 (48.0%)	25 (50.0%)
Kaplan-Meier Estimate of OS (Months) (95% CI) [a]	15.0 (10.1, NEst)	16.0 (11.0, NEst)
Kaplan-Meier Estimate of OS Rate (%) (95% CI)		
At 6 Months	83.6 (69.9, 91.5)	86.0 (72.9, 93.1)
At 12 Months	62.5 (47.3, 74.5)	60.8 (45.5, 73.0)
At 18 Months	42.2 (27.3, 56.4)	46.0 (31.0, 59.8)
Hazard Ratio of A vs B (95% CI) [b]	1.12 (0.65, 1.94)	
Log-Rank P Value [c]	0.6805	
Duration of OS Follow-up (Months) [d]		
>= 6 Months	40 (80.0%)	43 (86.0%)
>= 12 Months	27 (54.0%)	27 (54.0%)
>= 18 Months	9 (18.0%)	12 (24.0%)

NEst = Not Estimable.

a OS (months) = (date of death or censoring - date of randomization + 1)/30.4375, and the 95% CI is derived using log-log transformation.

b Hazard ratio and 95% CIs are calculated using the Cox proportional hazards model / statistical method.

c P value for comparing the overall survival functions from the two treatment groups is from.

d Duration of OS follow-up (months) = (date of death or censoring - date of randomization + 1)/30.4375.

Source: Listing 16.2.6.x

Data Extracted: ddmmyyyy

Source: .../final/draft1/prog/t-os.sas v9.4 Output file: t-os.pdf 30AUG2026:15:52