

Table 15.10.1.1.x: Individual Data and Summary Statistics of Plasma Concentration (ng/mL) at Protocol-Specified Sampling Times
PK Analysis Set

Treatment: Drug A

Participant ID	Pre-dose	Sampling Time (h)							
		1h	2h	4h	8h	12h	16h	18h	24h
100-006	5.03	85.86	122.53	81.79	50.35	43.51	27.90	29.04	17.03
100-016	6.76	98.79	98.39	BLQ	62.57	56.47	33.85	26.07	14.97
100-026	BLQ	113.52	99.60	76.71	54.26	44.62	31.42	27.23	16.37
100-036	6.71	111.17	83.49	BLQ	72.78	56.86	28.82	BLQ	21.19
100-046	6.56	90.86	100.65	100.51	BLQ	46.85	36.40	30.32	13.65
100-056	5.70	113.26	105.15	88.86	54.62	55.61	29.11	26.14	15.13
100-066	6.02	86.85	77.29	67.80	78.66	BLQ	BLQ	35.23	18.86
100-076	6.30	132.91	102.17	73.13	71.57	49.83	27.95	23.57	13.56
100-086	6.41	130.71	BLQ	79.32	70.96	46.81	BLQ	23.26	BLQ
100-096	BLQ	130.15	BLQ	97.28	65.25	49.29	32.42	29.28	19.07
101-002	5.05	94.18	84.30	95.45	59.97	40.32	29.65	22.36	16.00
101-012	5.47	119.59	124.23	65.79	74.84	48.91	BLQ	28.71	20.71
101-022	6.27	86.65	92.00	81.61	59.72	40.79	28.18	33.84	20.72
101-032	BLQ	108.20	BLQ	68.02	54.53	40.67	30.95	BLQ	21.09
101-042	6.59	137.42	92.95	69.99	60.14	48.13	33.06	24.55	18.51
101-052	6.33	129.96	121.95	73.46	64.85	47.06	28.55	33.41	19.33
101-062	6.25	96.58	BLQ	83.26	63.06	38.17	31.41	24.39	16.95
101-072	5.98	136.60	117.60	89.13	53.99	49.78	34.88	30.92	13.62
101-082	6.62	94.64	83.78	97.24	70.62	43.45	30.19	21.65	17.44
101-092	BLQ	BLQ	106.55	BLQ	52.49	BLQ	32.00	23.44	19.46
102-008	5.05	110.02	96.83	81.50	65.13	42.74	38.10	29.54	16.67
102-018	6.60	127.17	87.13	104.34	59.35	37.84	38.19	22.91	19.32
102-028	6.42	125.66	96.41	BLQ	64.79	48.29	33.86	22.47	18.76
102-038	6.03	118.38	78.13	70.85	65.69	55.16	33.12	30.50	15.64
102-048	5.51	83.13	120.54	BLQ	75.08	54.97	38.68	25.31	21.11
102-058	BLQ	133.08	126.02	86.46	49.81	54.27	31.65	32.05	13.25
102-068	6.64	137.84	97.88	68.87	56.08	38.87	35.17	27.27	13.95

BLQ = below the limit of quantitation; LLOQ = lower limit of quantitation; CI = confidence interval.

Values BLQ were treated as 0 for predose and postdose.

If > 1/3 of postdose values are BLQ at a time point, then the mean, StD, and %CV are not calculated at that time point.

LLQ was defined as 5.0 ng/mL for analyte GS-XXXX.

For each participant in PK analysis set, intensive serial PK sample collection (samples over 24-hr postdose) was performed at Day 1.

Data Extracted: ddmmmyyyy

Source: .../final/draft1/prog/t-pkpc.sas v9.4 Output file: t-pkpc.pdf 31AUG2026:14:39

Table 15.10.1.1.x: Individual Data and Summary Statistics of Plasma Concentration (ng/mL) at Protocol-Specified Sampling Times
PK Analysis Set

Treatment: Drug A

Participant ID	Pre-dose	Sampling Time (h)							
		1h	2h	4h	8h	12h	16h	18h	24h
102-078	6.58	104.18	82.56	72.78	72.50	41.25	BLQ	32.04	15.42
102-088	6.21	104.00	100.30	88.63	72.81	40.49	35.60	33.87	21.71
102-098	5.77	92.33	124.68	108.50	48.31	54.99	BLQ	34.81	18.61
103-004	5.32	BLQ	85.06	82.21	50.85	41.22	BLQ	35.26	14.18
103-014	6.41	105.38	100.29	70.41	66.97	34.50	BLQ	31.32	14.17
103-024	5.76	97.52	BLQ	95.39	74.56	38.26	40.05	23.39	19.42
103-034	6.10	94.67	81.65	105.87	62.43	BLQ	29.60	21.85	13.82
103-044	5.58	85.75	80.56	71.97	67.97	43.32	39.49	32.71	18.42
103-054	5.16	118.00	90.06	BLQ	BLQ	52.21	30.48	34.27	14.13
103-064	6.83	101.04	104.65	85.18	76.74	BLQ	36.15	28.26	21.58
103-074	BLQ	90.79	102.94	68.43	72.13	56.51	34.62	BLQ	14.47
103-084	5.84	BLQ	109.20	87.26	77.58	55.38	BLQ	30.58	20.03
103-094	5.21	90.13	125.52	96.84	48.28	35.91	39.63	33.30	15.24
104-010	6.97	BLQ	77.07	BLQ	74.28	55.20	39.03	31.98	BLQ
104-020	5.57	88.71	89.30	99.84	66.04	56.67	BLQ	24.64	15.24
104-030	6.69	98.10	111.88	72.42	73.06	36.52	38.49	24.09	18.95
104-040	6.97	107.71	104.09	BLQ	48.48	39.73	35.14	33.06	13.98
104-050	6.20	119.92	122.47	67.77	BLQ	49.99	40.42	26.47	17.50
104-060	5.65	BLQ	BLQ	67.82	75.13	50.18	37.77	34.22	15.42
104-070	6.04	92.87	88.63	81.40	47.83	55.97	27.43	28.60	18.10
104-080	5.91	110.24	103.06	92.95	55.97	41.36	31.79	BLQ	BLQ
104-090	BLQ	91.38	79.29	92.05	66.31	47.43	32.66	32.65	17.10
104-100	6.95	95.75	122.80	72.85	70.40	34.50	37.73	25.13	19.77

BLQ = below the limit of quantitation; LLOQ = lower limit of quantitation; CI = confidence interval.

Values BLQ were treated as 0 for predose and postdose.

If > 1/3 of postdose values are BLQ at a time point, then the mean, StD, and %CV are not calculated at that time point.

LLQ was defined as 5.0 ng/mL for analyte GS-XXXX.

For each participant in PK analysis set, intensive serial PK sample collection (samples over 24-hr postdose) was performed at Day 1.

Data Extracted: ddmmmyyy

Source: .../final/draft1/prog/t-pkpc.sas v9.4 Output file: t-pkpc.pdf 31AUG2026:14:39

Table 15.10.1.1.x: Individual Data and Summary Statistics of Plasma Concentration (ng/mL) at Protocol-Specified Sampling Times
PK Analysis Set

Treatment: Drug A

	Pre-dose	1h	2h	Sampling Time (h)		12h	16h	18h	24h
				4h	8h				
N	50	50	50	50	50	50	50	50	50
Mean	5.24	96.4	88.0	69.6	60.0	42.8	27.6	26.3	16.2
SD	2.20	36.1	35.9	32.7	17.8	14.4	13.6	8.85	4.84
% CV	42.0	37.5	40.8	47.0	29.7	33.7	49.1	33.6	29.9
95% CI(L)	4.62	86.2	77.8	60.3	54.9	38.7	23.8	23.8	14.8
95% CI(U)	5.87	107	98.2	78.9	65.1	46.9	31.5	28.8	17.6
Median	6.03	99.9	97.4	78.0	64.8	45.7	31.9	28.4	17.0
Min	BLQ	BLQ	BLQ	BLQ	BLQ	BLQ	BLQ	BLQ	BLQ
Max	6.97	138	126	109	78.7	56.9	40.4	35.3	21.7
Q1	5.32	90.8	82.6	68.0	54.3	39.7	28.2	23.6	14.2
Q3	6.56	118	107	89.1	72.1	52.2	36.2	32.1	19.3
N (LN-scale)	43	45	44	42	47	46	41	46	47
Geom. Mean	6.07	106	98.9	82.0	63.1	46.0	33.5	28.3	17.0
% GCV	9.54	15.4	15.5	14.8	15.2	15.4	11.8	15.4	15.3
95% GCI(L)	5.89	101	94.4	78.4	60.4	44.0	32.3	27.0	16.3
95% GCI(U)	6.25	111	104	85.9	66.0	48.2	34.7	29.6	17.8
# BLQ	7	5	6	8	3	4	9	4	3
>1/3 BLQ	No	No	No	No	No	No	No	No	No
% BLQ	14.0	10.0	12.0	16.0	6.00	8.00	18.0	8.00	6.00

BLQ = below the limit of quantitation; LLOQ = lower limit of quantitation; CI = confidence interval.

Values BLQ were treated as 0 for predose and postdose.

If > 1/3 of postdose values are BLQ at a time point, then the mean, StD, and %CV are not calculated at that time point.

LLQ was defined as 5.0 ng/mL for analyte GS-XXXX.

For each participant in PK analysis set, intensive serial PK sample collection (samples over 24-hr postdose) was performed at Day 1.

Data Extracted: ddmmyyyy

Source: .../final/draft1/prog/t-pkpc.sas v9.4 Output file: t-pkpc.pdf 31AUG2026:14:39

Table 15.10.1.1.x: Individual Data and Summary Statistics of Plasma Concentration (ng/mL) at Protocol-Specified Sampling Times
PK Analysis Set

Treatment: Drug B

Participant ID	Pre-dose	Sampling Time (h)							
		1h	2h	4h	8h	12h	16h	18h	24h
100-001	5.72	131.85	82.97	67.96	71.94	53.28	33.93	32.54	14.90
100-011	6.50	105.78	76.86	77.64	68.10	38.86	28.56	29.48	18.93
100-021	5.71	102.90	126.46	103.18	60.09	50.09	40.71	35.10	18.32
100-031	BLQ	116.80	100.71	100.26	74.91	36.93	35.34	21.36	20.88
100-041	6.57	124.58	113.94	99.18	49.35	55.56	36.48	34.77	20.97
100-051	6.88	128.99	86.27	85.24	66.47	BLQ	27.53	30.29	16.04
100-061	5.97	92.04	98.39	99.83	55.53	53.36	BLQ	32.35	16.50
100-071	5.77	86.28	92.89	BLQ	50.19	36.47	34.35	21.34	21.34
100-081	5.35	86.69	98.21	BLQ	70.55	40.41	32.64	33.36	21.48
100-091	6.54	125.41	80.20	103.01	60.56	44.42	40.57	22.82	21.22
101-007	BLQ	126.74	121.46	100.34	68.22	49.94	34.36	25.23	15.63
101-017	BLQ	95.40	108.20	75.75	65.96	47.06	39.66	23.39	21.43
101-027	5.02	133.48	109.45	98.13	65.06	43.28	BLQ	30.65	13.33
101-037	6.80	88.47	117.89	81.05	66.49	49.63	32.27	31.50	13.43
101-047	6.32	97.00	92.41	70.02	55.13	45.75	26.62	21.44	16.21
101-057	6.07	89.90	90.96	97.46	59.34	37.00	26.10	23.98	20.05
101-067	5.55	117.48	107.65	84.20	50.01	35.07	26.89	28.08	14.86
101-077	BLQ	BLQ	80.98	77.87	61.12	35.83	31.79	24.76	18.95
101-087	5.88	123.35	124.42	90.21	54.53	47.13	BLQ	35.17	15.71
101-097	BLQ	106.27	BLQ	74.18	68.09	46.20	36.43	29.26	17.21
102-003	BLQ	136.82	127.47	87.04	72.01	43.04	36.35	33.09	19.35
102-013	6.84	120.97	93.73	93.53	78.30	46.75	29.33	27.21	15.41
102-023	6.61	110.19	119.79	102.13	62.17	57.23	34.51	32.76	20.98
102-033	BLQ	106.47	79.65	99.26	59.39	46.95	40.67	28.98	19.22
102-043	6.87	86.19	83.03	107.24	48.52	41.73	38.23	25.13	14.35
102-053	6.22	109.63	115.29	BLQ	60.11	42.11	28.46	25.21	14.78
102-063	BLQ	111.61	78.59	103.16	BLQ	48.60	41.14	BLQ	20.31

BLQ = below the limit of quantitation; LLOQ = lower limit of quantitation; CI = confidence interval.

Values BLQ were treated as 0 for predose and postdose.

If > 1/3 of postdose values are BLQ at a time point, then the mean, StD, and %CV are not calculated at that time point.

LLQ was defined as 5.0 ng/mL for analyte GS-XXXX.

For each participant in PK analysis set, intensive serial PK sample collection (samples over 24-hr postdose) was performed at Day 1.

Data Extracted: ddmmmyyy

Source: .../final/draft1/prog/t-pkpc.sas v9.4 Output file: t-pkpc.pdf 31AUG2026:14:39

Table 15.10.1.1.x: Individual Data and Summary Statistics of Plasma Concentration (ng/mL) at Protocol-Specified Sampling Times
PK Analysis Set

Treatment: Drug B

Participant ID	Pre-dose	Sampling Time (h)							
		1h	2h	4h	8h	12h	16h	18h	24h
102-073	5.38	85.08	BLQ	89.48	48.85	49.66	41.37	27.14	BLQ
102-083	6.27	116.74	78.27	65.46	55.02	34.86	35.31	26.93	19.41
102-093	6.88	112.21	99.30	107.07	60.50	BLQ	38.52	33.32	13.54
103-009	6.01	95.03	81.44	73.46	58.65	40.14	39.03	34.04	18.21
103-019	5.31	116.42	99.10	74.58	BLQ	56.26	27.94	28.95	16.06
103-029	5.70	117.15	100.80	70.66	67.16	39.57	27.57	29.46	19.10
103-039	BLQ	118.17	123.99	65.80	77.54	38.48	31.06	24.13	17.30
103-049	6.18	88.18	99.81	90.14	61.77	56.52	33.53	31.20	20.94
103-059	5.99	101.41	109.29	65.89	77.11	53.11	BLQ	28.23	21.51
103-069	5.85	133.27	BLQ	99.40	66.79	55.92	29.00	31.84	19.71
103-079	6.08	96.40	97.82	72.71	60.46	43.49	40.65	BLQ	16.16
103-089	6.89	133.02	122.43	87.16	75.49	38.47	28.55	31.18	16.52
103-099	5.81	131.81	93.45	71.42	59.87	46.45	36.35	32.58	19.80
104-005	5.24	108.26	99.88	98.24	49.89	39.00	BLQ	34.10	18.42
104-015	5.81	128.83	85.52	67.03	52.27	48.53	BLQ	31.76	17.98
104-025	5.93	106.87	77.95	102.09	65.81	51.74	34.97	27.37	14.32
104-035	6.27	124.88	BLQ	106.34	BLQ	48.83	31.23	23.83	21.28
104-045	6.13	123.83	97.13	BLQ	76.76	40.04	BLQ	28.93	19.68
104-055	5.23	124.19	124.01	86.76	BLQ	46.58	34.17	24.19	13.98
104-065	5.40	119.91	103.35	75.17	53.76	50.03	33.61	24.85	15.78
104-075	5.98	BLQ	81.96	BLQ	54.90	BLQ	27.67	24.11	BLQ
104-085	5.31	100.79	96.11	105.50	59.88	BLQ	29.01	23.01	20.63
104-095	5.23	84.32	103.37	83.56	65.22	49.26	41.15	28.90	20.08

BLQ = below the limit of quantitation; LLOQ = lower limit of quantitation; CI = confidence interval.

Values BLQ were treated as 0 for predose and postdose.

If > 1/3 of postdose values are BLQ at a time point, then the mean, StD, and %CV are not calculated at that time point.

LLQ was defined as 5.0 ng/mL for analyte GS-XXXX.

For each participant in PK analysis set, intensive serial PK sample collection (samples over 24-hr postdose) was performed at Day 1.

Data Extracted: ddmmmyyyy

Source: .../final/draft1/prog/t-pkpc.sas v9.4 Output file: t-pkpc.pdf 31AUG2026:14:39

Table 15.10.1.1.x: Individual Data and Summary Statistics of Plasma Concentration (ng/mL) at Protocol-Specified Sampling Times
PK Analysis Set

Treatment: Drug B

	Pre-dose	1h	2h	Sampling Time (h)		12h	16h	18h	24h
				4h	8h				
N	50	50	50	50	50	50	50	50	50
Mean	4.92	107	91.7	78.7	57.4	42.0	29.1	27.4	17.2
SD	2.38	26.9	31.0	29.5	18.9	13.9	12.7	6.93	4.38
% CV	48.3	25.2	33.9	37.5	33.0	33.2	43.5	25.3	25.4
95% CI(L)	4.25	98.9	82.8	70.3	52.0	38.0	25.5	25.4	16.0
95% CI(U)	5.60	114	100	87.1	62.8	46.0	32.7	29.4	18.5
Median	5.83	111	98.0	86.0	60.5	46.0	33.1	28.9	18.3
Min	BLQ	BLQ	BLQ	BLQ	BLQ	BLQ	BLQ	BLQ	BLQ
Max	6.89	137	127	107	78.3	57.2	41.4	35.2	21.5
Q1	5.24	95.4	82.0	71.4	54.5	38.9	27.7	24.2	15.6
Q3	6.27	124	109	99.4	67.2	49.7	36.4	31.8	20.1
N (LN-scale)	41	48	46	45	46	46	43	48	48
Geom. Mean	5.98	110	98.5	86.4	61.8	45.2	33.5	28.2	17.8
% GCV	8.97	14.7	15.5	16.2	13.7	14.3	14.4	14.8	15.0
95% GCI(L)	5.81	105	94.1	82.3	59.4	43.3	32.0	27.1	17.0
95% GCI(U)	6.15	115	103	90.7	64.4	47.2	35.0	29.5	18.6
# BLQ	9	2	4	5	4	4	7	2	2
>1/3 BLQ	No	No	No	No	No	No	No	No	No
% BLQ	18.0	4.00	8.00	10.0	8.00	8.00	14.0	4.00	4.00

BLQ = below the limit of quantitation; LLOQ = lower limit of quantitation; CI = confidence interval.

Values BLQ were treated as 0 for predose and postdose.

If > 1/3 of postdose values are BLQ at a time point, then the mean, StD, and %CV are not calculated at that time point.

LLQ was defined as 5.0 ng/mL for analyte GS-XXXX.

For each participant in PK analysis set, intensive serial PK sample collection (samples over 24-hr postdose) was performed at Day 1.

Data Extracted: ddmmyyyy

Source: .../final/draft1/prog/t-pkpc.sas v9.4 Output file: t-pkpc.pdf 31AUG2026:14:39