From variational Monte Carlo to superfluid hydrodynamics in microwave-shielded Bose Einstein condensates

Xin-Yuan Gao,[1,*] Kai Yuen Lee,[1,*] and Yangqian Yan[1,2,3,†]

[1] *Department of Physics, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong, China*
[2] *State Key Laboratory of Quantum Information Technologies and Materials, The Chinese University of Hong Kong, Hong Kong SAR, China*
[3] *The Chinese University of Hong Kong Shenzhen Research Institute, 518057 Shenzhen, China*

(Dated: March 6, 2026)

We show that, within the manifold where the Jastrow pair factor is frozen to its ground-state form, time-dependent variational Monte Carlo (tVMC) based on a Jastrow–Feenberg wavefunction maps exactly onto irrotational superfluid hydrodynamics. The only microscopic input is an equation of state (EOS) extracted from a single variational Monte Carlo calculation of the uniform system, which yields a density-dependent chemical potential $\mu(\rho)$ for a generalized Gross–Pitaevskii equation (GPE). For a trapped microwave-shielded dipolar Bose gas, this reduced description reproduces the full tVMC breathing dynamics for $N = 8$–128 with below-1% frequency error and $\sim 2\%$ amplitude error, while lowering the computational cost by more than an order of magnitude. The framework turns correlated tVMC into a practical predictive tool for collective dynamics in shielded molecular gases.

Introduction.—Microwave-shielded polar molecules have emerged as a promising platform for quantum many-body physics, with recent experiments achieving Bose–Einstein condensation [1], observing shielded collisions [2, 3], and probing dipolar many-body effects [4–6]. A central theoretical challenge is to predict nonequilibrium collective dynamics—breathing oscillations, quenches, and expansion—from the microscopic interaction, which combines long-range dipolar attraction ($\sim C_3/r^3$) with short-range repulsive shielding ($\sim C_6/r^6$).

Mean-field Gross–Pitaevskii (GP) theory is computationally efficient but reduces this interaction to a contact nonlinearity, thereby missing the anisotropic and density-dependent correlations generated by the shielding mechanism. At the opposite end, time-dependent variational Monte Carlo (tVMC) [7, 8] can treat realistic interactions within a correlated many-body wavefunction, but its stochastic sampling at every time step severely limits accessible system sizes and evolution times.

This Letter introduces a general theoretical framework that retains the microscopic content of tVMC while recovering the efficiency of hydrodynamics. While demonstrated here for shielded molecular gases, this approach provides a pathway for describing collective dynamics in other strongly correlated fluids, such as quantum droplets, supersolids, and strongly interacting Rydberg gases, where standard mean-field theories often break down. Our central result is conditional but exact: once the two-body Jastrow factor u_2 is frozen, the time-dependent variational principle (TDVP) maps exactly onto irrotational superfluid hydrodynamics. We first verify from full tVMC that u_2 remains nearly stationary during the quench, then extract the uniform-system EOS from a one-time VMC calculation, and finally solve a generalized GPE with this EOS. The resulting dynamics reproduces the full tVMC benchmarks for $N = 8$ and $N = 128$ and extends straightforwardly to experimentally relevant large-N regimes beyond direct tVMC.

Model and variational dynamics.—We consider N bosons in a cylindrically symmetric harmonic trap with Hamiltonian

$$\hat{H}(t) = \sum_{i=1}^{N}\big[-\tfrac{1}{2}\nabla_i^2 + V_{\rm tr}(\mathbf{r}_i, t)\big] + \sum_{i<j} v(\mathbf{r}_{ij}), \qquad (1)$$

where $V_{\rm tr} = \frac{1}{2}[\omega_\rho(t)^2\rho^2 + \omega_z^2 z^2]$ (natural units $\hbar = m = 1$) and the two-body interaction models microwave-shielded dipolar molecules:

$$v(\mathbf{r}) = \frac{C_3}{r^3}(3\cos^2\theta - 1) + \frac{C_6}{r^6}(\sin^4\theta + w_{12}\sin^2\theta\cos^2\theta). \quad (2)$$

This effective potential, derived from Floquet scattering theory for dual-microwave-dressed molecules [9], is parametrized by coefficients C_3, C_6, and w_{12} that are determined by the microwave field configuration. In a realistic experiment these are fixed by the dressing scheme; here, for demonstration, we choose $C_3 = 10^{-3}$, $C_6 = 10^{-6}$, and $w_{12} = 2.0$ in oscillator units. We study a quench $\omega_\rho : 1 \to 2$ at $t = 0$ that excites radial breathing modes.

The many-body wavefunction is a truncated Jastrow–Feenberg [10, 11] ansatz:

$$\Psi(\mathbf{X}, t) = \exp\Big[\sum_i u_1(\mathbf{r}_i, t) + \sum_{i<j} u_2(r_{ij}, \hat{\mathbf{r}}_{ij})\Big], \qquad (3)$$

where $u_1(\mathbf{r}, t)$ is a complex, time-dependent one-body field and u_2 encodes two-body correlations. In cylindrical geometry we use a separable parametrization for u_1,



FIG. 1. (a) Leading-order HNC pair correlation $e^{2u_2(r,\theta=0)}$ for the N=8 ground state, showing a correlation hole at short range due to the shielding repulsion. (b) Breathing dynamics after the quench $\omega_\rho : 1 \to 2$ for N=128: $\langle\rho^2\rangle(t)$ and $\langle z^2\rangle(t)$ from full tVMC (red) and frozen-u_2 tVMC (blue dashed), validating the frozen-u_2 approximation.

while u_2 is expanded in partial waves as

$$u_2(r,\cos\theta) = u_{\rm cusp}(r) + \sum_{\ell=0,2,4} f_\ell(r)\, P_\ell(\cos\theta), \tag{4}$$

where $u_{\rm cusp}(r)$ captures the short-range scattering boundary condition and each f_ℓ is represented by Gaussian basis functions [12]. For the parameters studied here, the $\ell = 4$ channel is negligible and the $\ell = 2$ anisotropy is weak, so the pair correlation is close to isotropic.

Figure 1 shows the quality of the optimized pair correlations. The leading-order hypernetted-chain (HNC) approximation to the pair distribution function, $g_2^{\rm HNC}(r,\theta) = e^{2u_2(r,\cos\theta)}$ [13], reveals a correlation hole at short range [Fig. 1(a)], reflecting the interplay between dipolar attraction along $\theta = 0$ and the shielding repulsion. The pair correlation is nearly isotropic, consistent with the subdominant role of f_2.

The TDVP [14, 15] projects the Schrödinger equation onto the variational tangent space. Defining logarithmic derivatives $O_\alpha = \partial_{\theta_\alpha} \ln\Psi$ and the local energy $E_L = (\hat{H}\Psi)/\Psi$, the equations of motion read [7]

$$\sum_\beta S_{\alpha\beta}\,\dot\theta_\beta = -i\,F_\alpha, \tag{5}$$

where the overlap matrix and force vector are

$$S_{\alpha\beta} = \langle \Delta O_\alpha^* \,\Delta O_\beta\rangle_t, \tag{6}$$
$$F_\alpha = \langle \Delta O_\alpha^* \,\Delta E_L\rangle_t, \tag{7}$$

with $\Delta A = A - \langle A\rangle_t$ denoting connected fluctuations. Full Monte Carlo and convergence details are given in Ref. [12].

Frozen-u_2 reduction.—The key numerical observation is that during the quench the optimized two-body parameters change by less than 1% of their ground-state values, while the one-body field changes by order unity. Figure 1(b) already shows that a frozen-u_2 tVMC evolution tracks the full tVMC dynamics, and Fig. 2 visualizes the near-stationarity of the pair correlation itself. This motivates restricting the variational manifold to a fixed u_2.

When the variational degree of freedom is the continuous field $u_1(\mathbf{r},t)$, the TDVP logarithmic derivative becomes the microscopic density operator:

$$O(\mathbf{r};\mathbf{X}) = \frac{\delta \ln\Psi}{\delta u_1(\mathbf{r})} = \sum_i \delta^{(3)}(\mathbf{r}-\mathbf{r}_i) = \hat\rho(\mathbf{r}). \tag{8}$$

Because u_2 is real and frozen, the many-body phase is purely one-body and irrotational: $\mathbf{v} = \nabla(\mathrm{Im}\, u_1)$. No backflow terms arise.

Decomposing $u_1 = a + i\theta$ (real amplitude and phase), the TDVP action reduces to the standard hydrodynamic form [12]. Within this frozen-u_2 manifold the reduction is exact and yields the continuity and Euler equations:

$$\partial_t\rho + \nabla\cdot(\rho\mathbf{v}) = 0, \tag{9}$$
$$\partial_t\mathbf{v} + \nabla\Big(\tfrac{1}{2}v^2 + V_{\rm tr} + \mu_{u_2}[\rho]\Big) = 0, \tag{10}$$

where $\mu_{u_2}[\rho] = \delta E_{\rm int}[\rho;u_2]/\delta\rho$ is the effective chemical potential determined by the frozen correlations.

To obtain a closed local theory, we further invoke the local-density approximation (LDA). The internal energy then takes the form $E_{\rm int} = \int \rho\,\varepsilon(\rho)\,d^3r$, where $\varepsilon(\rho)$ is the energy per particle of the *uniform* system at density ρ with the same frozen u_2. The chemical potential becomes a local function:

$$\mu(\rho) = \frac{d[\rho\varepsilon(\rho)]}{d\rho} = \varepsilon + \rho\varepsilon'. \tag{11}$$

This is structurally identical to GP theory, except that the usual contact coupling gn is replaced by $\mu(\rho)$, which encodes the full effect of the realistic, anisotropic interaction through the frozen correlations.

The near-stationarity of u_2 during the quench is visualized directly in Fig. 2, which shows the change in the leading-order HNC pair correlation Δe^{2u_2} at four time snapshots. The variation remains below ~ 0.03 throughout (compared to peak values ~ 1), confirming that the pair correlations are effectively frozen.

Equation of state and generalized GPE.—We extract $\varepsilon(\rho)$ from a one-time VMC calculation on the uniform system (periodic boundary conditions) with u_2 frozen at its trapped ground-state values. Scanning 20–30 density values logarithmically from $\rho = 0.05$ to $\rho_{\max}$ (3.0 for $N = 8$; 15.0 for $N = 128$) with 8192 walkers, we fit the results to

$$\varepsilon(\rho) = a_1\rho + a_2\rho^2 + \cdots , \tag{12}$$

yielding the chemical potential $\mu(\rho) = 2a_1\rho + 3a_2\rho^2 + \cdots$.

This expansion gives a compact physical interpretation of the EOS. Relative to the standard GPE nonlinearity $\mu = g\rho$, the leading term $2a_1\rho$ corresponds to a *renormalized two-body coupling* $g_{\rm eff} = 2a_1$, which absorbs the effect of the frozen pair correlations into an effective contact interaction. The next-order term $3a_2\rho^2$, however, has the structure of an *effective three-body interaction* with coupling strength $g_3 = 3a_2$. This is not a genuine microscopic three-body force; rather, it emerges from integrating out the density dependence of the two-body correlations encoded in u_2. In density-functional language, it is the leading beyond-mean-field correction to the interaction energy [16].

For the shielded dipolar interaction studied here, we find $g_3/g_{\rm eff}^2 \sim 0.1$–$0.2$, indicating that the three-body term provides a 10–20% correction at the equilibrium density. For $N = 8$, the data require a weak cubic correction, whereas for $N = 128$ a quadratic fit with a_1=0.111 and a_2=0.001 already describes the relevant density range and remains stable ($d\mu/d\rho > 0$). The EOS is therefore close to linear at high density, reflecting the long-range character of the shielded dipolar interaction.

The VMC EOS for two interaction strengths is shown in Fig. 3. The leading coefficient a_1 scales linearly with C_3, while a_2 captures the density-dependent beyond-mean-field corrections.

The resulting generalized GPE,

$$i\partial_t\psi = \Big[-\tfrac{1}{2}\nabla^2 + V_{\rm tr} + g_{\rm eff}|\psi|^2 + g_3|\psi|^4 \Big]\psi, \tag{13}$$

is therefore a *two-plus-three-body* GPE whose couplings are fixed microscopically by VMC rather than by phenomenology. This provides a first-principles realization of the extended GPE forms discussed in the contexts of quantum droplets [17] and dipolar condensates [18, 19].

GPE solver.—We solve the generalized GPE

$$i\partial_t\psi = \Big[-\tfrac{1}{2}\nabla^2 + V_{\rm tr} + \mu(|\psi|^2) \Big]\psi, \tag{14}$$

where $\psi = \sqrt{\rho}\, e^{i\theta}$ and the Laplacian provides both quantum pressure and flow kinetic energy. The equation is discretized on a cylindrical (r, z) grid exploiting azimuthal symmetry, with a fourth-order finite-difference Laplacian (including L'Hôpital's rule at $r = 0$) and fourth-order Runge–Kutta time integration under a dispersive CFL condition $\Delta t \le C\Delta x^2/\pi$ ($C = 0.3$) [12].

The initial state is obtained by imaginary-time propagation to self-consistency: $\psi \to \psi - d\tau\, \hat{H}\psi$, normalized to N particles at each step, until the chemical potential converges to $\delta\mu < 10^{-8}$.

Validation against tVMC.—We benchmark the generalized GPE against frozen-u_2 tVMC at both weak and moderate interaction strength. For $N = 8$, the GPE ground state reproduces the tVMC moments to 0.7% and the energy to 0.15% (Table S1 [12]). The breathing frequency is $\omega_{\rm br} = 4.19 \approx 2\omega_\rho$ in both methods, and the oscillation amplitudes of $\langle\rho^2\rangle$ and $\langle z^2\rangle$ agree to within 1%. The quantum-pressure term is essential in this regime because the interaction energy is only $\sim 3.5\%$ of the total.

For $N = 128$, where the interaction energy reaches $\sim 21\%$ of the total, the same EOS-based GPE remains accurate. The ground-state moments agree with tVMC to 0.35% for $\langle\rho^2\rangle$ and 1.05% for $\langle z^2\rangle$, while the total energy differs by 2.8%, consistent with a residual LDA error at higher density. More importantly, the dynamical benchmark in Fig. 4(a) is essentially unchanged: $\omega_{\rm br} = 4.19$ in both approaches and the $\langle\rho^2\rangle(t)$ amplitude agrees to $\sim 2\%$.

The computational gain is substantial. For $N = 128$, the GPE evolution including imaginary-time state preparation takes ~ 70 minutes on a single CPU core, compared with ~ 6 hours on a modern GPU for the full tVMC. The reduction is larger still in total floating-point cost, since the stochastic tVMC evolution requires fresh Monte Carlo sampling at every Runge–Kutta stage.

Large-N prediction.—Having benchmarked the reduction, we use it in a regime inaccessible to direct tVMC. For a weaker interaction, $C_3 = 10^{-4}$, the VMC-derived EOS predicts sustained breathing oscillations for $N = 10{,}000$ over at least ten trap periods [Fig. 4(b)]. This illustrates the practical value of the approach: once the EOS is known, the many-body quench problem is reduced to a deterministic hydrodynamic evolution.

Discussion.—The success of the frozen-u_2 hydrodynamic reduction rests on a physical separation of scales: short-range pair correlations, governed by the interaction potential on lengths $r \lesssim a_{\rm ho}$, equilibrate much faster than the global density redistribution driven by the trap quench. This is manifested in the tVMC as $< 1\%$ variation of the u_2 parameters during the quench, while u_1 changes by order unity.

Two features distinguish the present framework from standard GP theory. First, the EOS is microscopically derived from the correlated many-body wavefunction



FIG. 2. Change in the leading-order HNC pair correlation, $\Delta e^{2u_2} = e^{2u_2(t)} - e^{2u_2(0)}$, during breathing dynamics for N=128. The pair correlations remain nearly frozen ($|\Delta e^{2u_2}| \lesssim 0.03$) while the one-body density undergoes order-unity changes.



FIG. 3. Chemical potential $\mu(\rho) = d(\rho\varepsilon)/d\rho$ from uniform-system VMC for C_3=10^{-3} (circles, black fit) and C_3=10^{-4} (squares, red fit). Data points are obtained by numerical differentiation of the VMC energy; lines show polynomial fits used in the GPE.

rather than inferred from a scattering-length reduction. Second, the hydrodynamic equation is not postulated: it follows directly from the TDVP once the dynamics is projected onto the frozen-u_2 manifold. In this sense, the approximation is transparent and diagnosable: its validity can be checked a posteriori by monitoring the time dependence of u_2.

The same construction also clarifies the origin of higher-order nonlinearities. The effective three-body term $g_3|\psi|^4$ in Eq. (13) is not inserted phenomenologically; it emerges from the density dependence of correlated two-body physics. This makes the method a useful nonperturbative route to extended GPEs for dipolar and molecular systems, where beyond-mean-field effects are often important but difficult to parameterize reliably.

The main quantitative limitation visible here is the $\sim 3\%$ energy offset at $N = 128$, which points to gradient corrections beyond the LDA [20]. More generally, the present reduction should hold when pair correlations remain slaved to the evolving density and the flow stays irrotational. It can then be extended naturally to other collective modes, quench protocols, and trap geometries relevant to current molecular-gas experiments [1, 21].

Conclusion.—We have demonstrated that correlated many-body dynamics from tVMC maps exactly onto superfluid hydrodynamics within the frozen-u_2 manifold, with the required equation of state obtained from a single uniform-system VMC calculation. The resulting generalized GPE reproduces the breathing dynamics of microwave-shielded dipolar bosons from $N = 8$ to $N = 128$ with below-1% frequency error and $\sim$ 2% amplitude error, while providing access to large-N regimes beyond direct tVMC. Beyond establishing a practical bridge between microscopic physics and predictive hydrodynamics for molecular gases, the "frozen-Jastrow" hydrodynamic concept introduced here establishes a powerful methodology. It has the potential to enable first-principles modeling of non-equilibrium dynamics in a wide array of quantum fluids governed by complex long- and short-range correlations, including quantum droplets, supersolids, and Rydberg atomic gases.

Computations were performed using JAX [22] for GPU-accelerated VMC. We would like to acknowledge financial support from the National Natural Science Foundation of China under Grant No. 124B2074 and No. 12204395, Hong Kong RGC Early Career Scheme (Grant No. 24308323) and Collaborative Research Fund (Grant No. C4050-23GF), the Space Application System of China Manned Space Program, Guangdong Provincial Quantum Science Strategic Initiative GDZX2404004, and CUHK Direct Grant No. 4053731.



FIG. 4. GPE dynamics from the VMC-derived EOS. (a) N=128 validation: $\langle\rho^2\rangle(t)$ and $\langle z^2\rangle(t)$ from frozen-u_2 tVMC (red) and GPE (blue), showing identical breathing frequencies within numerical resolution. (b) N=10,000 prediction with the weak-interaction EOS (C_3=10^{-4}), extrapolating to experimentally relevant system sizes inaccessible to direct tVMC.

* These authors contributed equally.
† yqyan@cuhk.edu.hk

[1] N. Bigagli, W. Yuan, S. Zhang, B. Ghosh, S. Kolkowitz, T. Tiberi, N. Bhatt, I. Stevenson, J. Ye, and W. Ketterle, Observation of Bose–Einstein condensation of dipolar molecules, Nature **631**, 289 (2024).
[2] L. Anderegg, S. Burchesky, Y. Bao, S. S. Yu, T. Karman, J. M. Doyle, and K.-K. Ni, Observation of microwave shielding of ultracold molecules, Science **373**, 779 (2021).
[3] D. I. Schuster, N. Li, S. Kotochigova, and E. Timmermans, Collisional shielding of ultracold molecules, Nat. Phys. **19**, 1450 (2023).
[4] T. Karman and J. M. Hutson, Microwave shielding of ultracold polar molecules, Phys. Rev. Lett. **121**, 163401 (2018).
[5] T. Karman and J. M. Hutson, Microwave shielding of ultracold polar molecules with imperfectly circular polarization, Phys. Rev. A **100**, 052704 (2019).
[6] X.-Y. Chen, S. Biber, A. Backes, P. Bitsch, R. Gao, I. Bloch, and F. Schäfer, Ultracold field-linked tetratomic molecules, Nature **626**, 283 (2024).
[7] G. Carleo, F. Becca, M. Schiró, and M. Fabrizio, Localization and glassy dynamics of many-body quantum systems, Sci. Rep. **2**, 243 (2012).
[8] G. Carleo, F. Becca, L. Sanchez-Palencia, S. Sorella, and M. Fabrizio, Light-cone effect and supersonic correlations in one- and two-dimensional bosonic superfluids, Phys. Rev. A **89**, 031602 (2014).
[9] F. Deng, X. Hu, W.-J. Jin, S. Yi, and T. Shi, Two- and many-body physics of ultracold molecules dressed by dual microwave fields, Nat. Commun. **16**, 5635 (2025).
[10] R. Jastrow, Many-body problem with strong forces, Phys. Rev. **98**, 1479 (1955).
[11] E. Feenberg, *Theory of Quantum Fluids* (Academic Press, New York, 1969).
[12] See Supplemental Material for details on the basis-function expansion, cusp treatment, TDVP derivation of the hydrodynamic equations, EOS fitting procedure, and numerical convergence tests.
[13] J.-P. Hansen and I. R. McDonald, *Theory of Simple Liquids*, 3rd ed. (Academic Press, 2006).
[14] P. A. M. Dirac, Note on exchange phenomena in the thomas atom, Math. Proc. Cambridge Philos. Soc. **26**, 376 (1930).
[15] A. D. McLachlan, A variational solution of the time-dependent Schrödinger equation, Mol. Phys. **8**, 39 (1964).
[16] L. P. Pitaevskii and S. Stringari, *Bose-Einstein Condensation and Superfluidity*, 2nd ed. (Oxford University Press, 2016).
[17] D. S. Petrov, Quantum mechanical stabilization of a collapsing Bose-Bose mixture, Phys. Rev. Lett. **115**, 155302 (2015).
[18] L. Chomaz, I. Ferrier-Barbut, F. Ferlaino, B. Laburthe-Tolra, B. L. Lev, and T. Pfau, Dipolar physics: a review of experiments with magnetic quantum gases, Rep. Prog. Phys. **86**, 026401 (2023).
[19] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein, and T. Pfau, The physics of dipolar bosonic quantum gases, Rep. Prog. Phys. **72**, 126401 (2009).
[20] F. Ancilotto, M. Barranco, and M. Pi, Density functional theory of helium droplets and clusters, Int. Rev. Phys. Chem. **36**, 621 (2017).
[21] J. Lin, J. Rosenberg, X. Zhu, L. Li, C. He, N. Bigagli, J. Devlin, and W. Ketterle, Microwave shielding of bosonic NaRb molecules, Phys. Rev. X **14**, 011033 (2024).
[22] J. Bradbury, R. Frostig, P. Hawkins, M. J. Johnson, C. Leary, D. Maclaurin, G. Necula, A. Paszke, J. VanderPlas, S. Wanderman-Milne, and Q. Zhang, JAX: composable transformations of Python+NumPy programs (2018).