Converting vertisol retention curves with a shrinking bulk density: two examples

Laikipia (Kenya) and Tomejil (Andalusia, Spain) · py_HB_texture · R. Muñoz-Carpena, 31 October 2026

Why

Retention curves are often measured as gravimetric water and made volumetric with one bulk density per sample. In a vertisol that misrepresents the soil: swelling clays shrink as they dry, so their bulk density rises along the curve, and one wet-state density understates the water held at the dry end. This note applies the same variable-density correction to two vertisols from two regions, shows it step by step for one sample of each, compares how much it changes the curves, and reports its effect on the texture and Ks the tool predicts from them.

The two examples

Soil	Data	Clay	One density used	Worked example
Laikipia, Kenya (vertic clays; Willard et al. 2026, HydroShare)	70 cores, gravimetric as published (the 16 samples flagged for incomplete dispersion left out)	median 55 %	the core's KALRO density (median 1.07 g/cm³)	WL_3LT07414: clay 58 %, sand 28 %, silt 14 %, density 1.07 g/cm³
Tomejil, Carmona, Spain (Chromic Haploxerert; Vanderlinden et al. 2017, 2021)	54 rings, 0–5 cm, gravimetric; no density per ring	56 %	the tillage treatment's mean: 1.085 (conventional) or 1.125 g/cm³ (direct drill; Ordóñez Fernández et al. 2007)	DD30 (direct drill), the median ring at pF 4.2

Steps, at each measured suction

1. Gravimetric water: $w = \theta / BD_1$, with BD_1 the one density above. Both soils' curves are already gravimetric as published, so w is taken directly; the standard mass–volume relation, with water density 1 g/cm³ (Hillel 1998), links it to the one-density curve $\theta_1 = w \cdot BD_1$.
2. Specific volume under normal shrinkage, where the soil loses volume equal to the water it loses: $v = 1/\rho s + w$, with ρs = 2.65 g/cm³, the conventional mineral particle density (Blake and Hartge 1986) (cm³ of soil per g of solids). This is the saturation line of the shrinkage curve, the "normal" phase of Haines (1923) and Stirk (1954), in the specific-volume form of McGarry and Malafant (1987); Bronswijk (1988) used it to model swelling clays.
3. Bulk density: $BD = \min(cap, 1/v)$. The cap stands for the density at which shrinkage stops. For Laikipia it is 1.6 g/cm³ (shrinkage limit w = 1/1.6 − 1/2.65 = 0.248), the value of the published volumetric series (Willard et al. 2026), which is also the dry-end density reported for the Tomejil vertisol (Vanderlinden et al. 2021). For Tomejil it is 1.75 g/cm³ (w = 0.194), the median oven-dry clod density of US Vertisols like these two (KSSL; clay 45–65 %, clod density at 33 kPa below 1.25 g/cm³; 1.54–1.90 for the middle 80 %; Soil Survey Staff 2014a and the KSSL database). Both caps give the same texture results (see *How general is the cap?* below). Where the soil is wetter than the core was dense, 1/v is below BD_1: the core had swollen, and the conversion lowers θ. This step is a simplification: measured shrinkage curves have structural, normal, residual and zero phases (Mitchell 1992; Groenevelt and Grant 2001; Peng and Horn 2005; Cornelis et al. 2006), and in the residual phase air enters and shrinkage slows. Here normal shrinkage runs up to the cap and then stops, with no residual phase, and the cap is not measured on each soil.

4. New volumetric water: `θ_new = w · BD`, the mass–volume relation of step 1 with the shrinking density. The one-density curve is `θ₁ = w · BD₁`.

Laikipia, sample WL_3LT07414 (BD_1 = 1.07 g/cm³): swollen at the wet end, at the 1.6 cap by 1500 kPa


h (kPa)	pF	θ one density	w (g/g)	v (cm³/g)	BD(w) (g/cm³)	θ new	new / one
0.00	–	0.703	0.657	1.034	0.967	0.635	0.90
0.98	1.00	0.630	0.589	0.966	1.035	0.610	0.97
5.51	1.75	0.600	0.560	0.938	1.066	0.598	1.00
7.79	1.90	0.544	0.509	0.886	1.128	0.574	1.05
9.81	2.00	0.496	0.464	0.841	1.189	0.551	1.11
13.85	2.15	0.479	0.448	0.825	1.212	0.543	1.13
19.57	2.30	0.430	0.402	0.779	1.283	0.516	1.20
21.95	2.35	0.365	0.341	0.719	1.391	0.475	1.30
31.01	2.50	0.297	0.278	0.655	1.526	0.424	1.43
49.15	2.70	0.290	0.271	0.648	1.543	0.418	1.44
69.43	2.85	0.285	0.266	0.644	1.553	0.414	1.45
491.50	3.70	0.277	0.259	0.636	1.573	0.407	1.47
1554.25	4.20	0.251	0.234	0.612	1.600	0.375	1.50

The values equal the published volumetric series for this core. At saturation θ_1 (0.703) is above the porosity of the core's own density (0.60): the core had swollen, so the shrinking density there (0.97 g/cm³) is below BD_1 and θ falls; from pF 1.75 on, θ rises.

Tomejil, ring DD30 (BD_1 = 1.125 g/cm³), points up to 1500 kPa

h (kPa)	pF	θ one density	w (g/g)	v (cm³/g)	BD(w) (g/cm³)	θ new	new / one
0.10	0.00	0.545	0.484	0.862	1.160	0.562	1.03
0.25	0.40	0.538	0.478	0.855	1.169	0.559	1.04
0.98	1.00	0.530	0.471	0.848	1.179	0.555	1.05
3.10	1.50	0.472	0.419	0.797	1.255	0.526	1.12
6.19	1.80	0.435	0.387	0.764	1.308	0.506	1.16
9.81	2.00	0.415	0.369	0.746	1.341	0.494	1.19
19.61	2.30	0.387	0.344	0.721	1.387	0.477	1.23
49.03	2.70	0.351	0.312	0.690	1.450	0.453	1.29
254.97	3.41	0.323	0.287	0.664	1.505	0.432	1.34
441.30	3.65	0.281	0.250	0.627	1.594	0.398	1.42
500.14	3.71	0.270	0.240	0.617	1.620	0.389	1.44
578.59	3.77	0.257	0.228	0.606	1.651	0.377	1.47
637.43	3.81	0.234	0.208	0.585	1.708	0.355	1.52
1206.22	4.09	0.210	0.186	0.564	1.750	0.326	1.56
1461.19	4.17	0.198	0.176	0.554	1.750	0.308	1.56

DD30's 12 WP4 readings beyond 1500 kPa (to 175 MPa) all sit at the cap, ratio 1.56; the tool fits curves up to 1500 kPa.



Laikipia (top) and Tomejil (bottom): the one-density and shrinking-density curves (a, d) and the density each uses (b, e), cap 1.6 g/cm³ for Laikipia and 1.75 for Tomejil; (c) how much the conversion changes θ across all 70 Laikipia samples and all 54 Tomejil rings.

How the two compare

	Laikipia (70 samples, cap 1.6)	Tomejil (54 rings, cap 1.75)
One density	the core's KALRO density, median 1.07 g/cm³	treatment mean, 1.085 / 1.125 g/cm³
Shrinking density at the wettest point (median)	1.01 g/cm³	1.16 g/cm³
Shrinking density at 1500 kPa (median)	1.60 g/cm³	1.75 g/cm³
Samples at the cap by 1500 kPa	68 of 68 with a 1500 kPa point	36 of 54
θ new / θ one density, wettest point	0.99 [0.94–1.01]	1.05 [1.01–1.08]
… at pF 2 (98 cm)	1.26 [1.17–1.34]	1.25 [1.20–1.29]
… at pF 2.5 (33 kPa)	1.39 [1.26–1.50]	1.32 [1.27–1.35]
… at pF 3.7 (490 kPa)	1.48 [1.35–1.60]	1.50 [1.46–1.54]
… at pF 4.2 (1500 kPa)	1.50 [1.37–1.62]	1.58 [1.56–1.61]
Texture as a new source, one density: clays read as clay; predicted vs measured clay	57.6 %; 40.0 % vs 54.1 %	85.2 %; 39.9 % vs 55.6 %
… shrinking density	100 %; 46.1 % vs 54.1 %	100 %; 53.4 % vs 55.6 %
Water left at 330 cm, as a share of saturation (reference clays: 82 %), one → shrinking density	50 % → 73 %	65 % → 81 %
Ks, kNN median, one → shrinking density	4.8 → 0.85 cm/h	2.3 → 0.093 cm/h

	Laikipia (70 samples, cap 1.6)	Tomejil (54 rings, cap 1.75)
Ks, physical estimate, one → shrinking density	14.4 → 3.19 cm/h	6.3 → 1.72 cm/h
kNN and physical Ks within a factor of 10 of each other, one → shrinking density	97 % → 79 %	93 % → 17 %
Within a factor of 10 of the SATURO infiltrometer (Laikipia, 26 samples, median 14.0 cm/h), kNN; physical, one → shrinking density	92 % → 54 %; 96 % → 92 %	not measured

Medians, with the 25–75 % range in brackets. Laikipia's wettest point is saturation (h = 0), Tomejil's h = 1 cm. Texture and Ks: each soil predicted as a new source (its own table removed from the reference) with the current tool (`test_caps.py`). The shrinking-density curves are the ones the tool's reference holds (for Laikipia, the published volumetric series), so these rows match the tool's own validation of both soils.

How general is the cap?

The two caps used here, 1.6 g/cm³ for Laikipia (as published) and 1.75 g/cm³ for Tomejil, can be checked against a general source, the US National Cooperative Soil Survey Soil Characterization Database (KSSL): every natural clod is weighed and measured at 33 kPa and again oven-dry (Saran-coated clod method; Grossman et al. 1968; Soil Survey Staff 2014a), the measurement behind the coefficient of linear extensibility (COLE) used to recognize Vertisols (Soil Survey Staff 2014b). In the model of step 3, the oven-dry clod density is the cap: the density a clod reaches once it has stopped shrinking. Drawn from the database's horizons with both densities:

KSSL horizons	Horizons (pedons)	Clod density at 33 kPa, median	Oven-dry clod density, median [10–90 %]	COLE, median
All Vertisols	4,188 (687)	1.30	1.80 [1.58–1.96]	0.115
Vertisols, clay 45–65 %	2,444 (581)	1.29	1.82 [1.62–1.97]	0.119
Vertisols, clay 45–65 %, clod density at 33 kPa below 1.25 (like Laikipia and Tomejil)	828 (324)	1.18	**1.75 [1.54–1.90]**	0.143
Vertic subgroups of other orders, clay 45–65 %	1,126 (415)	1.30	1.77 [1.50–1.94]	0.102
Any order, clay 45–65 %, COLE ≥ 0.09	4,367 (1,816)	1.27	1.81 [1.57–1.96]	0.121

Densities in g/cm³; COLE computed as $(\text{oven-dry} / 33\ \text{kPa density})^{1/3} - 1$. Of the Vertisols with clay 45–65 %, 92 % dry denser than 1.6 g/cm³ and 80 % denser than 1.7. Across them the oven-dry density rises with the 33 kPa density (oven-dry ≈ 0.615 × BD_{33} + 1.011, r = 0.50), which gives a per-sample cap: 1.67 for WL_3LT07414 (BD_1 1.07), 1.70 and 1.68 for the Tomejil treatments.

1.6 and 1.75 bracket the likely value. The oven-dry density is an upper bound for where normal shrinkage ends, since real clods keep shrinking slowly through the residual phase before they are fully dry (the long normal phase of Texas Vertisols in Yule and Ritchie 1980 is one measured example); the model of step 3 has no residual phase, so a 1.75 cap would push the dry end slightly too high, and 1.6 errs the other way. The texture result does not depend on which is used (each soil predicted as a new source, its own table removed from the reference; `test_caps.py`):

Shrinkage cap	Laikipia: exact class	Laikipia: clays read as clay (n = 66)	Laikipia: predicted clay (54.1 %)	Tomejil: rings read as clay (n = 54)	Tomejil: predicted clay (55.6 %)
none (one density)	54.3 %	57.6 %	40.0 %	85.2 %	39.9 %
1.4	57.1 %	60.6 %	40.0 %	96.3 %	49.2 %
1.6 (Laikipia, as published; Tomejil's dry density)	94.3 %	100 %	46.1 %	100 %	53.4 %
1.75 (KSSL median; Tomejil above)	94.3 %	100 %	47.3 %	100 %	53.4 %
per sample (0.615 × BD_1 + 1.011)	92.9 %	98.5 %	46.8 %	100 %	53.6 %

Raising the cap from 1.6 to 1.75 changes only the dry end, where the curves are already flat: single samples' predicted clay moves by up to 8.4 points (Laikipia) and 3.9 (Tomejil), the means by 1.2 points or less, one Laikipia sample changes class and every Tomejil ring keeps its class. A cap of 1.4, below the shrinkage limit of most vertisols, loses most of the gain for Laikipia (57 % exact) and part of it for Tomejil.

What the two examples show

- **The correction behaves alike through the pressure-plate range.** θ rises by about a quarter at pF 2, by a third or more at 33 kPa, and by about half at 1500 kPa (50 % for Laikipia, 58 % for Tomejil, median).
- **They differ at the wet end.** Near saturation the Laikipia cores had swollen beyond their ring density (θ_1 above the porosity of BD_1 in 42 of the 70 samples), so the shrinking density there is below BD_1 and θ falls slightly (median ratio 0.99, 0.90 for WL_3LT07414). Tomejil's rings sit on the shrinkage line, and θ rises by 5 %.
- **Both reach their cap at the dry end.** Every Laikipia sample with a 1500 kPa point reaches 1.6 (WL_3LT07414: w = 0.234 at 1500 kPa, below the limit of 0.248), and 36 of the 54 Tomejil rings reach 1.75.
- **Texture improves in both.** With one density both soils read too coarse and seemed to lose water steeply just below saturation (50 % and 65 % of their saturated water left at 330 cm, against 82 % for the reference clays). With the shrinking density they hold 73 % and 81 %, and every clay in both soils is read as clay. The predicted clay matches the measured mean for Tomejil (53 % against 56 %) but stays below it for Laikipia (46 % against 54 %). The result hardly depends on whether the cap is 1.6 or 1.75 (previous section).
- **Ks falls in both, and is low confidence either way.** Under normal shrinkage the water lost near saturation is volume lost, not pores draining, so the curve shows few large draining pores: the kNN Ks drops about 6-fold for Laikipia and 25-fold for Tomejil, and the physical estimate about 4-fold. The two estimates then agree within a factor of 10 for 79 % of the Laikipia samples and 17 % of the Tomejil rings (97 % and 93 % with one density). Against the SATURO infiltrometer on 26 Laikipia samples (median 14.0 cm/h), the kNN falls from 92 % to 54 % of samples within a factor of 10, while the physical estimate stays at 92 % (96 % with one density). The low kNN values may describe the swollen matrix, while field values carry cracks and macropores; neither conversion gives a true pore-size distribution for a swelling clay.
- **The agreement is partly built in.** Both use the same formula, and their one-density values are similar wet-state densities. The independent support is the texture result, reached separately in each soil.

> **Caveat.** Under normal shrinkage the converted water content equals the porosity of the shrinking soil at every step, so the model treats the sample as still saturated at 1500 kPa. Real vertisols let air in once structural pores and cracks open, so this conversion overstates the dry end, while one fixed density understates it by ignoring shrinkage. The truth lies in between. Measured shrinkage data would pin it down: ring or clod volumes at a few water contents, a shrinkage curve (e.g. Boivin et al. 2004), or a COLE value (Grossman et al. 1968), for either soil.

References

Blake, G.R. and Hartge, K.H. (1986). Particle density. In: *Methods of Soil Analysis, Part 1*, 2nd ed., Agronomy Monograph 9, ASA–SSSA, Madison, 377–382.

Boivin, P., Garnier, P. and Tessier, D. (2004). Relationship between clay content, clay type, and shrinkage properties of soil samples. *Soil Science Society of America Journal* 68: 1145–1153.

Bronswijk, J.J.B. (1988). Modeling of water balance, cracking and subsidence of clay soils. *Journal of Hydrology* 97: 199–212.

Cornelis, W.M., Corluy, J., Medina, H., Hartmann, R., Van Meirvenne, M. and Ruiz, M.E. (2006). A simplified parametric model to describe the magnitude and geometry of soil shrinkage. *European Journal of Soil Science* 57: 258–268.

Groenevelt, P.H. and Grant, C.D. (2001). Re-evaluation of the structural properties of some British swelling soils. *European Journal of Soil Science* 52: 469–477.

Grossman, R.B., Brasher, B.R., Franzmeier, D.P. and Walker, J.L. (1968). Linear extensibility as calculated from natural-clod bulk density measurements. *Soil Science Society of America Proceedings* 32: 570–573.

Haines, W.B. (1923). The volume changes associated with variations of water content in soil. *Journal of Agricultural Science* 13: 296–310.

Hillel, D. (1998). *Environmental Soil Physics*. Academic Press, San Diego.

McGarry, D. and Malafant, K.W.J. (1987). The analysis of volume change in unconfined units of soil. *Soil Science Society of America Journal* 51: 290–297.

Mitchell, A.R. (1992). Shrinkage terminology: escape from "normalcy". *Soil Science Society of America Journal* 56: 993–994.

Ordóñez Fernández, R., González Fernández, P., Giráldez Cervera, J.V. and Perea Torres, F. (2007). Soil properties and crop yields after 21 years of direct drilling trials in southern Spain. *Soil & Tillage Research* 94: 47–54.

Peng, X. and Horn, R. (2005). Modeling soil shrinkage curve across a wide range of soil types. *Soil Science Society of America Journal* 69: 584–592.

Soil Survey Staff (2014a). *Kellogg Soil Survey Laboratory Methods Manual*. Soil Survey Investigations Report No. 42, Version 5.0. USDA Natural Resources Conservation Service, Lincoln, Nebraska.

Soil Survey Staff (2014b). *Keys to Soil Taxonomy*, 12th ed. USDA Natural Resources Conservation Service, Washington, DC.

Soil Survey Staff. National Cooperative Soil Survey Soil Characterization Database (KSSL Lab Data Mart), https://ncsslabdatamart.sc.egov.usda.gov/ (SQLite snapshot used here, queried October 2026).

Stirk, G.B. (1954). Some aspects of soil shrinkage and the effect of cracking upon water entry into the soil. *Australian Journal of Agricultural Research* 5: 279–290.

Vanderlinden, K., Pachepsky, Y.A., et al. (2017). Water retention and preferential states of soil moisture in a cultivated vertisol. *Soil Science Society of America Journal* 81: 1–9.

Vanderlinden, K., Pachepsky, Y., Pedrera-Parrilla, A., Martínez, G., Espejo-Pérez, A., Perea, F. and Giráldez, J. (2021). Water retention and field soil water states in a vertisol under long-term direct drill and conventional tillage. *European Journal of Soil Science* 72: 667–678. doi:10.1111/ejss.12967

Willard, L., Muñoz-Carpena, R., Venort, T., Gitonga, J., Maltais-Landry, G., Caylor, K. and Palm, C. (2026). A multi-domain field dataset to evaluate agricultural impacts at the edge of a Kenyan savanna. HydroShare, doi:10.4211/hs.99080b48206e4b2c82e716516d463a61 (data descriptor in review, *Scientific Data*).

Yule, D.F. and Ritchie, J.T. (1980). Soil shrinkage relationships of Texas Vertisols: I. Small cores; II. Large cores. *Soil Science Society of America Journal* 44: 1285–1291 and 1291–1295.