Algoritmo diagnóstico y de alta sanitaria para personal de salud en unidades de atención y que presentan síntomatología de infección respiratoria aguda

Toma de muestra para SARS-CoV-2 Aislamiento domiciliario estricto



- 1) Retorno a labores al cumplir 10 días del inicio de los síntomas y 72 horas sin fiebre, sin tratamiento antipirético.
- Reforzar medidas básicas de prevención y uso de EPP

- Retorno a labores hasta cumplir 72 horas sin fiebre, sin tratamiento antipirético.
- 2) Reforzar medidas básicas de prevención y uso de EPP.

No es necesaria la toma ni resultado de una segunda muestra

Referencias bibliográficas

Centers for Disease Control (CDC)

CDC Interim U.S. Guidance for Risk Assessment and Public Health Management of Healthcare Personnel with Potential Exposure in a Healthcare Setting to Patients with Coronavirus Disease (COVID-19) April 30, 2020

https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-risk-assesment-hcp.html

https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html

https://www.cdc.gov/coronavirus/2019-ncov/travelers/index.html

CDC Criteria for Return to Work for Healthcare Personnel with Confirmed or Suspected COVID-19

https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/hcp-return-work.html

NSW Health self isolation guidelines for people confirmed to have COVID-19 infection. 16 June 2020.

 $https:/\!/www.health.nsw.gov.au/Infectious/covid-19/Pages/default.aspx$

COVID-19: management of staff and exposed patients or residents in health and social care settings. 14 June 2020.

https://www.gov.uk/government/publications/covid-19-management-of-exposed-healthcare-workers-and-patients-in-hospital-settings

