

Supplementary Material to “Beyond of the joint survival and longitudinal data analysis model for binary data via two stages”

Sandro Lucena, Lizandra C. Fabio and
Jalmar M. F. Carrasco

More figures for the simulation study section in the manuscript.




Figure 1: Randomized quantile, deviance, Martingale, NMSP, and NRSP residuals (from top to bottom) for $n = 150$ and censoring 0%, 10%, 30%, and 50% (from left to right), for the model described in Scenario 2, $\max(m_i) = 2$.




Figure 2: Quantile, deviance, Martingale, NMSP, and NRSP residuals (from top to bottom) for $n = 150$ and censoring 0%, 10%, 30%, and 50% (from left to right), for the model described in Scenario 2, $\max(m_i) = 5$.




Figure 3: Randomized quantile, deviance, Martingale, NMSP, and NRSP residuals (from top to bottom) for $n = 150$ and censoring 0%, 10%, 30%, and 50% (from left to right), for the model described in Scenario 2, $\max(m_i) = 11$.




Figure 4: Randomized quantile, deviance, Martingale, NMSP and NRSP residuals (from top to bottom) for $n = 150$ and censoring 0%, 10%, 30%, and 50% (from left to right), for the model described in Scenario 3, with $\max(m_i) = 2$.




Figure 5: Randomized quantile, deviance, Martingale, NMSP and NRSP residuals (from top to bottom) for $n = 150$ and censoring 0%, 10%, 30%, and 50% (from left to right), for the model described in Scenario 3, with $\max(m_i) = 5$.



Figure 6: Randomized quantile, deviance, Martingale, NMSP and NRSP residuals (from top to bottom) for $n = 150$ and censoring 0%, 10%, 30%, and 50% (from left to right), for the model described in Scenario 3, with $\max(m_i) = 11$.



Figure 7: Randomized quantile residuals, deviance, Martingale, NMSP and NRSP (from top to bottom) for $n = 150$ and censoring 0%, 10%, 30%, and 50% (from left to right), for the model described in Scenario 4 with $\max(m_i) = 2$.



Figure 8: Randomized quantile, deviance, Martingale, NMSP, and NRSP residuals (from top to bottom) for $n = 150$ and censoring 0%, 10%, 30%, and 50% (from left to right), for the model described in Scenario 4 with $\max(m_i) = 5$.



Figure 9: Randomized quantile, deviance, Martingale, NMSP, and NRSP residuals (from top to bottom) for $n = 150$ and censoring 0%, 10%, 30%, and 50% (from left to right), for the model described in Scenario 4 with $\max(m_i) = 11$.